



Terrestrial Ecosystem Mapping of the Coastal Douglas-Fir Biogeoclimatic Zone

for:

**Mr. Bill Zinovich
Integrated Land Management Bureau
Planning Officer**

by:

**MADRONE ENVIRONMENTAL SERVICES LTD.
1081 Canada Avenue, Duncan, B.C. V9L 1V2**

June 21, 2008

Dossier 07.0359

Executive Summary

This report presents the results of Terrestrial Ecosystem Mapping (TEM) of the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone in British Columbia. An expanded legend and map legend also accompany this document for interpreting the mapping deliverables.

The CDFmm subzone occupies about 252,000 hectares of land in the Georgia Depression on southwest portions of mainland BC, the central and southern Gulf Islands and southeastern Vancouver Island. This project excludes the Lower Fraser Valley and Gulf Islands National Park Reserve (GINPR). Therefore, the resultant study area covers 212,660 hectares for which TEM mapping was completed.

Elevation boundaries for the CDFmm range from sea level to approximately 150 m, except for in the Cowichan Valley where boundary elevations were raised to 380 m. There were isolated mountains around the Cowichan area and on Saltspring Island which extended above the CDFmm boundary; therefore small amounts of the Coastal Western Hemlock very dry maritime (CWHxm) subzone have also been included in the map deliverables.

The temperature and climate in the CDFmm is mild and warm, leading to unique assemblages of vegetation and ecosystems. The subzone covers less than 0.3% of the Province, yet contains a diverse set of landscapes, vegetation, and ecosystems.

The variation of ecosystems from north to south in the CDFmm is widespread. Ecosystems include sand dune communities in the central Gulf Islands, and large river estuaries along eastern Vancouver Island, to Garry oak sites in the Cowichan Valley, Saanich/Victoria and the southern Gulf Islands. A range of anthropogenic units are also common in the CDFmm, such as cultivated fields and urban townships.

Four ecosections span the CDFmm biogeoclimatic subzone from north to south, including the Strait of Georgia, Georgia Lowland, Nanaimo Lowland, and Southern Gulf Islands. Ecosections are based on differences in physiography and macro climatic process at a sub-regional level.

Soil and terrain types present in the CDFmm today are influenced by historical glaciation events and contemporary erosion and deposition. Therefore soils are mainly derived from marine, glacio-marine, morainal, and colluvial sources.

The Biogeoclimatic Ecosystem Classification system (BEC) groups similar sets of landscapes into site series. These are mainly forested site series which can repeat across biogeoclimatic subzones. Non-forested ecosystems, wetlands and sparsely-vegetated areas also occur, such as Garry-oak meadows, estuaries and rock outcrops. Non-forested and wetland ecosystems are also derived from the BEC system naming conventions. The Wetland and Riparian Ecosystem Classification system (WREC) is used for naming various wetland sites. The typical site conditions, such as soil, terrain and climate, combined with the interaction of vegetation, animals and insects, make up the ecosystems in the CDFmm. Two-letter TEM mapcodes are used for mapping upland sites and four-letter WREC codes are used for wetland ecosystems.

Site modifiers are applied to the TEM mapcodes to further describe atypical situations, such as shallows soils and steep slopes. The succession of forested and non-forested ecosystems can be partly described by their structural features and relative age. Numerical structural stage codes, ranging from 1 to 7 are therefore added to the TEM mapcode to explain vegetation structure.

Douglas-fir dominates the tree canopy in the upland CDFmm ecosystems with lesser amounts of grand fir, arbutus, bigleaf maple and western redcedar, depending on site position, moisture regime and nutrient status. The understory of a typical forested ecosystem is characterized by the shrubs salal, Oregon-grape and oceanspray along with scattered herbs, and a continuous layer of feathermosses depending on light conditions and slope position. Tree species and understory vegetation variability across the subzone is a result of numerous factors including soil type, slope position, moisture regime, nutrient status, as well as widespread natural and anthropogenic disturbances.

The Conservation Data Centre lists thirty-five ecological communities and 218 wildlife species and plants at risk in the CDFmm. Their rare status is due to the limited range of the CDFmm in the Province and from past habitat loss arising from natural and anthropogenic disturbances.

The purpose of the project was to complete ecosystem mapping for the CDFmm study area in order to provide baseline information that can be used in support of future land-use planning initiatives.

The provincial TEM standards were followed for mapping and capturing the digital terrain and ecosystem data, using level-five survey intensity. Terrain and ecosystem mapping, field sampling and air photo interpretations were completed at a scale of 1:16,000 using the appropriate provincial standards and methods.

A stratified sampling strategy was designed to collect field data from as many types of ecosystems as possible throughout the study area. Field inspections were distributed across the study area on a wide range ecosystems and site conditions on accessible lands in the CDFmm. Field sampling resulted in over 1458 total plots, as well as 331 background plots, resulting in 1234 field checked polygons, which provided ecosystem information for final ecosystem mapping. Level-five survey intensity was accomplished with 9% polygon inspection (including background plot data).

Within the CDFmm, forest ecosystem and anthropogenic units cover 61% and 33% of the landscape, respectively, while non-forested ecosystems and wetlands encompass the remaining 6%. The most common forested unit is the Douglas-fir—Salal site series, which makes up 37% of the study area. The dominant wetland ecosystem is the Pink spirea—Sitka sedge swamp representing only 0.6% of the study area. Garry oak ecosystems occupy about 0.5% of the study area, but would increase to around 1% with the GINPR mapping incorporated.

Anthropogenic and natural disturbances have altered the forest age structure of the CDFmm ecosystems over time, mainly from urban development, agriculture, forestry, and fire. Young forests, between the ages of 40 and 80 years, occur over 71,562 ha or 45% of the CDFmm landscape. Immature (pole/sapling) forests, less than 40 years old, cover 20,026 ha or 12% of the study area. Mature forest ecosystems, between 80 and 250 years old, occupy 20,519 ha or 13% of the subzone. Old forests over 250 years old only occur on 610 ha of the study area.

Twenty-five at-risk ecosystems were mapped in the study area, twenty-one red-listed and four blue-listed communities; however rare plant field sampling was limited by survey timing.

Over 30% of the land area has been converted to urban, rural, agricultural, and industrial use in the CDFmm. However, this value probably under-estimates the actual conversion as of 2008; our mapping was based on aerial photos taken at various times in the past 15 years.

Consequently we have missed more recent disturbances. Approximately two-thirds of these anthropogenic sources represent non permeable surfaces with the remaining third as permeable surface, such as cultivated fields. Most of these patterns are associated with land use.

Overall, with the addition of ecosystem mapping results from the lower Fraser Valley and the Gulf Islands National Park Reserve, land use planning in the CDFmm can be integrated into a single database and used to guide priorities for regional outcomes for conservation and management.

ACKNOWLEDGEMENTS

This project was the result of the co-operation, organization, and dedication of many people and organizations. Funding for this project was provided by the Integrated Land Management Bureau (ILMB), under the administrative direction of Mr. Bill Zinovich (currently Land & Resource Specialist with Crown Lands and Resources, Nanaimo) and through the B.C. Ministry of Environment (MOE) via Ms. Mary Jo Hamey (Ecosystem Specialist, Nanaimo).

Islands Trust provided additional funding for higher field sampling intensity and additional map products on Saltspring Island, carried by the vision of Ms. Ardice Todosichuk (Ecosystem Protection Specialist).

BC Parks provided an additional support for even more detailed sampling and additional field and interpretive data collection and extension products on current and proposed provincial parks and ecological reserves of Saltspring Island under the auspices of Karen MacDowell and Peggy Burfield. In addition to the funding provided by the B.C. Government, this project was also sponsored by the Bulkley Valley Research Centre, a not-for-profit society based out of Smithers, BC.

Project management was led by Jane Thomson, assisted by Tania Tripp and Jackie Churchill. Bioterrain mapping was done by Wanda Miller, Michelle Trommelen, Pamela Williams, and Sonia Meili, with internal quality assurance (QA) provided by Gordon Butt and Brian Roberts, and external QA by Sid Tsang and Deepa Spaeth-Filatow (MoE). Ecosystem mapping and report writing were done by Jodie Krakowski, Helen Reid, Claudia Houwers, Chris Clement, Caroline Astley, Jackie Churchill, and Tyler Innes with external QA and feedback by Jo-Anne Stacey, Corey Erwin, Kim Everett, Ted Lea, and Carmen Cadrin (MoE). Field crews included the bioterrain and ecosystem mappers mentioned above. Andrew Neale Digital Mapping provided cascade control. Photos were monorestituted by Chartwell Consultants Ltd. Data entry was supported by Jackie Churchill, Kyle Rezansoff, and Jenny Reid. Julie Cowie provided database QA, VENUS data entry for year 2 and queries to support the results section of this report. GIS applications and final map products were completed by Jane Thomson, Dana Luxmoore, and Brett Korteling. Additional discussions with Andy MacKinnon, Sari Saunders, Todd Golumbia, regional district staff and many others provided valuable insight and clarification.

LIST OF ACRONYMS

Acronym	Description
asl	Above Sea Level
B.C.	British Columbia
BEC	Biogeoclimatic Ecosystem Classification
CDC	Conservation Data Centre, a unit within the B.C. Ministry of Environment
CDF	Coastal Douglas-Fir biogeoclimatic zone
CDFmm	Coastal Douglas-Fir biogeoclimatic zone, moist maritime subzone
CWH	Coastal Western Hemlock biogeoclimatic zone
CWHxm	Coastal Western Hemlock biogeoclimatic zone, dry maritime subzone
GEL	Georgia Lowland
GIF	Ground Inspection Form
GINPR	Gulf Islands National Park Reserve
GIS	Geographic Information System
GPS	Global Positioning System
ILMB	Integrated Land Management Bureau
MAL	Ministry of Agriculture and Lands
MOE	Ministry of Environment
MOFR	Ministry of Forests and Range
NAL	Nanaimo Lowland
QA	Quality Assurance
RISC	Resource Information Standards Committee
RIC	Resource Inventory Committee
SEI	Sensitive Ecosystem Inventory
SGI	Southern Gulf Islands
SOG	Strait of Georgia
SSI	Saltspring Island
TEM	Terrestrial Ecosystem Mapping
TRIM	Terrain Resource Information Management
WREC	Wetland and Riparian Ecosystem Classification

TABLE OF CONTENTS

1.0 INTRODUCTION	1
1.1 Location.....	2
1.2 Background	4
1.3 Project Study Area.....	6
1.4 Project Timeline	8
1.4.1 Year 1.....	8
1.4.2 Year 2.....	8
2.0 BIOPHYSICAL SETTING.....	9
2.1 Ecoregion Classification.....	9
2.2 Biogeoclimatic Ecosystem Classification	9
2.3 Geologic Setting	12
2.4 Glacial History	13
2.5 CDFmm Vegetation	15
3.0 OBJECTIVES AND DELIVERABLES	15
4.0 METHODOLOGY	16
4.1 Background Research and Data Sources	17
4.2 Pre-typing Aerial Photographs.....	18
4.3 Cascade Control	18
4.4 Monorestitution.....	19
4.5 Development of a Working Legend	19
4.6 Field Sampling	19
4.6.1 Field Sampling – Year 1.....	20
4.6.2 Field Sampling – Year 2.....	21
4.7 Final Typing and Ecosystem Labelling.....	23
4.8 Database.....	24
4.9 Map Products	25
4.10 Project Limitations	25
5.0 RESULTS AND DISCUSSION	27
5.1 Soil Landscapes.....	27
5.2 Ecosystem Overview.....	29

5.3 Subregional Location Descriptions	33
5.4 Ecosystem Representation and Condition in the Study Area	51
5.4.1 Site Series and Ecosystem Units	51
5.4.2 Structural Stage.....	60
5.4.3 Disturbance.....	62
5.4.4 Invasive species.....	65
5.4.5 Rare Elements.....	66
6.0 RECOMMENDATIONS.....	68
7.0 CONCLUSION	69
8.0 REFERENCES CITED.....	71
APPENDIX I. RED- AND BLUE-LISTED ECOSYSTEMS AND SPECIES IN THE CDFMM.....	77
APPENDIX II. AIR PHOTOS USED FOR THE PROJECT	94
APPENDIX III. THEMATIC MAPS.....	100
APPENDIX IV. TEM MAP LEGEND.....	102
APPENDIX V. TERRAIN LEGEND.....	104
APPENDIX VI. EXPANDED TEM LEGEND	106
APPENDIX VII. VASCULAR AND NON-VASCULAR PLANT SPECIES OBSERVED DURING FIELD ASSESSMENTS IN THE STUDY AREA.....	108



TERRESTRIAL ECOSYSTEM MAPPING Coastal Douglas-Fir Biogeoclimatic Zone

1.0 INTRODUCTION

This project was initiated by Integrated Land Management Branch (ILMB) and the Ministry of Agriculture and Lands (MAL) (together referred to in this report as the client) as a means to characterize the ecosystems of the Coastal Douglas-Fir moist maritime (CDFmm) biogeoclimatic subzone. This area has been identified as a high priority for comprehensive land-use planning and conservation initiatives. Accordingly the client needs accurate and comprehensive baseline information that can be used to support these initiatives.

We used Terrestrial Ecosystem Mapping (TEM) methodology to achieve this goal. TEM is a standardized set of protocols for bioterrain and ecosystem mapping supported by field data collection and geographic information systems (GIS) mapping and interpretation. Final deliverables for this project include

1. Digital TEM maps for the study area.
2. A set of hardcopy TEM maps themed by structural stage.
3. An expanded legend showing all mapped ecosystems.
4. All field data.
5. This technical report.

The report starts with an introduction of the study area location followed by a description of the biophysical setting in section two. The project objectives and deliverables are expressed in section three and the ecosystem mapping methods are explained in section four. The results of the CDFmm mapping project are presented in section five, including ecosystem and terrain descriptions for ten geographic areas progressing north to south. The results are summarized into tables and graphical representations and discussed further in sections six and seven. The map legend (Appendix IV) and expanded legend (Appendix VI) can be found appended to the end of the report along with other relevant reference material.

1.1 Location

The CDFmm occurs in southwestern B.C., covering ecosystems along the eastern coastline of Vancouver Island, the southern Gulf Islands, parts of the Sunshine Coast and a portion of the Fraser Valley. However, for this TEM project and report, the Fraser Valley and the Gulf Islands National Park Reserve were excluded.

On Vancouver Island, Deep Bay marks the northern extent of the CDFmm while Metchosin marks the southern boundary. From Deep Bay moving south, the subzone extends along the Strait of Georgia from sea level to an approximate elevation of 150 m above sea level (asl) and includes the major centres of Nanaimo, Duncan, and Victoria. The CDFmm covers or partially covers all of the Gulf Islands south of Cortes Island including: Texada, Hornby, Denman, Lasqueti, Gabriola, Valdes, Galiano, Thetis, Kuper, Saltspring, North Pender, South Pender, Mayne, Saturna, Sidney, and smaller islets in between. Across the Strait of Georgia, the CDFmm covers portions of Lund, Powell River, Sechelt, and the Fraser Valley for a total area of approximately 252,000 hectares (Figure 1).



Figure 1. Geographic extent of the CDFmm biogeoclimatic subzone.



1.2 Background

The CDFmm covers only a very small proportion of the land mass of B.C., yet it falls within one of the most densely populated areas of the province. Its mild, warm climate makes it attractive for residential, agricultural, commercial and industrial development. The cool-Mediterranean climate and ecosystem conditions that characterize the CDFmm continue south through Puget Sound, with only the northernmost tip in Canada.

The extent and condition of naturally-occurring ecosystems and wildlife have been directly impacted by anthropogenic disturbances such as logging, agriculture, invasive species, land alienation, resource extraction, altered drainage patterns, urban sprawl, and fire suppression. The continuing pressure for disturbance, combined with the restricted distribution of the subzone and its ecosystems, has resulted in mounting interest for comprehensive conservation planning.

Logging has historically been one of the primary industries in the region. With its proximity to population centres and markets, and gentle topography, the productive, accessible Douglas-fir forests of the CDFmm were the first to be harvested in BC. Around the same time logging began, the fertile lowland soils were cleared for agriculture. Growth of commercial and industrial activities contributed to land use changes (BCMELP, 1999a).

The area has experienced extensive land use changes associated with agriculture (namely forest clearing, drainage alteration, draining of wetlands) and forest harvesting (namely conversion of old-growth to second-growth forest). Now however, the area supports a rapidly growing population, drawn by the relatively mild climate and natural beauty. Today the spread of urban and suburban development is the most significant factor affecting the remaining low-elevation CDF ecosystems.

Accompanying the impacts of urban development is the loss of biodiversity due to the suppression of wildfire. Mature Douglas-fir trees are adapted to have thick bark that protects the tree from low-intensity fires which historically occurred in the CDFmm area every 100 to 300 years (Stoffels, 2000).

Fires have been largely eliminated from the landscape over the past century and as a result many of the fire-adapted indigenous plant species and communities are being replaced by an influx of non-native invasive plants which thrive in this modified disturbance regime (i.e., fire suppression).

Ecosystems throughout the CDFmm are currently listed as critically imperiled in a global context by the B.C. Conservation Data Centre (CDC). Currently, 35 plant communities are provincially red or blue-listed in the CDFmm (B.C. C.D.C. 2008) (Appendix I). Most forested site series and wetlands are on these lists, including 28 red-listed forested communities, herbaceous meadows, woodlands, and wetlands and 7 blue-listed herbaceous and forested wetlands. Of these 35 communities, 17 are listed as globally “imperiled” (S2) or “critically imperiled” (S1), while 3 are secure or apparently secure (where their occurrence in B.C. represents the northernmost extent of a larger range), and 11 are not yet ranked.

A similar search of the database for red and blue-listed species occurring in the CDFmm yielded 218 records, including unique populations and subspecies or varieties (Appendix I). Several marine species, including fish, mollusks, and marine mammals, were listed since their range falls within or overlaps the CDFmm. Of the 218, 30 occurrences were endemic to B.C., including six populations of genetically unique stickleback. There were 101 blue-listed taxonomic units and 117 red-listed units, including the six stickleback populations. Of the total, 16 were globally listed S1 or S2, with the vast majority having secure or apparently secure populations in terms of their global occurrence. Several subspecies or varieties were lacking in information on that specific taxonomic unit, but the species overall were ranked as globally secure.

The CDFmm portion of the lower mainland and southern Vancouver Island is unique in the province in that a relatively large portion of the landbase is privately owned. This contributes to the complexity of developing an overall strategy for managing the landbase.

1.3 Project Study Area

The study area for this project includes the entire CDFmm excluding the Gulf Islands falling within the Gulf Islands National Park Reserve (GINPR) and approximately 42,000 hectares within the Fraser Valley.

The islands within the GINPR (North and South Pender Islands, Saturna Island, Prevost Island, Mayne Island, Sidney Island and surrounding islets) were not included in this project because they were recently TEM mapped by Parks Canada and will complement this project. The CDFmm in the Fraser Valley was considered a lower priority and is already heavily modified, so was not included in the study area. The total area TEM mapped by Madrone totals 212,660 hectares (Figure 2).

In some cases, particularly on warm aspects, the CDFmm extends above 150 m. The adjacent biogeoclimatic unit above the CDFmm on eastern Vancouver Island and the Gulf Islands is the eastern variant of the very dry maritime subzone of the Coastal Western Hemlock zone (CWHxm1). The western (slightly cooler and wetter) variant, CWHxm2, is directly above the CWHxm1 in the study area. Within the study area CWHxm1 and CWHxm2 were mapped on peaks of Saltspring Island and Cowichan Valley (Figure 2).



Figure 2. Project study area and biogeoclimatic subzones.



1.4 Project Timeline

The CDFmm TEM project was initiated by ILMB (the client) in fall 2006. The ILMB was the primary sponsor of the project; however, co-sponsors included Islands Trust Fund and The Bulkley Valley Centre. In-kind contributions were made by Ministry of Environment and B.C. Parks. The project was divided into two years of work, described in more detail below in sections 1.4.1 and 1.4.2. Year 1 of the project was scheduled from October 2006 to March 31st 2007. Year 2 commenced in October 2007 and was completed June 2008.

1.4.1 Year 1

From the onset of this project it was determined by all sponsoring parties that a CDFmm TEM pilot project should be completed prior to mapping the entire CDFmm study area. Saltspring Island (SSI) was chosen as the pilot project study area and was allocated priority for mapping, field work, reporting, and map production. The deliverables for year one included a preliminary TEM map and technical report for SSI, 15% field verification of Crown land TEM polygons on SSI, 10% field verification of private land TEM polygons on SSI, preliminary bioterrain and ecosystem pretyping, and a summary report for the remainder of the study area, excluding the Saanich Peninsula and Greater Victoria where current air photos were unavailable. A total of \$175,000 was funded by ILMB and \$60,000 by Islands Trust for year one of the project.

1.4.2 Year 2

Year 2 of this project began in late October 2007. The project continued with 7% field verification of TEM polygons on Crown lands and 17% verification of TEM polygons on privately owned land within Islands Trust jurisdiction. Madrone field staff completed over 930 field plots throughout the entire study area (excluding SSI) between October 2007 and February 2008. Madrone also contacted private landowners for permission to access areas on the Gulf Islands. The deliverables for year two included final Resource Information Standards Committee (RISC) standard digital TEM deliverables for the entire study area, a set of TEM maps and this accompanying technical report. A total of \$285,000 was funded by ILMB, \$160,000 by the Bulkley Valley Centre and \$26,000 by the Islands Trust.

2.0 BIOPHYSICAL SETTING

The ecosystems mapped in the CDFmm include forested units and non-forested units, wetlands, and sparsely-vegetated ecosystems. There is a great diversity and variability of ecosystems in the CDFmm subzone as a result of the interactions between climate, terrain, and vegetation. Two classification schemes in B.C. help to place the ecosystems into their respective vegetative, terrain and climatic context. The Ecoregion Classification and the Biogeoclimatic Ecosystem Classification (BEC) systems offer a framework for describing the variation of vegetation, climate and topography in the CDFmm subzone.

2.1 Ecoregion Classification

The CDFmm is situated within the Georgia Depression ecoprovince, which lies between the Vancouver Island Mountains and Coast Mountains. Within this ecoprovince, the CDFmm covers four ecosections: the Strait of Georgia (SOG) ecosection and portions of the Georgia Lowland (GEL), Nanaimo Lowland (NAL) and Southern Gulf Islands (SGI) ecosections. The climate in this sub-region is cool mesothermal with long dry, warm summers and relatively mild, wet winters during which temperatures seldom fall significantly below freezing.

Localized meso- and micro-scale effects create some fine-scale variability relative to the regional mean values. Aspect has a pronounced effect on growing season duration and moisture availability. Perhaps the most important factor affecting terrestrial ecology is the summer moisture deficit, which arises from the relatively dry summers from July through September.

2.2 Biogeoclimatic Ecosystem Classification

The BEC system is a framework that groups similar sets of landscapes into a site classification. Sites are classified on their potential to produce similar vegetation communities within similar environmental site conditions. Site series are representative ecosystems in each biogeoclimatic subzone in the Province. Site series are specific to a subzone and primarily correspond to forested ecosystems that repeat across each biogeoclimatic subzone.

The typical site conditions, such as soil, terrain and climate combined with the interaction of vegetation, animals and insects make up the ecosystems in the CDFmm (Green & Klinka, 1994). Additional non-forested ecosystems, such as wetlands, are also based on the BEC system (Mackenzie & Moran, 2004).

Although the focus of this mapping exercise is the CDFmm, we also mapped adjacent areas in the CWHxm subzone, which lies above and along the western edge of the CDFmm. In the project area, the CWHxm was mapped on the peaks of Saltspring Island and Cowichan Valley however the full extent of the CWHxm was not mapped (Figure 2).

The CDFmm is in the rainshadow of the Vancouver Island Mountains as well as the Olympic Mountains in Washington. It therefore receives less precipitation than adjacent biogeoclimatic units. The Pacific Ocean moderates temperatures throughout the area, creating long growing seasons and pronounced growing season water deficits on zonal and drier sites. The CDFmm represents the mildest climate in Canada.

The adjacent CWH zone is characterized by a cooler, wetter climate during the growing season. The CWHxm (dry maritime) subzone, adjacent to the CDFmm in the project area, is further subdivided into eastern and western variants (CWHxm1 and CWHxm2, respectively), with the CWHxm2 slightly wetter and cooler than the CWHxm1, and mapped immediately above it at the highest points on Saltspring Island (Mount Sullivan and Hope Hill), above approximately 600 m.

Table 1 includes some representative climate station data from Environment Canada stations throughout the CDFmm to provide an indication of the variability and mean values for temperature and precipitation.

Table 1. Climate Normal Data (1971-2000) for Stations Throughout the CDFmm.

Station	Parameter	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Cortes Island	Daily mean (°C)	3.4	4.6	6.5	9.1	12.7	15	18	18	15	10.1	5.9	3.7	10.2
50° 4.800' N	Std. Dev. (°C)	1.7	1.4	1.3	1.2	1.1	1.1	1.1	1.1	1.2	0.8	1.5	1.6	1.3
125° 1.800' W	Daily max (°C)	5.4	7	9.4	13	16.6	19	22	22	18	12.7	8	5.5	13.2
06.10 m	Daily min (°C)	1.4	2.2	3.5	5.6	8.8	12	14	14	11	7.4	3.8	1.8	7.0
Station ID	Rainfall (mm)	148	121	111	89	70.2	65	51	59	72	150	207	168	1308.8
1021950	Snowfall (cm)	24.8	12.2	5.5	2	0	0	0	0	0	0.6	4.9	19.9	69.9
Delta Ladner South	Daily mean (°C)	2.8	4.5	6.6	8.9	12.2	15	17	17	14	9.5	5.4	3.1	9.6
49° 4.200' N	Std. Dev. (°C)	1.8	1.5	1.1	1.1	0.7	1	0.8	1	1	1	1.9	1.9	0.5
123° 4.800' W	Daily max (°C)	5.8	7.9	10.5	14	17.2	20	23	22	19	13.8	8.7	6	13.9
02.00 m	Daily min (°C)	-0.3	1	2.6	4.2	7.2	9.9	11	11	8.6	5	2	0.2	5.3
Station ID	Rainfall (mm)	104	102	92.2	70	56.8	45	34	33	57	93.2	155	134	976.5
1102417	Snowfall (cm)	10.6	7.9	0.7	0	0	0	0	0	0	0.1	2.7	9.5	31.6
Gabriola Island	Daily mean (°C)	3.7	4.3	6.1	8.5	11.7	14	17	17	14	9.3	5.6	3.3	9.5
49° 9.000' N	Std. Dev. (°C)	1.3	1.7	0.9	0.8	1.3	1	1.1	0.8	1.1	0.9	1.7	1.5	0.7
123° 43.800' W	Daily max (°C)	6.6	7.7	10	13	16.6	19	22	22	19	13.4	8.8	6	13.8
46.00 m	Daily min (°C)	0.8	0.8	2.2	4	6.6	9.6	11	11	8.4	5.1	2.4	0.6	5.3
Station ID	Rainfall (mm)	116	96.8	85.1	57	44.9	41	26	28	39	80.9	143	127	884.3
1023042	Snowfall (cm)	13.7	9.1	1.8	0	0	0	0	0	0	0.3	3.9	10.9	39.7
Nanaimo A	Daily mean (°C)	2.7	4.2	6.1	8.8	12.3	15	18	18	15	9.7	5.4	2.9	9.8
49° 3.000' N	Std. Dev. (°C)	1.6	1.5	1.1	1	1.2	1.1	1.1	1.1	1.2	0.8	1.6	1.7	0.6
123° 52.200' W	Daily max (°C)	6.2	8.2	10.9	14	17.8	21	24	24	21	14.6	9.1	6.1	14.7
28.40 m	Daily min (°C)	-0.8	0	1.3	3.4	6.7	9.7	12	12	8.6	4.8	1.5	-0.4	4.9
Station ID	Rainfall (mm)	142	123	106	63	49.9	45	26	32	39	97	191	166	1078
1025370	Snowfall (cm)	27.2	16.3	5.6	0.2	0	0	0	0	0	1.1	7.8	22.8	80.9
Saanichton CDA	Daily mean (°C)	4	5.1	6.7	9	12	15	17	17	14	10.2	6.4	4.3	10
48° 37.200' N	Std. Dev. (°C)	1.6	1.4	1.2	1	1	1	0.9	1	1	0.8	1.5	1.5	0.5
123° 25.200' W	Daily max (°C)	6.7	8.1	10.2	13	16.4	19	22	22	19	13.8	9.2	6.8	13.8
61.00 m	Daily min (°C)	1.3	2.1	3.2	4.9	7.5	9.9	12	12	9.8	6.6	3.6	1.8	6.2
Station ID	Rainfall (mm)	127	98.7	75.5	47	38.9	35	21	28	32	79.3	154	145	879.8
1016940	Snowfall (cm)	10.9	4.5	1.2	0	0	0	0	0	0	0.3	1.2	8.2	26.2
Victoria Airport	Daily mean (°C)	3.8	4.9	6.4	8.8	11.8	14	16	16	14	9.8	6.1	4	9.7
48° 39.000' N	Std. Dev. (°C)	1.6	1.5	1.1	1	1.1	0.9	0.7	0.9	1	0.7	1.5	1.5	0.6
123° 25.800' W	Daily max (°C)	6.9	8.4	10.5	13	16.6	19	22	22	19	14.2	9.5	6.9	14.1
19.20 m	Daily min (°C)	0.7	1.4	2.3	4.1	6.9	9.3	11	11	8.4	5.3	2.7	1	5.3
Station ID	Rainfall (mm)	122	98.8	75.8	45	36.5	32	20	24	30	75.6	144	138	841.4
1018620	Snowfall (cm)	15.2	9	2.4	0	0	0	0	0	0	0.2	3.3	13.8	43.8
Qualicum River	Daily mean (°C)	3	4	5.6	8.3	11.7	15	17	17	13	9.1	5.4	3.3	9.3
49° 23.400' N	Std. Dev. (°C)	1.6	1.5	1	0.9	1	0.9	0.9	1	1	0.7	1.6	1.6	0.5
124° 37.200' W	Daily max (°C)	5.7	7.3	9.5	13	16.4	19	22	22	18	13	8.3	5.8	13.3
07.60 m	Daily min (°C)	0.3	0.8	1.6	3.9	7	9.9	12	11	8.4	5.2	2.4	0.7	5.2
Station ID	Rainfall (mm)	177	151	115	69	50.9	47	28	37	52	137	214	186	1264
1026565	Snowfall (cm)	16.2	12.4	3.5	0.2	0	0	0	0	0	0.4	3.7	13.8	50



2.3 Geologic Setting

The CDFmm straddles the Georgia Strait, occupying a broad physiographic unit called the Georgia Depression by Holland (1976). Within this unit, Holland further identified three areas, namely the Nanaimo Lowland, Georgia Lowland, and Fraser Lowland.

The Nanaimo Lowland is located on the western side of the Georgia Strait, and includes southeast Vancouver Island and the Gulf Islands. The area is composed of flat to gently tilted sedimentary rocks of the Nanaimo Group. The northeast-southwest orientation of the Gulf Islands reflects the southeast dip of these rocks; the ridges generally consist of harder sandstones, and the valleys in between are typically mudstones or shales.

The Georgia Lowland forms a narrow coastal strip on the eastern side of the Georgia Depression, including the Sunshine Coast up to Powell River. It is a diverse landscape shaped by complex tectonic activity, bedrock structure, glacial events and post-glacial deposits.

A third subdivision of the Georgia Depression is the Fraser Lowland which includes the Fraser Valley.

Bedrock outcrops mapped by the B.C. Geological Survey¹ and in part, observed in the field inspections include:

- Upper Cretaceous Nanaimo Group: undivided sedimentary rocks occurring along the eastern coast of Vancouver Island and the majority of the Gulf Islands.
- Middle to Upper Triassic Karmutsen Formation: basaltic rock occurring predominantly on Texada Island, Lasqueti Island, and North and South Thormanby Islands, and in pockets along the east coast of Vancouver Island and other lesser islands.
- Late to Early Cretaceous unnamed dioritic intrusive rocks along the west coast of the mainland in the Sunshine Coast and Powell River areas.

¹The Map Place – BCGS Geology Map: <http://webmap.em.gov.bc.ca/mapplace/minpot/bcgs>. Accessed March 25, 2008.

- Middle to Upper Triassic Vancouver Group: limestone, marble and calcareous rocks of the Quatsino Formation occurring in pockets on Texada Island.
- Middle to Upper Devonian Mount Sicker Group: Nitnat and MacLaughlin Ridge Formations – calc-alkaline volcanics and volcaniclastics occurring in pockets in the Cowichan Valley area and on Saltspring Island.
- Late Devonian Saltspring Plutonic Suite: granodioritic intrusive rocks occurring in pockets on southern Saltspring Island and in the Greater Victoria area.
- Late Triassic Mount Hall Gabbro: gabbroic to dioritic intrusives occurring in pockets on southern Saltspring Island, in the Greater Victoria area, and on lesser islands nearby.

2.4 Glacial History

The Georgia Depression experienced multiple episodes of glaciation and deglaciation during the Pleistocene Epoch (beginning 1.8 million years ago and ending 10,000 years ago) of the Quaternary Period (Bichler *et al.*, 2002). The most recent glacial cycle was Fraser Glaciation (Clague, 1994). The early phases of this glaciation were characterized by the growth of alpine glaciers on both sides of the Georgia Depression, as well as the expansion of the Cordilleran Ice Sheet, covering much of western Canada. At the peak Glaciation, the Cordilleran Ice Sheet overrode Georgia Strait and enveloped Vancouver Island. Quaternary sediments associated with this glaciation cover the majority of the study area.

Approximately 15,000 years before present, the climate began to warm and the ice sheets slowly melted and retreated. The retreat of Fraser Glaciation ice was essentially complete by about 10,000 years, which coincides with the beginning of the Holocene Epoch. Sediments deposited during this non-glacial period are typically formed by modern processes still observable in the study area today.

Sea levels fluctuated above and below present levels during the Quaternary Period due to subsidence of the land surface related to the weight of overlying ice, isostatic uplift associated with the removal of the ice, global sea level changes, and tectonism. The land mass of Vancouver Island has risen (relative to sea level) by 50 m – 300 m in elevation due to isostatic rebound since deglaciation. Within the study area, this uplift has resulted in marine and glaciomarine deposits at elevations up to 180 m asl.

The majority of the study area is covered by glacial and related sediment deposits laid down during a period of high sea level between 19,000 and 13,000 years ago (Bichler *et al.*, 2002). These glacial-related sediments originated largely from marine, deltaic, and fluvial environments near the margins of Pleistocene glaciers (Clague, 1994).

Several deposits have been characterized. Quadra sands are outwash sediments that typically consist of cross-stratified, well-sorted glaciofluvial sands. These sediments were deposited in front of advancing glaciers, and can now be found up to 100 m asl. Vashon Drift consists of silty sandy till and sandy gravelly glaciofluvial and glaciolacustrine sediments initially deposited along the margins of overriding ice lobes. As the study area started to become ice-free some 13,000 years ago, thick deposits of the Capilano Sediments marked the ice retreat with glaciomarine and marine sediments deposited on the seafloor. These deposits are now seen as raised deltas and intertidal beach sediments. The Capilano Sediments also consist of glaciofluvial deposits and can be found up to 180 m asl in the study area.

In a general cross-section from the coast to inland uplands, low-lying areas adjacent to shorelines commonly consist of glaciomarine and marine deposits, and may also include finer textured deposits possibly associated with impounded lakes or locally protected marine embayments. Glaciofluvial sands and gravels are also common throughout low-lying areas, river valleys, and where streams exit confined gully systems forming fans. On middle and upper slopes, surficial material commonly includes silty, sandy diamictic till deposits, with colluvial deposits and exposed bedrock outcrops.

2.5 CDFmm Vegetation

The coastal variety of Douglas-fir is the principal tree species in the CDF. Douglas-fir forests originally blanketed most of the area, with the exceptions of scattered bedrock outcrops, wetlands, meadows and a small number of non-forested sites. This productive forest has provided the foundation for industrial forestry.

Natural and anthropogenic disturbances have changed the vegetation structure and stand age of upland ecosystems in the study area. When the tree canopy is removed in this subzone a characteristic succession of pioneer vegetation establishes.

Other tree species found in the CDFmm include redcedar, grand fir, bigleaf maple, arbutus, and Garry oak. Hundreds of plant species occur in the subzone, including a variety of shrubs, herbs, mosses and lichens. A more detailed description of ecosystems in the CDF are provided later in this report and in the expanded legend.

3.0 OBJECTIVES AND DELIVERABLES

The purpose of this project was to complete ecosystem mapping for the CDFmm study area. The mapping was to be completed at a scale of 1:16,000 using Resource Information Standards Committee (RISC) standards at modified survey intensity 5, following the *Standard for Terrestrial Ecosystem Mapping in British Columbia* (RISC, 1998b). In addition to the standard attributes collected for TEM, ILMB and MOE requested that further TEM parameters be collected, including continuous riparian polygons, disturbance history, separating smaller pure polygons containing sensitive ecosystems, and completion of conservation evaluation forms during collection of ground inspections and full plots.

The final project objective was to produce the following deliverables:

- A complete set of organized, boxed, and labeled hardcopy air photos with field plot locations identified and final ecosystem and terrain polygons delineated.
- A complete ecosystem and terrain attribute database for all mapped polygons.

- A field sampling program following a modification of the Survey Intensity Level 5 using a combination of Ecosystem Field Forms (FS882), Ground Inspection Forms (GIFs, FS212) and visual checks in a ratio of 5:20:75, With sampling on Crown land over a minimum 7% of polygons and 15% on Islands Trust lands.
- A seamless GIS layer of the final mapped ecosystem polygons merged with the GINPR TEM polygons (if available) in ArcInfo coverage format (RISC, 2000, 2002, 2003).
- A GIS layer of the field plot locations in ArcInfo coverage format.
- A complete error-free VENUS database containing all field data.
- A GIS layer and associated database containing all background plots from relevant ecosystem mapping projects that were used to support mapping.
- A database of Conservation Data Centre (CDC) Conservation Evaluation forms (BCMOE, 2006) completed at each full and ground inspection plot.
- A final expanded legend and report summarizing methods, results, interpretations, conservation recommendations and a detailed description of each mapped ecosystem.
- A TEM map legend.
- Final thematic maps depicting the spatial distribution of old, mature, young and immature forests, Garry oak ecosystems, non-forested areas, riparian/wetland areas, and anthropogenic or urban areas.

4.0 METHODOLOGY

Mapping was completed according to the methodology outlined in the *Standard for Terrestrial Ecosystem Mapping in British Columbia* (RISC, 1998b). The TEM method utilizes the hierarchical biogeoclimatic ecosystem classification system approach. Ecoregions and biogeoclimatic zones represent the broad scale regional and climatic landscape units. Ecoregions are further subdivided into ecosections; biogeoclimatic zones are subdivided into subzones, which are themselves divided into variants.

Polygons are then delineated based on differences in bioterrain and ecological features. Bioterrain mapping identifies terrain features and landforms, while ecological mapping identifies site series (plant communities), site modifiers (e.g., aspect, slope) and structural stage. These bioterrain and ecological features are determined through digitized image or aerial photograph interpretation and verified by field sampling.

The following sections describe in further detail how these methods were applied to the CDFmm TEM project. As this project was carried out in a condensed time frame, most activities were concurrent, rather than in the order listed.

4.1 Background Research and Data Sources

In year one of the project, a mosaic of monochrome and colour aerial photographs of varying scales and years were acquired from ILMB for the majority of the project area. These photos were then prepared for interpretation. Preparation included boxing, north arrows and delineation of the study area boundary. In some areas there were gaps in the photo coverage and additional photos were ordered. In some areas where photos were not available or outdated, photos were acquired in Year 2. The majority of the photos were flown by the Province with the remainder flown by McElhanney Consulting Services Ltd. and Intergrated Mapping Technologies Inc. A full list of the aerial photographs and roll information is available in Appendix II.

Background materials on the geomorphology, vegetation ecology, land management, sensitive ecosystem inventory, rare element occurrences, Garry oak communities, aggregate resources, soils, and rare species were researched. Where permission was granted, numerous projects including sensitive ecosystem inventory and TEM that was previously carried out in portions of the CDFmm on private, Federal and Provincial Crown land were collected to provide additional information and data (BCMELP, 1999, BCMELP, 2001; Dunster & Booth, 2001; Madrone, 2002; Madrone, 2003; Madrone, 2005, Madrone 2008b).

Digital GIS datasets were provided by ILMB and Islands Trust. Access to Terrain Resource Inventory Mapping (TRIM) data for the entire project area, as well as orthophoto and satellite imagery, parks and protected areas, digital elevation models, hillshades, sensitive ecosystems inventory (SEI), and land tenure, was provided by ILMB.

The Islands Trust provided GIS datasets containing administrative boundaries, land tenure and prior Islands Trust ecosystem mapping. Madrone also obtained ecoregion, ecosection, and biogeoclimatic boundaries from the ILMB land and resources data warehouse.

4.2 Pre-typing Aerial Photographs

Photos were pre-typed for bioterrain (surficial geology material, texture, thickness, expression, processes, modifiers, and drainage) following the *Guidelines and Standards to Terrain Mapping in British Columbia* (RIC, 1996), the *Terrain Classification System for British Columbia* (Howes and Kenk, 1997) and the *Standard for Terrestrial Ecosystem Mapping in British Columbia* (RISC, 1998b). Bioterrain pre-typing was revised based on internal QA. Completed pre-typed photos of representative areas were sent to external quality assurance by Ms. D. Spaeth-Filatow, P. Geo., MoE, to ensure accuracy and consistency across the study area.

After adjustments were made to incorporate the QA, these sections were then pre-typed for ecosystem mapping, which consisted of subdividing bioterrain polygons where multiple discrete ecosystem types, aspects, and/or stages occurred, and adjusting the existing bioterrain polygons to correspond with the ecosystem polygons (RISC, 1998b). Pre-typed linework for SaltSpring Island was then sent to C. Cadrin and T. Lea, MoE, for QA.

4.3 Cascade Control

During year one of the project, all of the aerial photographs were cascade-controlled by Andrew Neale Digital Mapping. This procedure entails transferring geospatially referenced control points from high level TRIM photography into each typed photo. Control points situated close to the four corners of each typed photo are critical to ensure the accuracy of the monorestitution setup and transfer. The cascade control procedure followed the provincial standards (RISC, 2000, 2004).

The area covered by this project encompassed over 1400 air photos, requiring cascade control for approximately 700. Each section of the project area was cascade-controlled prior to pre-typing. In Year 2, new photos were received for the Saanich Peninsula and Thetis / Kuper Islands Trust Area and were subsequently cascade-controlled prior to pre-typing.

4.4 Monorestitution

Data capture of the pre-typed, cascade controlled aerial photographs was completed by Chartwell Consultants Ltd. following monorestitution guidelines (RISC, 2000, 2004).

In Year 1, Saltspring Island, Powell River, Texada Island, Lasqueti Island and Sechelt were monorestituted. In Year 2, the remainder of the study area was monorestituted.

Monorestitution following field work maximizes efficiency, after pre-typed line work was finalized through field verification and QA. This sequence minimizes the need for repeated digital editions and extensive cross-tracking of edits on photos and databases as polygons are adjusted. For this project, monorestitution was almost always completed prior to field work (excluding the Cowichan Valley and Saanich Peninsula) to provide field crews with maps to target field sampling based on polygon distribution and representation. This approach resulted in extensive post-field work and post-QA linework edits, which significantly added to the workflow for all project components.

4.5 Development of a Working Legend

Bioterrain attributes such as surficial material type and surface expression, soil drainage, soil depth, slope, aspect, and slope position were used to guide the development of an ecosystem working legend. Biogeoclimatic subzone profiles were created using the terrain features that typically correspond with the development of predictable ecosystem types. Preliminary field data and existing background information were integrated to develop a working legend prior to beginning field work (RISC, 1998b: Table 6.1). This legend was refined following ground truthing where modifications warranted, and ultimately integrated into a series of landscape profiles for the study area (See Figures 3 and 4 in Section 5.2)

4.6 Field Sampling

Field sampling methods followed provincial standards for TEM data collection (RISC, 1998a, 1998b). Sampling efforts were conducted by two teams, each led by an ecosystem and a bioterrain specialist. Polygons were sampled using three types of plots: full ecosystem plots with site, soil, vegetation and wildlife descriptions (FS882); ground inspection plots (FS212); and visual inspections (also recorded on FS212).

Spatial coordinates of each plot were recorded by a handheld GPS unit in addition to pin-pricks on the aerial photographs. A digital photograph was taken at each full and ground plot, as well as most of the visual plots.

Conservation evaluation forms (BCMOE, 2006) were filled out for all full and ground plots in the project area, except in protected and disturbed areas. Parameters include rare elements occurrences, disturbance sources, resilience, fragmentation, and presence of invasive species. These evaluations are used to prioritize areas for conservation efforts.

A stratified sampling strategy was designed to collect field data from as many types of ecosystems as possible throughout the study area. Full and ground plots were concentrated on sites likely to support zonal ecosystems, infrequently-occurring ecosystems, rare species, mature stands, and typical ecosystems representing all site series and positions on the edaphic grid mapped. Additional plots were included to confirm structural stages, site series, proposed new or non-correlated ecosystem types, and ecosystem types that were difficult to identify from the air photos (e.g., fluctuating water table sites and some disturbed sites).

Visual inspections were used to confirm site series, structural stages, identification of some disturbed and anthropogenic areas, and areas not accessible for more detailed plots because of access limitations. Disturbance features and riparian zones were noted, as well as other features of interest.

In order to clarify the boundary within transitional zone between the CDFmm and CWHxm subzones, elevational transects from high to low elevation were completed in accessible areas. Cool and warm aspects were assessed to identify the variability associated with topography in the study area. Pre-typed photos, draft maps, and plot data for areas where data potentially supported localized biogeoclimatic unit (i.e., zone, subzone, or variant) delineation adjustments were sent to the Regional Ecologist for approval of proposed changes.

4.6.1 Field Sampling – Year 1

Field work on Saltspring Island was conducted in January, March and April of 2007. Most sites were visited by vehicle, on foot, and by water taxi.

We aimed to sample 25% of polygons across Saltspring Island, with 50% polygon visitation rates in provincial parks and candidate areas identified by BC Parks at a ratio of 5:20:75 full:ground:visual sampling intensity. A total of 523 plots were assessed in the field: 25 full plots, 114 ground inspection plots, and 384 visual inspections. Sampling included public and private lands with landowner consent, supported by the Islands Trust.

4.6.2 Field Sampling – Year 2

Field work for the remainder of the study area was carried out between October 2007 and March 2008. For all areas sampled, most plots were visited by vehicle, on foot, or by water taxi. A draft field sampling plan was prepared in early September 2007. Following approval, Madrone field teams and provincial correlators completed plots on Maple Mountain and Mt. Richards in the Cowichan Valley on October 16 and 17, 2007. Field correlations ensured consistent field data collection for all field crews in accordance with the RISC (1998a) standards.

In early October 2007, a field crew of six Madrone staff completed 179 plots within Powell River, Texada Island and Savary Island. Provincial correlators Corey Erwin, Kim Everett and Sid Tsang conducted field work QA and reported on their findings to Madrone crews, who made the requisite adjustments to field sampling and data collection for subsequent plots.

Between October and December 2007, a field crew of two sampled 180 plots in the Cowichan Valley. This region contains the highest elevations in the study area. Results indicated a local revision of the CDFmm/CWHxm boundary was necessary. In early March 2008, Regional Ecologist Andy Mackinnon accompanied Madrone crews in the field and recommended that the CDFmm/CWHxm boundary should be moved from its present 150 m to 380 m.

A total of 62 plots were collected over four days on Denman and Hornby Islands in mid-December, 2007 by one crew of two. Field work assessing 76 plots across Sechelt and Lasqueti Island was conducted in mid-December, 2007. Sechelt field work was completed by one team of two; Lasqueti Island fieldwork was completed by local vegetation ecologist Doug Hopwood and one Madrone bioterrain team member.

A total of 215 plots were collected in the Bowser, Qualicum, Parksville, Nanaimo, and Ladysmith areas between November 2007 and February 2008. Kim Everett and Sid Tsang conducted field work QA in the Qualicum area.

The remainder of the Gulf Islands field work (Gabriola Island, Valdes Island, Thetis Island, and Galiano Island) was completed in early 2008, assessing 136 plots.

The Saanich Peninsula and Capital Regional District were sampled in February 2008. A total of 87 plots were collected in this area.

A total of 76 full plots, 399 ground inspection plots and 983 visual checks were completed by the Madrone team in 2007/8 (Table 2). Sampling included public and private lands (within the Islands Trust). Additional visual plot data were employed from existing projects (TEM, SEI, ecological assessments, and conservation evaluations) to fulfill the sampling intensity required to meet the project objectives while minimizing duplication of previous efforts. Plot data was entered into the provincial database VENUS 5.1 and checked for errors.

Table 2. Plot Summary: Sampling Intensity by General Location

Region	Full plots	Ground plots	Visual plots	Total plots	Conservation evaluation forms
Powell River	4	14	48	66	18
Texada Island	6	32	75	113	38
Sechelt and Lasqueti Island	4	25	47	76	17
Denman and Hornby Islands	4	16	42	62	20
Parksville and Qualicum	8	47	98	153	53
Gabriola Island	2	9	35	46	5
Valdes Island	1	1	9	11	2
Thetis Island	1	6	15	22	7
Nanaimo and Ladysmith	2	30	30	62	32
Cowichan Valley	8	55	117	180	58
Saltspring Island	25	114	384	523	0
Galiano Island	3	11	43	57	14
Saanich Peninsula and Greater Victoria	8	39	40	87	38
Totals	76	399	983	1458	302

4.7 Final Typing and Ecosystem Labelling

Following field work, preliminary bioterrain and ecosystem line work was adjusted. Ten percent of the photos from each section of the study area were then sent to Sid Tsang for final bioterrain mapping QA to ensure the mapping met the RISC standard.

Draft hard-copy maps were then created displaying final polygons. Each was assigned a unique number labeled within each polygon centroid. These maps were then used in conjunction with the aerial photographs to complete the ecosystem mapping. The ecosystem mappers reviewed the aerial photographs and applied ecosystem labels integrating all available data sources (e.g., field plots, background research, and bioterrain labels).

Each ecosystem label is coded using a standard format to describe the vegetation, site conditions, structural stage, stand composition, and disturbance. Ecosystems are assigned up to three ecosystem codes, each comprising a quantified proportion of the polygon area using deciles.

Ecosystems, with the exception of wetlands are coded using two upper case letters indicated in the provincially correlated TEM code list², including sparsely vegetated, non-vegetated, and anthropogenic units. Wetlands are classified following the Wetland and Riparian Ecosystem Classification system (WREC) which is based on the BEC system (Mackenzie & Moran, 2004). They are assigned a four character code, and are considered a site series number in the database. The first two characters are letters (the first letter uppercase, and the second lower) while the second two characters are numbers. Where an ecosystem does not correspond to a designated unit, new codes were proposed to the Provincial correlators based on the field data collected.

Each ecosystem code has assumed (typical) site condition modifiers and may also have atypical site modifiers coded using individual lower case letters. The site series and/or site modifier is followed by the numerical structural stage designation (1 through 7). Structural stages describe the dominant seral stage for the ecosystem unit (RISC, 1998a).

² <http://www.env.gov.bc.ca/ecology/tem/list.html>

An optional structural stage modifier (a single lower case letter) further characterizes the structural stages. A stand composition modifier (single upper case letter) may follow the structural stage and/or structural stage modifier to differentiate forest stands based on proportions of coniferous and broadleaf canopy. For this project, the coniferous modifier was assumed for all forested stands with a stand composition modifier applied when the situation was otherwise.

Disturbance type and subtypes were also mapped when apparent. Logging and land clearing represented the primary type of disturbance mapped in the study area.

Portions of the study area were submitted to Corey Erwin for the initial ecosystem mapping QA. Feedback was addressed by the ecologist that mapped the area and reviewed by the other mappers to ensure consistency throughout the study area. An internal QA process was established between Madrone ecosystem mappers, featuring continual communication, and detailed review of map work.

4.8 Database

As areas were completed and finalized, ecosystem and bioterrain map codes were entered into the standard Excel database (RISC, 2002). Once an area was complete, the database was imported into ArcGIS 9.2 and joined to the ecosystem polygons based on the unique polygon identifier. This preliminary join would highlight correspondence errors in the database.

The revised database had a one-to-one relationship between the spatial and non-spatial datasets. Mapsheet, ecosection and subzone fields were populated. The final database was then imported back into Excel and subjected to further QA using the TEM Data Capture application (DC Tool). This tool highlights errors where there are non-standard ecosystems and bioterrain attributes. The final database was confirmed as error-free by the DC Tool utility, with the exception of non-standard or updated codes (e.g., wetlands) that the tool does not recognize.

4.9 Map Products

Following finalization of the northern half of the study area, a meeting between Madrone, ILMB and MoE was held in early March 2008 to ensure that the final maps would reflect the vision of the client. The client's highest priority was that the maps illustrated where Crown land intersected the forested ecosystems of the CDFmm and depicted the age classes (i.e., structural stages) of those forests. Mapped locations of Garry oak ecosystems, non-forested ecosystems, and wetland/riparian areas were also requested.

The study area was subdivided into 9 1:50,000 maps for ease of interpretation and application. The maps were created in ArcGIS 9.2 using a personal geodatabase as the platform for spatial data storage and management. A custom annotation file was created for the entire study area containing park names, cities and towns, unincorporated areas, major streams and rivers, major highways, and island names. The annotation overlays the following base data: parks and protected areas, Crown land ownership boundaries, major roads, and TRIM streams, rivers, and lakes.

A thematic layer was created that categorized the TEM data into eight classifications: old forest (structural stage 7), mature forest (structural stage 6), young forest (structural stage 5), immature forest (structural stage 3 – 4), Garry oak ecosystem (any Garry oak ecosystem type), non-forested (all non-forested ecosystem types), wetland/riparian (defined ecosystem types), and non-natural (any anthropogenic map unit). Non-forested ecosystems included rock outcrops, meadows, woodlands, coastal bluffs, beaches, and shorelines. Wetland ecosystems included marshes, fens, bogs, swamps and estuaries. Non-natural units included urban areas, agricultural areas, industrial sites, mine sites, roads, and golf courses.

Final thematic maps can be found in Appendix III.

4.10 Project Limitations

Project limitations included accessibility; project scheduling; historic and contemporary disturbance; minimum polygon size; and air photo coverage, year, and scale.

The air photo coverage provided for the project area varied in age, scale and flight direction. Most of the air photos for the entire project were at either 1:16,000 or 1:15,000, with small areas of overlapping flightlines of different scales. In the Cowichan Valley a significant portion of the map was derived from 1:10,000 scale air photos which resulted in a much smaller mean polygon size (9.5 ha) as compared to the rest of the study area (12.5 ha).

The airphotos used to map the study area spanned 25 years (1980 to 2005). A high percentage of the changes that have occurred in the CDFmm have taken place in the past 25 years, particularly due to urbanization but also due to logging. These changes are not accurately reflected in the mapping, thus creating inconsistency of mapping throughout the entire CDFmm, and in particular, of site disturbance, and the loss of ecosystems from urbanization. The use of both colour and black and white air photos (especially in the Cowichan Valley) may have resulted in some inconsistency in interpretation.

Access to private property was the primary limitation to sampling. Public land was inaccessible when surrounded by unroaded private lands. While every effort to sample consistently across the study area was made, not all ecosystem types could be directly observed.

In an effort to minimize this limitation, we based our mapping around these questionable areas on supporting information and background research.

The field sampling schedule (October through February, but including March and April in Year 1) greatly limited the effectiveness of the sampling with regard to complete species lists and general data collection accuracy. Field crews were rushed to complete a significant amount of sampling in an extremely short time frame during the winter months. Some of the data collected on Hornby Island and in the Nanaimo / Ladysmith area was completed in the snow; we suspect that the species lists collected in these plots are incomplete. In general, the winter sampling limited the efficacy of the ecological sampling, particularly with respect to annual indicator plants and classification of species guilds. In some sites, especially those dominated by herbaceous meadows and Garry Oak ecosystems, repeated sampling throughout the year results in a more comprehensive inventory of species composition since communities represented vary substantially throughout the growing season (Erickson and Meidinger, 2007).

The widespread (extensive) nature of disturbance and modification to the ecosystems throughout the study area was a critical factor affecting ecosystem classification and interpretation. Prevalence of exotic species, forest harvesting and regeneration of forest species that differ from those that would occur under natural succession and agricultural utilization altered the species and ecosystem dynamics of the study area.

To ensure sites were classified appropriately, topography, surficial geology, geomorphological history, soils, drainage, and (where available) local historical information was also evaluated where diagnostic plant communities differed from those expected at a site.

5.0 RESULTS AND DISCUSSION

Section 5.1 describes observed soil types and patterns across the landscape with respect to the regional terrain and geomorphology, and representative plant communities for various soil types. The results of the mapped ecosystems and dominant vegetation types are captured in Section 5.2. Section 5.3 contains bioterrain and vegetation descriptions for the ten geographic subregions in the study area. Finally, Section 5.4 summarizes the results integrated across the entire CDFmm study area.

5.1 Soil Landscapes

The following soil descriptions are based on background information summarized on the Ministry of Sustainable Resource Management's website describing the soil landscapes of B.C.³, as well as observations made in the field.

Glaciation has produced a wide range of parent materials from which the majority of the soils of the study area have formed. Soil development in the area also occurs on modern sediments such as fluvial and marine deposits, and also from physical and chemical weathering of exposed bedrock.

The soil Orders identified in the project area include Brunisols, Podzols, Regosols, and to a lesser extent, Gleysols and Organics. The most common soil Great Groups in the area are Dystric Brunisols.

³ Ministry of Sustainable Resource Management Soils website: *The soils landscape of British Columbia* <http://srmwww.gov.bc.ca/soils/landscape/part3.html>. Accessed February 17, 2008

Sombic Brunisols are the next most common. These soils typically occur in the drier, forested sites (i.e., site series CDFmm/01 to /04) supporting predominantly Douglas-fir, Garry oak, and arbutus. These sites may contain grand fir, which prefer slightly moister and richer sites. The most widespread parent materials were medium to coarse-textured till. Also occurring throughout the study area, but to a lesser extent are glaciomarine and glaciofluvial deposits. We also encountered exposed bedrock outcrops in many of our sample plots.

Humo-Ferric Podzols are also common in the project area. They occur predominantly in lower elevation inland valleys that support well established coniferous forests. These soils were also found in drier forested sites with moderately dense Douglas-fir and western hemlock. A moderately dense understory layer was found at most observed sites (which were dominated by younger forests). Podzolic soils are typically associated with glaciomarine and till deposits, and less frequently on glaciofluvial deposits.

Regosols occur where there has been little to no soil development overlying parent materials. Surficial materials associated with Regosols in the project area include fluvial and glaciofluvial deposits, thin colluvial deposits, and weathered bedrock. They occur on recent fluvial deposits such as along smaller streams and rivers, where the vegetation includes black cottonwood, red alder, and willow.

We also encountered Regosols on steep bedrock-controlled slopes. Some of these sites did not support forest vegetation.

Gleysols were rare in the area. They occur on very gently sloping to flat terrain, or low-lying moisture receiving areas associated with marine, glaciomarine or fluvial deposits, or saturated thick till deposits. These soils develop on the wetter ecological sites (i.e., site series CDFmm/06-09, and /12-/14). Typical vegetation includes red alder, black cottonwood, willows, skunk cabbage, and hydrpohytic sedges. However, Gleyed subgroups of Brunisols and Podzols were common.

Organic soils occur where the decay of organic residues is inhibited by a lack of oxygen caused by submersion or saturation (i.e., CDFmm/10 and /11). Organic soils were observed in the project area in wetlands and along forested areas adjacent to wetlands. Fibrisols were the most commonly observed organic Great Group, and typically consist of undecomposed sphagnum and forest litter.

Moderately decomposed organic soils (Mesisols) are less common, and were usually indicated by sedges, hardhack, mosses, willows, grasses, and reeds (i.e., fens). Mesisols are typically richer in nutrients because seepage inflows from adjacent mineral soils rather than from precipitation alone, as is the case with Fibrisols. Well-decomposed Humisols occur only rarely in the project area. Vegetation found on Fibrisols in the study area typically consists of shore pine with Labrador tea and sedges with hardhack.

5.2 Ecosystem Overview

There were a wide range of ecosystems represented throughout the CDFmm (Figures 3 and 4), and many are in disturbed states. This diversity of ecosystems consists of 14 forested site series (including three floodplain units and three fluctuating water table units), 17 wetland and estuary ecosystems, 19 non-forested ecosystems (including nine non-forested and ten sparsely-vegetated site series) and 14 anthropogenic units.

Forested ecosystems consist primarily of second-growth stands. Regenerating stands range from recent clearcuts to dense young stands and maturing forests. The past and current history of land clearing for agricultural practices has contributed to widespread invasive and weedy species colonizing many habitats, especially open sites. Urbanization and some industrialization have contributed to a high loss of ecosystems, particularly in the most southern portion of the study area.

The majority of the CDFmm was actively logged over the last 80 – 90 years, creating large areas of younger forest and alienated lands resulting from land use conversion. Substantial portions of the CDFmm are still dominated by forested ecosystems, consisting of a patchwork of different structural stages, but with negligible old-growth remaining (structural stage 7). In general, forest soils within the CDFmm are mesotrophic to rich (compared to forest soils in other biogeoclimatic zones), and support productive forest stands (Nuszdorfer *et al.*, 1990).

Mature forests of the CDFmm are dominated by the coastal variety of Douglas-fir (*Pseudotsuga menziesii* var. *menziesii*) throughout most of the regional landscape.

Tree species composition of forest stands varies considerably as a result of human disturbance, and also varies from that expected under natural succession regimes due to silviculture and disturbance. Mature and old growth forests are rare in the CDFmm. Less than 1% of the subzone remains in old forest condition as a result of logging and clearing of land for human settlement (BCMWLAP, 2004).

The understory vegetation in upland Douglas-fir forests is characterized by salal (*Gaultheria shallon*), sword fern (*Polystichum munitum*), Oregon-grape (*Mahonia* spp.) and feathermosses.

On drier sites and bedrock outcrops, Douglas-fir stands are joined by arbutus (*Arbutus menziesii*) and occasionally Garry oak (*Quercus garryana*). Baldhip rose (*Rosa gymnocarpa*), common snowberry (*Symphoricarpos albus*), oceanspray (*Holodiscus discolor*) and electrified cat's-tail moss (*Rhytidiadelphus triquetrus*) are frequent in the understory. Forests on lower slopes with moist to subhygric soils will typically contain western redcedar (*Thuja plicata*), red alder (*Alnus rubra*), grand fir (*Abies grandis*), and bigleaf maple (*Acer macrophyllum*) in the canopy, with ladyfern (*Athyrium filix-femina*), spiny wood fern (*Dryopteris expansa*), salmonberry (*Rubus spectabilis*), Indian-plum (*Oemleria cerasiformis*), skunk cabbage (*Lysichiton americanum*), and leafy mosses in the understory.

A variety of other tree species thrive in the mild coastal climate of the CDFmm, adapted to a gradient of site moisture and nutrient regimes.

Less common trees include shore pine (*Pinus contorta* var. *contorta*), Sitka spruce (*Picea sitchensis*), Pacific yew (*Taxus brevifolia*), bitter cherry (*Prunus emarginata*), western flowering dogwood (*Cornus nuttallii*), black cottonwood (*Populus balsamifera* ssp. *trichocarpa*), and rarely, trembling aspen (*Populus tremuloides*).

The disturbed character of many ecosystems presented challenges typical of the CDFmm when identifying and classifying sites in the field. The typical suite of characteristics – including indicator plants, species presence and distribution, and structural attributes – often differed considerably from what was expected in the undisturbed, mature to old seral stands used to develop the classifications (Green and Klinka 1994). This highlighted the importance of assessing soil and bioterrain characteristics, and evaluating site features that did not rely solely on vegetation, particularly mesoslope position, drainage, and disturbance history.




Figure 3. Landscape profile of CDFmm ecosystems in the project area.




Figure 4. Landscape profile of CWHxm ecosystems in the project area.

5.3 Subregional Location Descriptions

This section provides an overview of each of the ten subregions in the study area based on the sequence of mapping from north to south. Table 3 outlines the major communities and geographic location of each subregion. The following descriptions provide notes on special features with respect to their location, typical surficial material and soils as well as dominant vegetation and ecosystems.

Table 3. Geographic Locations of Subregions Mapped in the Project Area

Map	Subregion	Major Communities and Islands
1, 2	Powell River	Powell River, Lund, Hernando Island, Savary Island, Harwood Island
2	North Texada Island	North Texada Island
3	Sunshine Coast, southern Texada Island, and Lasqueti Island	Northern Sechelt area, Thormanby Island, Southern Texada Island, Jedediah Island, Lasqueti Island
4	Denman and Hornby Islands	Denman Island, Hornby Islands
4, 5	Parksville and Qualicum	Bowser, Fanny Bay, Qualicum, Parksville
5, 6, 7	Nanaimo, Ladysmith and Gabriola Island	Nanoose Bay, Nanaimo, Gabriola Island, Newcastle Island, Ladysmith
7, 8	Saltspring Island	Saltspring Island
7, 8	Galiano Island	Galiano Island
7, 8	Chemainus and Cowichan	Chemainus, Crofton, Valdes Island, Thetis Island, Reid Island, Kuper Island, Duncan, Cowichan Bay, Cobble Hill, Mill Bay, Maple Bay
8, 9	Saanich Peninsula and Greater Victoria	Saanich Peninsula, Greater Victoria

Powell River – Maps 1 and 2

Location

This subregion follows the lower elevations on the east side of the Strait of Georgia from Lund to Powell River, including Hernando, Savary, and Harwood Islands.

Bioterrain

Surficial materials within the Powell River subregion are predominantly thick, gravelly and diamictic sandy glaciomarine deposits found at elevations below 100 m asl. Above this, silty sandy diamictic till mantles were observed, grading into the glaciomarine deposits further downslope.

Gravelly, sandy glaciofluvial deposits are located in the southern area where several active and inactive gravel quarries were located. Hummocky and undulating bedrock outcrops are located throughout the area with thin soil overburden and well to very rapidly drained soils. Thin organics were found in small discrete wetland areas accompanied by imperfectly to poorly drained soils.

Glaciomarine materials support Orthic Humic Gleysols with moderately well to imperfectly drained soils on gentle to level ground, and Orthic Humo-Ferric Podzols with well drained soils on steeper gradients. Orthic Dystric Brunsols and Orthic Humic Regosols were observed on rapidly-drained glaciofluvial sites. In general, the glaciomarine-derived soils are moderately well drained, and the till and glaciofluvial deposits are well drained.

Savary Island is comprised of the thick glaciofluvial Quadra Sands with numerous debris slides and slumps observed along steep, eroding cliffs. During the Holocene, wind action has created eolian deposits in the form of dunes in the central and eastern portions of the island.

Vegetation and Ecosystems

The mainland portion of this subregion, including the land surrounding the City of Powell River, has been intensively logged and developed, with the remaining young forests dominated by Douglas-fir, often consisting of planted stands. Similarly, residential and industrial development on the adjacent islands has reduced the distribution of mature forests.

Several rare plants and vegetation communities were found on the shoreline dune ecosystems of Savary Island, including the provincially red-listed plant contorted-pod evening-primrose (*Camissonia contorta*) and the red-listed vegetation community dominated by large-headed sedge (*Carex macrocephala*). The steep sand dunes and cliffs support the Dunegrass–Beach pea site association and exposed soil map units, unless plot data showed otherwise. Douglas-fir is the most common tree species on these sites, with pockets of arbutus. The well-drained soils provide optimal conditions for shore pine on Savary Island. No Garry oak vegetation communities exist in this subregion of the project area.

The CDFmm/CWHxm transition was apparent in the field assessment plots around Lund and Savary Island where western hemlock was found scattered throughout.

North Texada Island – Map 2

Location

The CDFmm occurs along the northwest portion and western edges of Texada Island up to 150 m elevation.

Bioterrain

In this subregion, glaciomarine sediments were expected to occur up to approximately 100 m elevation; however, thick sandy glaciofluvial sediments, and thin diamictic sandy till mantles were found. Well drained glaciofluvial deposits are common throughout the area on gentle slopes and following local depressions. Glaciofluvial deposits give way to morainal and colluvial deposits, as well as bedrock bluffs and hummocks on steeper slopes. The tills tend to be in the form of well drained veneers to thin veneers; however, thicker deposits with moderately well drained soils occur in local depressions.

In the north along the west coast, a complex of thin deposits till and decomposed bedrock were common. Bedrock exposures are common on Texada Island, with numerous exposures of granitic rock. There are also numerous limestone quarries. Thin colluvial deposits are common along the bases of steep slopes. Other colluvial deposits occur along the west central island slopes along the shoreline where slumping was observed in oversteepened glaciofluvial slopes.

Thin, gravelly sandy glaciomarine and marine deposits are less common, occurring only in the Gillies Bay area and along low-lying shorelines. These areas are fine-textured and moderately-well drained. Organic deposits were mapped around small wetlands throughout the area, and were predominately with fibric to mesic veneers.

Dystric and Eutric Brunisols are the most common soils on the island and they develop on both till and glaciofluvial deposits. Orthic Humo-Ferric Podzols were observed in till veneers, and an Orthic Humic Regosol was observed on a fluvial plain.

Vegetation and Ecosystems

Douglas-fir forests dominate Texada Island; however, most areas have been extensively logged. Recent clearcuts and blocks re-planted within the last five years are common.

The abundant deer populations on Texada heavily browse Douglas-fir and select understorey shrubs, modifying ecological succession in the area. Limestone quarries occupy large areas in the northern part of Texada.

The Douglas-fir forests on northern Texada Island are predominately zonal ecosystems (Douglas-fir–Salal). The island contains a relatively high percentage of slightly wetter forests (Western redcedar–Douglas-fir–Oregon beaked moss), higher than the other subregions mapped in the CDFmm. These forests occur on flat or very gentle slopes (less than 5%), where compact marine sediments impede water drainage. Several moist, rich sites (Western redcedar–Skunk cabbage) containing Sitka Spruce, western redcedar and red alder occur on organic veneers. The wetter richer ecosystems (Western redcedar–Grand fir–Foamflower) were generally found alongside drainages between wetlands and from wetlands to the ocean. Rare fern species are known to occur on Texada within this ecosystem types.

Numerous wetlands are located on northern Texada Island and differ in ecosystem type according to drainage patterns. The dominant wetland types include fens and swamps (Sitka sedge–peat moss fen, Pink spirea–Sitka sedge swamp, Sitka willow–Pacific willow–Skunk cabbage swamp) which were dominated by sedges, hardhack and willows and red alder, respectively. Rare plants are associated with these wetland plant communities.

Rocky shoreline bluffs exist along the coast of north Texada Island containing a diverse vegetation assemblage in sharp contrast to the Douglas-fir forest ecosystems inland on deeper soils. The shoreline vegetation communities develop on bedrock and very shallow soils and are composed of several tree, shrub, herb, lichen and moss species that are well adapted to salt spray. A mixture of shore pine, Douglas-fir, Rocky Mountain juniper, common juniper, Nootka rose, and evergreen huckleberry occupy the tree and shrub layers. Life forms range from mature trees to stunted, windswept shrubs. An assortment of forbs, grasses, and sedges can occur in the herb layer, including entire-leaved gumweed and sea plantain. Several open water polygons complexed with sedge wetlands occur on northern Texada Island, supporting rare plant species and vegetation communities.

Sunshine Coast, Lasqueti Island and Southern Texada Island – Map 3

Location

The Sunshine Coast portion of the study area includes Sechelt and extends from Secret Cove to south of Smuggler Cove Marine Provincial Park. Lasqueti Island is located off the east coast of Vancouver Island between Parksville and southern Texada Island. This subregion also includes North and South Thormanby Islands between the Sunshine Coast and Texada Island.

Bioterrain

The Sechelt study area is draped by thick silty, sandy glaciomarine deposits on lower and middle slopes, and to a lesser extent gravelly, sandy glaciofluvial deposits along the lower slopes of the southern part of the subregion where glacial rivers flowed into the ocean. Thinner glaciomarine deposits change to well drained diamictic, sandy till veneers along the middle and upper slopes. Undulating and hummocky bedrock outcrops are distributed throughout the area, but predominantly on the middle and upper slopes, as well as along the shoreline. There are several organic deposits in the area that are associated with small lakes and active floodplains.

North Thormanby Island consists of thick, well drained glaciofluvial deposits (likely Quadra Sands) overlying a thin till veneer. Several slumps and small slides have occurred along the steep, high sandy bluffs adjacent to the shoreline. South Thormanby Island is dominated by a moderately well to well-drained silty sandy glaciomarine mantle. Thin, discontinuous till veneers and thin weathered bedrock veneers are typical on the higher-elevation bedrock outcrops found along the northern and eastern areas of this island.

Lasqueti and the smaller adjacent islands have undulating terrain with a few high elevations that were not inundated by marine and glaciomarine sediments. The majority of the low-lying land is covered by thick, sandy glaciomarine deposits with well to imperfectly drained soils. There are several hummocky bedrock exposures where rubbly, sandy colluvial deposits overlie these glaciomarine sediments. Diamictic sandy till mantles are present from 100 m to more than 150 m, and typically have well to moderately well-drained soils. There are a few areas with glaciofluvial deposits: along ancient river valleys on Lasqueti, and isolated areas along the smaller islands where Quadra Sands likely deposited.

Southern Texada Island is composed of a well-drained sandy gravelly glaciofluvial veneer and blanket, grading to well to moderately well-drained diamictic sandy till veneer along upper slopes.

Soils associated with these areas are Dystric Brunisols and Orthic Humo-Ferric and Humic Podzols. Brunisols are associated with the glaciomarine and colluvial deposits, while Podzols are found in the till and glaciofluvial sediments.

Vegetation and Ecosystems

Lasqueti Island contains diverse wetland ecosystem types. These wetlands are generally surrounded by wet, rich forest ecosystems (Western redcedar–Skunk cabbage) or the drier rich ecosystem (Western redcedar–Grand fir–Foamflower). The forests of Lasqueti Island were varied, but notably were not dominated by the zonal Douglas-fir–Salal type. They were unique within the CDFmm in that dry Douglas-fir–Shore pine–Arbutus forests were abundant, complexed with moss covered rock outcrops (Wallace’s selaginella–Cladina community), in many places grading into dry, rich Douglas-fir–Grand-fir–Oregon-grape forests. Several larger polygons containing older forests on private land were mapped.

On the Sunshine Coast mainland, ecosystems consisted primarily of zonal Douglas-fir–Salal forests transitioning on rocky areas with shallow soils to dry Douglas-fir–Shore pine–Arbutus forests. Wetter, richer ecosystems are localized along streams and creeks (typically Western redcedar–Grand fir–Foamflower). On the wet (subhydric to hydric and fluctuating water table) sites, ecosystems ranged from fluctuating water tables (Western redcedar–Indian-plum and Western redcedar–Slough sedge) to the richer, slightly drier (subhygric) ecosystems (Western redcedar–Grand fir–Foamflower).

North Thormanby and South Thormanby support different forested ecosystem types. North Thormanby is relatively flat, and forests are generally richer zonal (Douglas-fir–Salal) mature forests, with some pockets of wetter forests (Western redcedar–Grand fir–Foamflower). Steep slopes and cliffs with seepage are abundant around the island.

South Thormanby forests are dominated by a mix of Douglas-fir–Shore pine–Arbutus forests, moss covered rock outcrops, and Douglas-fir–Salal forest. Some portions of the island contain moist pockets with fluctuating water tables characterized by Western redcedar–Vanilla-leaf and Western redcedar–Indian-plum associations. Several streams and small drainages contain the richer forested ecosystems.

Denman and Hornby Islands – Map 4

Location

Denman Island and Hornby Islands are just east of the east coast of Vancouver Island adjacent to Deep Bay. This subregion also covers the communities of Fanny Bay and Bowser.

Bioterrain

On both islands, surficial materials consist of predominantly glaciomarine sediments of variable thickness. These deposits extend from the shoreline to the island centres, and range from well to poorly drained. Morainal deposits occur in the central portion of the islands on hill tops above 100 m asl. Steep, exposed bedrock bluffs occur on the western sides of the islands. Both islands are comprised of Nanaimo Group sedimentary rock which consists of sandstone, conglomerate, and mudstone. Organic soils are scattered throughout the area in local depressions and on gentle terrain. These materials vary in thickness from veneers to blankets and are poorly to very poorly drained. Orthic Dystric Brunisols were found in all full plots associated with glaciomarine deposits.

Vegetation and Ecosystems

Mesic zonal Douglas-fir forests dominated Hornby Island; with small areas of the wetter richer ecosystem (Western redcedar–Grand fir–Foamflower). Several Garry oak ecosystems (Garry oak–Brome/mixed grasses) occurred throughout Hornby Island, including the southern tip, in the Heron Rocks area. Associated with the Garry oak ecosystems were native grasses and wildflowers. Native grasslands are extensive along the ocean edge of Helliwell Park and were classified and mapped as Fescue-Camas ecosystems. These sites have very shallow soils over bedrock and are very sensitive. A small trembling aspen stand was also identified at Heron Rocks (Trembling aspen–Slough sedge). Cultivated fields were common throughout the landscape, as well as rural areas where houses, small fields and trees were conspicuous features.

The ecosystems of Denman are similar to Hornby Island. A notable feature of Denman Island is the large recently harvested area in the northeastern portion of the island, which covers approximately one third of the island. The remainder of Denman Island consisted of cultivated fields and rural areas. Mesic zonal forests (Douglas-fir–Salal) were the dominant forest type, with drier forests on thin soils (Douglas-fir–Shore Pine–Arbutus) above the coastline. Fluctuating water table ecosystems (Western redcedar–Vanilla-leaf, Western redcedar–Indian-plum) occurred on level areas on compacted glaciomarine deposits, while the richer ecosystems, (Western redcedar–Grand fir–Foamflower), occurred along streams and between wetlands. Several large and numerous shrub dominated wetlands (Pink spirea–Sitka sedge swamp, Sitka willow–Pacific willow–Skunk cabbage swamp) were found on Denman Island.

Parksville and Qualicum – Maps 4 and 5

Location

The Parksville and Qualicum mapping area extends from several kilometres north of Bowser south to Nanoose, from the ocean to 150 m in elevation.

Bioterrain

The Parksville and Qualicum areas are generally flat, with some undulating terrain along the western boundary. Gravelly sandy marine and glaciomarine deposits of varying thickness dominate the terrain below 110 m asl, and are well to imperfectly drained.

Gravelly sandy glaciofluvial sediments are mapped further inland and along river systems, grading to glaciomarine sediments towards the coast. Drainage in these subregions varies from well to imperfect. The Englishman and Qualicum Rivers have produced thick fluvial deposits along the perimeters of their respective valleys, and there is evidence of similar deposits along their post-glacial floodplains.

Diamictic, sandy morainal deposits are less common, occurring predominantly along the western margins of the study area, on steeper slopes, and in steep cutbanks of creek channels where underlying sediments are exposed. Bedrock outcrops are uncommon, and occur as scattered hummocks throughout the area.

Duric Dystric Brunisols with strongly cemented horizons were found in the thick glaciofluvial and till deposits, with Gleyed Dystric Brunisols found in a marine fan deposit. Orthic Humo-Ferric Podzols were observed on gently sloping and undulating glaciofluvial and glaciomarine deposits. Orthic Humic Regosols occurred in shallow weathered bedrock deposits, as well as with fine textured thick glaciomarine deposits. We also encountered Orthic Humic Gleysols developed where organic veneers overlay fine textured thick glaciomarine deposits.

Vegetation and Ecosystems

This area is generally flat with landforms dominated by glaciomarine and glaciofluvial deposits. These deposits in many places contained compacted horizons, which restricted percolation and created seasonally fluctuating water tables. Fluctuating water table ecosystems were widespread. The wettest type (Western redcedar–Slough sedge) was most frequent adjacent to wetlands, while the most common type mapped was Western redcedar–Indian-plum. On slopes steeper than 5%, drainage was free and forests graded into Douglas-fir–Grand fir–Oregon-grape or Douglas-fir–Salal ecosystem types, depending on nutrient availability. Typical species on these fluctuating water tables were Sitka spruce, western redcedar, and Douglas-fir. Skunk cabbage and lady fern dominated the understory, which developed over a thick layer of peat, corresponding most closely to the Western redcedar–Indian-plum association. These rich sites were mapped in the Bowser area. The rich moist ecosystem type Western redcedar–Grand fir–Foamflower was consistently found in riparian areas with deep soils.

Where the duric horizon was absent, the sandy glaciofluvial soils supported nutrient-poor forests of slow growing, occasionally stunted Douglas-fir and shore pine with understory indicators of low nutrient status, reflecting their subxeric, poor site conditions. This combination corresponded to site series CDFmm/01 (Douglas-fir–Salal; Green and Klinka, 1994), but were typically drier and more nutrient-poor, transitional to site series CDFmm/02 (Douglas-fir–Shore Pine–Arbutus). These forests were extensive in the Parksville and Qualicum areas.

In this subregion, there were few areas of contiguous mature and older forests. Rath Trevor Park and Englishman River Estuary contained the oldest forests in the area. Other mature and older forests were being harvested during field assessments.

Nanaimo, Ladysmith and Gabriola Island – Maps 5, 6, and 7

Location

The Nanaimo subregion extends from Nanoose south through the City of Nanaimo to the Town of Ladysmith, from sea level to 150 m asl.

Bioterrain

The Nanaimo area generally transitions from flat to gently sloping beach deposits along the eastern coastline, to undulating terrain and rolling hills to the west. Slope steepness and the presence of scattered bedrock controlled slopes increases from the west of Highway 19 to the 150 m contour line that serves as the western boundary.

Below 110 m a.s.l., marine and glaciomarine deposits of varying thickness dominate the terrain paralleling the coastline, and particularly east of Highway 19. Pockets of hummocky to undulating bedrock slopes with thin soil veneers of coarse mixed fragments are scattered throughout this coastal lowlands. Coarse-textured glaciofluvial deposits and ablation tills are common (but in places there was uncertainty as to the origin of these deposits; some could have been glaciomarine). Surficial materials to the west of Highway 19 are till blankets and veneers on steeper terrain, deep glaciofluvial materials in valley bottoms and fan deposits where streams leave confined gully systems. Colluvium is rare, as are the steep bedrock slopes. The bedrock observed consisted primarily of sedimentary rocks of the Nanaimo Group.

The sediments in the Nanaimo area and the soils they support are typically coarse, gravelly-sandy, and well-drained; however, drainage can vary greatly depending on slope position and depth to the impermeable horizon. Soils are typically poorly developed Orthic Dystric Brunisols and Regosols. Humo-Ferric Podzols are primarily associated with well-established forests. We encountered isolated areas of imperfectly to poorly drained soils (Gleysols) in depressions and where perched water tables are formed over less permeable underlying sediments. Organic soils are rarely found in these sites.

On Gabriola Island, glaciomarine sediments are the dominant surficial material. These deposits range from gravelly sandy to sandy gravelly in texture, with drainage ranging from rapidly to well-drained. Morainal deposits occur only in the higher elevations at the centre of the island.

Soils tend to be thin near the coast, especially in the northern portion of the island where surficial materials are dominated by well to moderately-well drained veneers to thin veneers of weathered bedrock and glaciomarine sediments, with some sedimentary bedrock exposures. Steep slopes of bedrock and colluvium occur sporadically along the northeast and southwest sides of the island. Bedrock on Gabriola Island consists of Nanaimo Group sedimentary sandstone, conglomerate, and mudstone.

Vegetation and Ecosystems

The few non-fragmented and undisturbed ecosystems that occur throughout the subregion are primarily on Crown land. Many small younger zonal forest patches (structural stage 5) occur throughout the rural areas, in parks, and on Provincial and Federal Crown land. Drier forests complexed with well to rapidly drained moss-covered rock outcrops (Wallace's selaginella– Cladina communities) are common on the elevated sites where soils are shallow.

Garry oak sites are scattered in the Nanaimo area, and are generally associated with moss and herb-covered rock outcrops. Garry oak sites located in the Linley Valley–Departure Bay area exist on the margin of current residential and commercial development. More extensive Garry oak sites are mapped in the Nanoose area, particularly on the Department of National Defense lands. Scattered pockets occur on Provincial Crown land and some on private lands.

Several large herbaceous–moss communities containing rare plant species were mapped around Harewood Plains, Cable Bay, and the Linley Valley in Nanaimo. The Linley Valley has a wide range of mature forested ecosystems with high wildlife value, including riparian, zonal, Garry oak and dry forests. The Nanaimo River estuary is also a significant contiguous natural area, although highly disturbed. The previously undescribed map unit Nootka rose–Pacific crab apple was mapped along the river. A high percentage of the Nanaimo area has been mapped as urban and rural.

Saltspring Island – Maps 7 and 8

This area has been mapped in detail during Year 1 and has been described in Madrone (2008a).

Galiano Island – Maps 7 and 8

Location

Galiano Island is located east of Saltspring Island between Vancouver Island and the Fraser Valley and northwestern Washington State.

Bioterrain

Galiano Island consists of long narrow bedrock ridges and steep bluffs overlooking small valleys and shorelines (i.e., Lovers' Leap Viewpoint, Montague Harbour area, and Bluffs Park). Mount Sutil (323 m) and Mount Galiano (341 m) are the highest points on the island. There several other steep bedrock-exposed slopes aligned northwest to southeast. North-south passes across the island are limited to the area between Pebble Beach and Retreat Cove and the eastern tip of the island. Surface water drainage is controlled by the ridges, with predominantly seasonal streams flowing perpendicularly from the ridges to the beaches.

The steep, well to rapidly drained bedrock ridges and colluvial veneer complexes lead to interior lowlands with small valleys. Hummocky terrain in the lowland areas consists of well to moderately well drained, silty sandy tills, and pockets of gravelly sandy glaciofluvial blankets (observed to the east of Laughlin Lake only). In valleys, and on uniform slopes at and below 100 m, are finer textured glaciomarine and marine deposits that lead to beaches surrounding the island. Drainage in these small valleys and on the low gradient slopes leading to the shoreline ranged from well to imperfectly drained depending on the proportion of silts and clays. Soils encountered in both the till, marine and glaciomarine environments ranged from Orthic to Gleyed Dystric Brunisols. In some valley depressions, a few small wetlands and lakes have formed, with associated organic deposits typically developing moderately well decomposed soils.

Vegetation and Ecosystems

Galiano Island is the driest of the Gulf Islands characterized by long narrow ridges of sandstone which run the full length of the island. Stands of dry Douglas-fir – arbutus forests (Douglas-fir–Shore Pine–Arbutus) complexed with mesic zonal Douglas-fir forests (Douglas-fir–Salal) grow along these ridges. A mosaic of Garry oak ecosystems (Garry oak–Brome/mixed grasses) interspersed with grass (Fescue–Camas) and moss dominated ecosystems (Cladina–Wallace’s selaginella) occur along south facing slopes.

Along the base and between these ridges is a combination of rich, moist and productive forests of western redcedar and bigleaf maple (Western redcedar–Grand fir–Foamflower) mixed with zonal Douglas-fir forests (Douglas-fir–Salal). Where the underlying drainage is poor shrub and herb-dominated wetlands are present. Relative to the other Gulf Islands, Galiano Island has more mature and older forest (greater than 140 year old) and fewer cultivated fields.

Chemainus and Cowichan – Maps 7 and 8

Location

This subregion includes Chemainus, the Cowichan Valley, Valdez Island, Thetis Island, and Kuper Island. The western boundary extends up to 150 m.

Bioterrain

In the Cowichan Valley, glaciomarine sediments of variable thickness are the dominant sediments of the flat to undulating slopes below 100 m. These occur mainly south of the Cowichan River, especially on the south shores of Cowichan Bay. These marine sediments are easily seen in the scarp faces in the community of Cowichan Bay extending south to Mill Bay. Soil textures in our plots were typically silty to clayey silt. Drainage was typically moderate to imperfect.

Extensive deposits of sandy gravelly glaciofluvial materials also occur throughout the area. Soils that have developed on these materials are typically well drained and give rise to Orthic Dystric Brunisols (Jungen, 1985) on gentle slopes with well drained to moderately well drained soils.

Fluvial deposits are more common towards the estuaries of both the Cowichan and Chemainus Rivers. Smaller fluvial deposits occur in thin bands along the main creeks in the study area such as Stocking, Porter, Bonsall, and Hollings Creeks. Soils active fluvial environments along Hollings Creek are typically classified as a moderately well-drained Cumulic Humic Regosol.

Till is common above 100 m in this subregion. Most of the higher topography occurs north of the Cowichan River and includes the slopes of Mount Tzouhalem (497 m), Maple Mountain (525 m), Mt. Richards (353 m), Stoney Hill (200 m), and Grouse Hill (191 m).

These upper slopes are typically covered with moderately to rapidly drained sandy diamictic veneers and blankets of till. Orthic Dystric Brunisols are common (Jungen, 1985). Three full plots were completed in till-derived polygons. All were on gentle to moderately steep slopes that were well drained. Two were classified as Orthic Dystric Brunisols and the third was classified as Orthic Sombric Brunisol. Colluvial deposits and bedrock outcrops also occur on these slopes.

Organic deposits are rare in this area. The largest area of organic soil occurs along Richards Creek north of Somenos Lake, and another pocket occurs west of Dougan's Lake. These large areas of organic soils are used for agriculture; consequently no full plot data was collected in these areas. Soils data in an organic deposit were collected in a pocket of humic organic material towards Shawnigan Lake on Crown land at the end of Owl Road. This area of very poorly drained organic material was classified as a Typic Humisol.

Vegetation and Ecosystems

The Cowichan Valley is a developed rural landscape, characterized by a mosaic of cultivated fields, forests and residential areas. Urban centres occur along the main highway corridor. There are extensive shoreline and coastal bluffs, most of which have been modified for residential housing. In the Cowichan Valley, most sampling was conducted on Cowichan Valley Regional Parks (CVRD) land, District of North Cowichan municipal forests, and Crown land. The majority of the landscape in the Cowichan Valley contains second-growth forest.

Zonal forest types (Douglas-fir–Salal) are very diverse in this area, and can be found on moderately dry (SMR 2) to submesic to mesic (SMR 4) site conditions. Mesic zonal sites occur on deep, medium textured soil, gentle slopes, and are dominated by salal in the understory. Douglas-fir dominates the canopy with admixtures of western redcedar, bigleaf maple, and grand fir. Submesic zonal sites tend to occur on shallow and/or coarse soils, steep slopes, warm aspects, and upper to crest mesoslope positions. Drier sites support less salal which is replaced by tall Oregon-grape. In submesic to subxeric Douglas-fir–Salal forests, arbutus becomes more common in the canopy, with less western redcedar, grand fir, and bigleaf maple.

Ecosystems that tend to develop on drier sites, such as the Douglas-fir–Shore Pine–Arbutus forest, typically occur in a mosaic with the non-forested Wallace’s selaginella–Cladina community; this combination was quite common in the area. Both ecosystem types occurred on very shallow (often <10 cm) soils (i.e., Orthic Regosols to Non-soil) and warm aspects, occasionally occurring on cool aspects. The Douglas-fir–Oniongrass forest type was much less common, and almost always developed on warm aspect, upper slopes.

The Western redcedar–Grand fir–Foamflower association was also very common in this subregion in draws, adjacent to creeks or in moisture-receiving lower slopes. This forest type is distinguished from the Douglas-fir–Salal forest type predominantly by abundant bigleaf maple, with varying amounts of grand fir, western redcedar, and red alder, with Douglas-fir less frequent and salal only a minor understorey component. The understory is very lush in structural stages that contain canopy gaps, with rich site indicator species such as foamflower and vanilla-leaf.

Garry oak ecosystems are present but limited in extent and distribution. Most were found on dry rocky outcrops such as those found on Cobble Hill, Mt. Tzouhalem, and Maple Mountain. Scattered large Garry oak trees occur around the east side of Somenos Lake throughout residential developments, and scattered groves can be found in cultivated fields. Historically these ecosystems were mapped over a much broader range of sites than their current distribution (Lea, 2006).

The Douglas-fir– Grand fir –Oregon grape forest type is most often restricted to side slopes of creek draws, occurring both on till and glaciomarine substrates. This forest type occurred in various site conditions, but was relatively limited in this area. These forests have a constant presence of Douglas-fir, western redcedar, and sword fern.

Active floodplain ecosystems most often occur along main river corridors such as those of the Cowichan, Chemainus and Koksilah Rivers, but can occur to a lesser extent along some of the smaller tributaries. Quite often these riparian forests are mature or old forests, older than surrounding forests.

Extensive stands of grand fir, western redcedar and bigleaf maple characterize high-bench floodplain sites while stands of black cottonwood occur on mid-bench sites. Low-bench floodplain sites are shrub dominated by willow species and red alder. All have high wildlife value.

Ecosystems characteristic of fluctuating water tables, particularly the Western redcedar–Indian-plum were very common, while the wetter analogue Western redcedar–Slough sedge was restricted to wetland margins.

Extensive wetland complexes occur around Somenos Lake, as well as smaller scattered wetlands throughout the Cowichan Valley. The most frequent wetlands are the shrub swamp types: the Spirea–Sitka sedge and the Sitka willow–Pacific willow–Skunk cabbage associations. Cattail marshes were typically associated with lakes, ponds, and the margins around shallow open water. The estuary at Cowichan Bay was characterized by the Nootka rose–Pacific crab apple shrub-dominated ecosystem in complex with estuary meadows and marshes.

Saanich Peninsula and Greater Victoria (CRD) – Maps 8 and 9

Location

The CDFmm encompasses the entire Saanich Peninsula and Greater Victoria. This subregion is characterized by an urban landscape with developed shorelines and coastal bluffs.

Bioterrain

The Saanich/Victoria subregion can be roughly divided into four physiographic sections based on surficial materials and surface shape (Stanley-Jones and Benson, 1973).

The largest section in the Saanich/Victoria map area is the mountainous Highlands and Goldstream section in the west area along Finlayson Arm. The topography is moderately rugged, including Mount Finlayson (416 m) and Mount Work (446 m). Surficial materials in this area are typically well to moderately well-drained sandy diamictic till veneers. Outcrops of bedrock are common on high peaks and steep slopes, and the bedrock is hummocky to undulating.

The hummocky nature of the bedrock gives rise to small pockets of wetter soils in the lower areas. Colluvial deposits were observed along the steep slopes of along Finlayson Arm.

The second largest section is comprised of nearly level to gently rolling glaciomarine deposits that are characteristic of the Saanich Peninsula up to approximately 100 m. Plot data gathered in this area showed parent material that was typically silty glaciomarine with moderate to imperfect drainage. Horth Hill (136 m), Mount Newton (302 m), and Bear Hill (220 m) all rise above the rolling countryside on the peninsula and have well-drained sandy till veneers. Bedrock outcrops were also common. Due to the competent nature of the bedrock, colluvial deposits were uncommon. Glaciomarine materials are also found on the southwest side of the study area in Metchosin below 100 m (e.g., at Witty's Lagoon).

The third section is an extensive deposit of glaciofluvial sediments in the vicinity of Langford and Colwood. These deposits can be easily observed at the large gravel pit operations on Metchosin Road south of Esquimalt Lagoon. Soils in this area are sandy gravelly in texture and typically well to moderately well drained.

The fourth section is the Greater Victoria area. Much of this area was mapped as anthropogenic, indicating surficial materials have been significantly altered by excavation and pavement.

Vegetation and Ecosystems

Several provincial and regional parks, such as Mt. Douglas, Mill Hill, and Mt. Tolmie have excellent examples of Douglas-fir forests below Garry oak ecosystems.

These sites display clear vegetation zonation with elevation: Douglas-fir-Salal on deep, medium textured soil and gentle slopes; Douglas-fir-Shore pine-Arbutus forests on shallow, coarse soils and warm aspects; and Garry oak-Brome/mixed grasses woodlands on upper to crest slope positions. In general, the parks around Victoria and Saanich Peninsula represent fragmented islands of forested and non-forested ecosystems intersected by development and transportation corridors. Many of the remnant patches of forest tend to occur on sites with shallow soils and on rocky outcrops (e.g., Mt. Douglas and Mt. Tolmie). Douglas-fir-Salal forests in the subregion tend to occur on shallow soils with submesic to subxeric moisture regimes.

Most of the extant forests are second-growth (structural stages 5 and 6), with patches of older forest in John Dean Park.

The Western redcedar-Grand fir-Foamflower site series in this area was typically restricted to the bases of draws, riparian areas, and moisture-receiving lower slopes. These forest types often had a substantial hardwood component comprised of bigleaf maple and red alder, with varying amounts of grand fir and western redcedar. The understory was variable, but could support very dense vegetation featuring indicator species typical of richer sites.

Sites with fluctuating water tables occur throughout the Saanich Peninsula. The Western redcedar-Indian-plum ecosystem type was by far the most abundant, occurring frequently on level sites and particularly through channels of cultivated fields in sites with deep glaciomarine sediments. The wetter sites supporting the Western redcedar-Slough sedge ecosystem were less abundant, typically adjacent to small wetlands. The Western redcedar-Vanilla-leaf fluctuating water table ecosystem type was uncommon in this subregion.

Garry oak ecosystems are far more abundant in the Saanich Peninsula than in other areas. Two main Garry oak units occur: those developed on rocky outcrops and those on deep soils with a parkland character. The former was mapped as QB with a shallow soil modifier; the latter was mapped as QB with no shallow soil modifier. The rocky outcrop Garry oak units were most abundant near Victoria proper, represented on Mt. Douglas, Mt. Tolmie, and in Saanich and adjacent areas around the Highlands, Mt. Work, Gowlland Tod Provincial Park, and around Prospect and Elk Lakes.

The parkland Garry oak unit was most commonly found interspersed through the cultivated fields of Central and North Saanich where rich soils have developed from deep glaciomarine deposits to support large, robust Garry oak trees, in sharp contrast with the stunted Garry oak trees found on rocky outcrops.

The estuaries of Goldstream, McKenzie Bight, Witty's Lagoon and Island View Beach were a complex of the Nootka rose–Pacific crab apple shrub ecosystem and estuary marsh and meadows. Non-forested wetlands are uncommon in this area, and, where present, consist of shrub-dominated wetlands such as the Sitka willow–Pacific willow–Skunk cabbage swamp type. This unit occurs around Swan Lake and Blenkinskop Lake, among others, usually in a complex with marsh ecosystems such as the Cattail marsh.

5.4 Ecosystem Representation and Condition in the Study Area

A total of 17,621 ecosystem polygons were delineated in the CDFmm project area (212,660 ha), with an average polygon size of 11.6 hectares (ranging from 0.20 to 7,093 ha). The vast majority of mapped polygons (17,188 or 98%) were in the CDFmm subzone; the remaining polygons are in the CWHxm subzone. A list of the TEM units mapped in the project area is provided in the expanded legend in Appendix V. The legend includes site descriptions, assumed modifiers, site characteristics, photographs and plot reference numbers of each ecosystem unit mapped in the study area.

The project area does not include the Gulf Islands National Park Reserve (GINPR, established in 2003), which features nearly 3,500 ha of diverse ecosystems on primarily sandstone-derived, well-drained, drier landforms that are strongly influenced by ocean spray at lower elevations. Incorporating ecosystem information from the mapping completed in the GINPR would yield a more accurate representation of ecosystem distribution and status across the entire CDFmm. However, the defined scope for this project is restricted to interpretation and discussion of the previously defined project area.

5.4.1 Site Series and Ecosystem Units

The following results illustrate the ecosystems mapped in the CDFmm, including forested, non-forested, sparsely-vegetated, wetland and anthropogenic units. Tables 4 – 10 list the various ecosystem units mapped, the total area of each unit, and the percentage of the total project area.

In the CDFmm, 14 forested site series (including three floodplain units and three fluctuating water table units) comprised 61% (127,411 ha) of the study area. Natural non-forested ecosystems (including nine non-forested and ten sparsely-vegetated site series) made up 4% (9,888 ha) of the CDFmm. Lastly, 17 wetland and estuary ecosystems and 14 anthropogenic units were mapped totaling 2% (3,394 ha) and 33% (69,048 ha), respectively (Figure 5).




Figure 5. Distribution of ecosystem categories in the CDFmm by area and percentage.

The most common forested ecosystems within the CDFmm study area were the zonal Douglas-fir-Salal, the richer/wetter Western redcedar-Grand fir-Foamflower forest type, and the drier Douglas-fir-Shore Pine-Arbutus forest type comprising 37%, 7%, and 5% of the study area, respectively. Forested fluctuating water table sites (Western redcedar-Vanilla-leaf, Western redcedar-Indian-plum, Western redcedar-Slough sedge) were also fairly common collectively, representing 6% or almost 12,500 ha. Douglas-fir-Oniongrass, Shore pine-Sphagnum, Western redcedar-Skunk cabbage, and the forested floodplain ecosystems were the least common in the CDFmm study area (Table 4 and Figure 6).

Table 4. Ecosystems Mapped in the CDFmm Project Area.

Map code	Site series	Ecosystem name	Area (ha)	Area (%)
Forested site series				
DS	01	Douglas-fir–Salal	77063	36.9
DA	02	Douglas-fir–Shore Pine–Arbutus	11044	5.29
DO	03	Douglas-fir–Oniongrass	1083	0.52
DG	04	Douglas-fir–Grand fir–Oregon-grape	3794	1.82
RK	05	Western redcedar–Douglas-fir–Oregon beaked moss	3935	1.89
RF	06	Western redcedar–Grand fir–Foamflower	15399	7.38
LS	10	Shore pine–Sphagnum	154	0.07
RC/Ws53	11	Western redcedar–Skunk cabbage	925	0.44
Forested floodplain site series				
RS	07	Western redcedar–Snowberry high bench floodplain	396	0.19
CD/Fm50	08	Black cottonwood–Red-osier dogwood medium bench floodplain	224	0.11
CW	09	Black cottonwood–Willow low bench floodplain	38	0.02
Forested fluctuating water table site series				
RV	12	Western redcedar–Vanilla-leaf	3107	1.49
RP	13	Western redcedar–Indian-plum	7956	3.64
CS	14	Western redcedar–Slough sedge	1426	0.68
Total Forested			127411	61




Figure 6. Distribution of forested ecosystem units in the CDFmm by area, including floodplain and fluctuating water table site series: note log area scale.

Natural non-forested ecosystems, including non-forested and sparsely vegetated ecosystems, represented 4% of the CDFmm study area, but, incorporating data from the Gulf Islands National Park Reserve would likely increase the coverage of these ecosystem types.

Typically the majority of these ecosystems were rare throughout the CDFmm. The most common ecosystems were Cladina-Wallace's selaginella which occurred over approximately 1.4% of the study area and were generally associated with rock outcrops. Garry oak ecosystems (Garry oak-Oceanspray, Garry oak-Moss, Garry oak-Brome/mixed grasses) only comprised about 0.5% within the CDFmm (Table 5 and Figure 7).

Table 5. Natural Non-Forested (Non-Forested and Sparsely Vegetated) Ecosystems Mapped in the CDFmm Project Area

Map code	Site series	Ecosystem name	Area (ha)	Area (%)
Non-forested ecosystems				
AS	00	Trembling aspen-Slough sedge	12	0.006
FC	00	Fescue-Camas	381	0.18
GO	00	Garry oak-Oceanspray	4	0.002
LM	00	Dunegrass-Beach pea	127	0.06
OM	00	Garry oak-Moss	26	0.01
OR	00	Oceanspray-Rose	5	0.002
RA	00	Nootka rose-Pacific crab apple	978	0.47
QB	00	Garry oak-Brome/mixed grasses	81	0.04
SC	00	Cladina-Wallace's selaginella	2866	1.37
Sparsely vegetated units				
BE		Beach	526	0.25
CL		Cliff	132	0.06
GB		Gravel bar	137	0.07
LA		Lake	1911	0.92
MU		Mudflat	68	0.03
OW		Open water	552	0.27
PD		Pond	145	0.07
RI		River	312	0.15
RO		Rock outcrop	1623	0.78
TA		Talus	3	0.001
Total Natural Non-forested			9888	4




Figure 7. Distribution of natural non-forested ecosystem units in the CDFmm by area, including non-forested and sparsely vegetated ecosystems: note log area scale.

Wetland and estuary ecosystems, including bogs, fens, marshes, and swamps, represented only 2% of the CDFmm study area. The most common ecosystems are Pink spirea–Sitka sedge swamp, Sitka willow–Pacific willow–Skunk cabbage swamp and Cattail marsh, which comprised 0.6%, 0.4%, and 0.2% of the study area, respectively (Table 6 and Figure 8).

Table 6. Wetland and Estuary Ecosystems Mapped in the CDFmm Project Area

Map code	Site series	Ecosystem name	Area (ha)	Area (%)
Wetland and estuary ecosystems				
Ed01	00	Tufted hairgrass-Meadow barley estuarine meadow	134	0.06
Ed03	00	Arctic rush-Alaska plantain estuarine meadow	21	0.01
Em01	00	Widgeon-grass tidal flat	34	0.02
Em02	00	Glasswort-Sea-milkwort estuarine marsh	131	0.06
Em03	00	Seashore saltgrass	50	0.02
Em05	00	Lyngbye's sedge estuarine marsh	232	0.11
Wb50	00	Labrador tea-Bog-laurel-Peat-moss bog	6	0.003
Wf51	00	Sitka sedge-Peat-moss fen	135	0.07
Wf52	00	Sweet gale-Sitka sedge fen	30	0.01
Wf53	00	Slender sedge-White beak-rush fen	13	0.006
Wm05	00	Cattail marsh	347	0.17
Wm06	00	Great bulrush marsh	1	0.001
Wm50	00	Sitka sedge-Hemlock-parsely marsh	74	0.04
Wm51	00	Three-way sedge marsh	22	0.01
Ws50	00	Pink spirea-Sitka sedge swamp	1258	0.60
Ws51	00	Sitka willow-Pacific willow-Skunk cabbage swamp	892	0.42
Ws52	00	Red alder-Skunk cabbage swamp	11	0.01
Total wetlands and estuaries			3394	2




Figure 8. Distribution of wetland and estuary ecosystem units in the CDFmm by area: note log area scale.

Anthropogenic units represented a disproportionately large 33% of the CDFmm study area. Ecosystems converted primarily to rural residential land use covered approximately 9.5% of the CDFmm study area while urban areas accounted for about 9.4%. These areas were common throughout the entire CDFmm but were most prevalent near or adjacent to the major city centres, such as Victoria, and Nanaimo. Fields, including those in use for cultivation and grazing, or fallow fields, covered approximately 10.7% (Table 7 and Figure 9).

Table 7. Anthropogenic Units Mapped in the CDFmm Project Area

Map code	Site series	Ecosystem name	Area (ha)	Area (%)
Anthropogenic units				
CF		Cultivated field	22348	10.7
CO		Cultivated orchard	176	0.08
CV		Cultivated vineyard	74	0.04
ES		Exposed soil	538	0.26
GC		Golf course	1266	0.61
GP		Gravel pit	903	0.43
IN		Industrial	1555	0.75
MI		Mine	675	0.32
RE		Reservoir	41	0.02
RN		Railway surface	33	0.02
RZ		Road surface	1708	0.82
RW		Rural	19857	9.52
TZ		Mine tailings	180	0.09
UR		Urban	19696	9.44
Total anthropogenic			69048	33




Figure 9. Distribution of anthropogenic map units in the CDFmm by area: note log area scale.

The CWHxm accounted for 2% of the project study area. In the CWHxm, 13 forested site series (including two floodplain units and three fluctuating water table units) comprised 91% (3,757 ha) of the mapped area. Natural non-forested ecosystems, including four non-forested, four sparsely-vegetated and four wetland ecosystems, made up 4% (174 ha). A total of four anthropogenic unit types were mapped over 5% (210 ha) of the CWHxm within the study area (Figure 10).




Figure 10. Distribution of ecosystem categories in the CWHxm by area and percentage.

The most common forested ecosystems mapped within the CWHxm study area were the zonal mesic Western hemlock–Douglas-fir–Oregon beaked moss, the slightly drier Douglas-fir/Western hemlock–Salal, and the richer mesic Western hemlock/western redcedar–Deer fern associations, which comprised 45%, 16%, and 8% of the CWHxm study area, respectively. Forested fluctuating water table sites and the forested floodplain ecosystems were rare within the study area (Table 8 and Figure 11).

Table 8. Forested Ecosystems Mapped in the CWHxm Project Area

Map code	Site series	Ecosystem name	Area (ha)	Area (%)
Forested site series				
HK	01	Western hemlock–Douglas-fir–Oregon beaked moss	1880	45.4
DC	02	Douglas-fir–Shore pine–Cladina	195	4.69
DS	03	Douglas-fir–Western hemlock–Salal	680	16.4
DF	04	Douglas-fir–Sword fern	215	5.18
RS	05	Western redcedar–Sword fern	266	6.42
HD	06	Western hemlock–Western redcedar–Deer fern	324	7.82
RF	07	Western redcedar–Foamflower	121	2.92
RC (Ws53)	12	Western redcedar–Sitka spruce–Skunk cabbage	12	0.28
Forested floodplain site series				
SS	08	Sitka spruce–Salmonberry high bench floodplain	1	0.02
CW (FI50)	10	Black cottonwood–Willow low bench floodplain	1	0.02
Forested fluctuating water table site series				
RB	13	Western redcedar–Salmonberry	33	0.79
RT	14	Western redcedar–Black twinberry	28	0.67
CS	15	Western redcedar–Slough sedge	4	0.10
Total Forested			3757	91




Figure 11. Distribution of forested ecosystem units mapped in the CWHxm by area, including floodplain and fluctuating water table site series: note log area scale.

Natural non-forested ecosystems, including non-forested, sparsely vegetated, and wetland ecosystems, represented 4% of the CWHxm study area mapped. The most common ecosystem was the Arbutus–Hairy manzanita which occurred on 1.3% of the study area (53 ha). Wetland ecosystems collectively accounted for approximately 0.5% of the CWHxm (Table 9 and Figure 12).

Table 9. Natural Non-Forested (Non-Forested, Sparsely Vegetated, and Wetland) Ecosystems Mapped in the CWHxm Project Area.

Map code	Site series	Ecosystem name	Area (ha)	Area (%)
Non-forested ecosystems				
AM	00	Arbutus–Hairy manzanita	53	1.29
FC	00	Fescue–Camas	5	0.12
QB	00	Garry oak–Brome/mixed grasses	1	0.02
SC	00	Cladina–Wallace’s selaginella	29	0.70
Sparsely vegetated units				
CL		Cliff	22	0.52
LA		Lake	30	0.74
OW		Open water	1	0.01
RO		Rock outcrop	11	0.28
Wetland ecosystems				
Wb50	00	Labrador tea–Bog-laurel–Peat-moss bog	1	0.02
Wf53	00	Slender sedge–White beak-rush fen	14	0.33
Ws50	00	Pink spirea–Sitka sedge swamp	6	0.14
Ws51	00	Sitka willow - Pacific willow–Skunk cabbage swamp	2	0.05
Total natural non-forested			174	4




Figure 12. Distribution of natural non-forested ecosystem units mapped in the CWHxm by area, including non-forested, sparsely vegetated, and wetland ecosystems: note log area scale.

Anthropogenic units represented 5% of the CWHxm study area mapped with rural residential land use and cultivated fields comprising 3.07% and 1.57%, respectively (Table 10 and Figure 13).

Table 10. Anthropogenic Units Mapped in the CWHxm Project Area.

Map code	Site series	Ecosystem name	Area (ha)	Area (%)
Anthropogenic units				
CF		Cultivated field	65	1.57
ES		Exposed soil	18	0.43
RW		Rural	127	3.07
RZ		Road surface	1	0.02
Total anthropogenic			210	5




Figure 13. Distribution of anthropogenic map units mapped in the CWHxm by area: note log area scale.

Ecosystem accounts for each map unit are provided in the expanded legend (Appendix VI). Each unit includes a description of the site, soil and terrain characteristics observed in each ecosystem, and a representative photo.

5.4.2 Structural Stage

Young forests (structural stage 5) are the most common structural stages in the CDFmm study area, representing 45% of the landscape (Figure 14). Young forests typically range from 40 – 80 years old in this area, with self-thinning evident, producing a forest canopy which is beginning to differentiate into distinct layers.

These forests reflect logging or another disturbance that has occurred over the past 40 – 80 years (forest harvesting being the most common in the project area). In the CWHxm within the study area, young forests and immature forests (structural stage 4 and 5) were the most typical at 41% and 37%, respectively (Figure 15).

There was extremely little old growth, (structural stage 7) mapped in the CDFmm zone. Only 610 ha of old forest were mapped on Lasqueti Island and small pockets scattered on southern Vancouver Island. Immature forest (structural stage 4) and mature forest (structural stage 6) represented 12% and 13% of the CDFmm study area, respectively. Lastly, structural stages 1 – 3 combined occurred over 30% of the landscape representing logged or otherwise disturbed forests, natural non-forested ecosystems (non-forested, sparsely vegetated, wetland), and a small percentage of anthropogenic areas (Figure 14).




Figure 14. Distribution of structural stages in the CDFmm by area and percentage.




Figure 15. Distribution of structural stages in the CWHxm by area and percentage.

5.4.3 Disturbance

The CDFmm subzone covers a diverse range of landforms, land uses, ecosystems, and infrastructure. Historic and modern settlement and resource extraction have displaced many naturally occurring ecosystems across the region, limiting their distribution.

Disturbance is extensive across the east side of Vancouver Island, especially near the major centres (e.g., Greater Victoria and Nanaimo), and to a lesser extent, on the Gulf Islands, and Sunshine Coast. Disturbances consisted primarily of urban and rural residential development, followed by logging and agriculture.

Larger protected areas on eastern Vancouver Island such as East Sooke Park and Thetis Lake Park harbour substantial representative areas of CDFmm forests, but are still impacted to varying degrees by disturbance and invasive species. All of the Gulf Islands have had more or less intensive human disturbance, while some retain a variety of remnant CDFmm ecosystems.

Intensive agriculture is relatively common, especially in glaciomarine and glaciolacustrine deposits. These rich, gently undulating to flat sites support productive crop growth and forage. Wetland modification through drainage, excavation, and vegetation change also impacts on ecosystem representation and ecological integrity.

Seasonally flooded fields are recognized and mapped as sensitive ecosystems through the Sensitive Ecosystem Inventory (SEI) program due to their high habitat value (Ward *et al.* 1998); however the CDFmm TEM mapping project includes these areas as cultivated fields.

Cultivated fields are widely distributed throughout the CDFmm, from the Saanich Peninsula to the Cowichan Valley and Salt Spring Island. Cultivated fields predominantly occur on deep glaciomarine deposits and are heavily used for haying, grazing, and agriculture. Some small-scale orchards and vineyards have been established on gently to moderately sloping well-drained terrain. Between major urban centers and intermixed with cultivated fields, rural residential development is predominant.

Rural residential and cultivated field modification involves land conversion or alienation such that natural processes and functions are hampered or modified to the extent that native plant communities are not supported.

This does not negate their potential function, should they be restored or set aside for reclamation. Seasonally flooded fields, hedgerows, ditches, and windbreaks also provide habitat for a variety of wildlife species, serving to maintain some degree of landscape habitat connectivity and support forage species, shelter, roosting sites, nest locations, and prey populations.

Forest harvesting has been occurring in the region for more than 100 years. The dominant silvicultural system is clearcutting with and without reserves, with lesser amounts of single-tree and group selection systems. Intensive agriculture and hobby farms are relatively common and widespread, especially in glaciomarine and glaciolacustrine deposits. Glaciofluvial sites and some gentler till deposits support orchards and vineyards. The entire area has been subject to varying degrees of wetland diking, drainage, and excavation.

Another modern threat to CDF forests is fire suppression. The characteristic Garry oak ecosystems represent a disturbance driven, fire-dependent disclimax type and has had widespread declines concurrent with fire suppression (MacDougall and Turkington 2004; MacDougall et al. 2004).

Fire frequency is suspected to have historically fluctuated based on cultural practices (McDougall and Turkington 2004; McCoy et al. 2006). The Garry Oak Ecosystems Restoration Team (GOERT) has collected a wide range of contemporary and historical baseline data to provide context regarding the extent, condition, composition, and resilience of these ecosystems, including their responses to restoration treatments.

Within the CDFmm study area, 40% of the landscape was determined to be disturbed by anthropogenic factors, such as urbanization and farming, as well as logging, fires and biotic effects, such as grazing (Figure 16). Similar findings occurred in the CWHxm study area with anthropogenic disturbances and logging making up 42% of the landscape (Figure 17). These values, however, likely drastically underestimate the actual amount of disturbance, especially within the CDFmm, as the air photos used for mapping the majority of the study area were out-dated, going back as far as 1980. Based on field observations, many areas within the CDFmm were recently disturbed and therefore not reflected on the air photos. The area from Deep Bay to Cowichan, which includes Qualicum, Parksville, Nanaimo, Ladysmith, and Duncan have air photos ranging in age from 1980 to 1998 (Appendix II).

Other than Victoria and Saanich, these areas have likely been the most altered in the past 10 to 20 years and contain substantially more disturbed areas (urbanized, agricultural and logged) than accounted for in this study.




Figure 16. Distribution of disturbance types in the CDFmm by area and percentage.




Figure 17. Distribution of disturbance types in the CWHxm by area and percentage.

5.4.4 Invasive species

Exotic and indigenous invasive plants displace native and endemic species, particularly those with narrow ecological niches and slow growth and recruitment. Transportation corridors and recreational areas, agricultural, residential, and disturbed sites all increase the presence and abundance of weedy invasive species which tend to flourish in pioneer habitats.

The past century has seen the gradual introduction of many exotic species into the CDFmm, many of which have become naturalized. The number and abundance of these species has been dramatically increasing over the past several decades, driving a concerted effort by local, regional, and provincial agencies and non-profit groups to gather data and develop control and eradication strategies.

The most prevalent exotic species in the study area are: Scotch broom, English ivy, foxglove, Daphne-laurel, agronomic grasses, gorse, English hawthorn, English oak species, holly, and Himalayan blackberry. There are many others that are localized to particular habitats, such as rush skeletonweed, thistles, giant cow-parsnip, white poplar, knotweed species, knapweed species, carpet burweed, yellow-flag iris, Eurasian milfoil, purple loosestrife, and spartina cordgrass. Typical strategies of successful invasive and weedy species include the ability to colonize mineral seedbeds, rapid early growth, prolific reproduction, robust seed banks, vegetative spread, apomixes and/or self-fertilization, and dense monotypic growth habits.

These characteristics modify existing microsites in such a way that native species are out-shaded, displaced spatially, unable to access sufficient moisture and nutrients, or restricted in reproductive capacity.

Displacement of rare indigenous species by exotic and indigenous invasive plants has also contributed substantially to the decline of Garry oak ecosystems (MacDougall and Turkington 2004) and ecosystems of the CDFmm in general. Introduced and feral domestic livestock serve as vectors for many weedy species. Other invasives have been deliberately introduced as ornamentals, or forage crops. Site disturbance that exposes mineral seedbeds hastens colonization by weedy invasive species. Scotch broom, many common agronomic grasses, gorse, English hawthorn and oak, holly, and a wide range of other life forms aggressively colonize nearly all sites in the CDFmm. Their adaptations to disturbance, including rapid growth, prolific reproduction, robust seed banks, vegetative spread, and dense monotypic growth habits, tend to render sites unsuitable for many native species, particularly those that require specialized microhabitats.

Wetland ecosystems (including ponds, marshes, lakes, streams, estuaries, and fens) have experienced similar invasive species which have caused changes in ecological representation and condition. Purple loosestrife, Eurasian milfoil, yellow flag iris, cordgrass species, and many other escaped cultivated plants have altered the hydrological regime, decreased habitat diversity, and displaced native aquatic and estuarine plants.

Many particularly insidious weeds affecting wetlands originated via escapes from cultivation. These habitat types are increasingly less frequent throughout the CDFmm and their sensitivity to ecological disturbance makes them particularly vulnerable to degradation (MacDougall et al. 2006).

5.4.5 Rare Elements

Forestry and intense development pressure has resulted in fragmentation and permanent loss of habitat. The British Columbia Conservation Data Centre (CDC) has listed 35 ecological communities in the CDFmm, representing all ecosystems within this zone and subzone, as provincially at risk (red-or blue-listed) (CDC 2008: see Appendix I). Most of these ecosystems are described as “imperiled” (S2) or “critically imperiled” (S1).

The vegetation of the CDFmm includes ecosystems and species that are considered rare due to their limited occurrences and restricted range. Most of these are at the northern limits of their distribution and include species that occupy seaside, aquatic, rock outcrop, and forested habitats. A few species of animals, plants, and mosses are entirely restricted or endemic to the CDFmm zone, such as Macoun's meadowfoam. Endemic species and unique assemblages of ecosystems provide genetic-, species-, and landscape-level components that are important for maintaining biodiversity. Due to its restricted distribution and the aforementioned impacts, many plant and wildlife species found in the CDFmm are listed federally under Schedules of the *Species at Risk Act*, and others are on the provincial Red and Blue lists due to their rarity and/or special concern. These species may also be afforded additional protection under legislation such as the *Forest and Range Practices Act* (FRPA) and associated regulations. The CDC lists 63 plants and animals (including six unique populations of stickleback and six owl species/subspecies) whose habitat overlaps or falls completely within the CDFmm [March 2008]. Animal species include invertebrates, insects, fish, insectivorous mammals, raptors, and bats. Plant species include a range of taxa, although many are annual flowering plants (CDC 2008).

Provincially, the CDC also lists 218 plants and animals as either blue-or red-listed in the CDFmm; including 127 vascular plant species, three non-vascular plants, 53 vertebrate animals, and 35 invertebrate animals. Of these, 101 plants and animals are blue-listed and 117 are red-listed.

Federally, the Committee on the Status of Endangered Species in Canada (COSEWIC) ranks 78 plants and animals with conservation status, 45 of these being endangered, denoting species facing imminent extirpation or extinction. And lastly, the *Species at Risk Act* (SARA) lists 68 of these plants and animals on their Schedule 1, which classifies wildlife species at risk as being either extirpated, endangered, threatened, or special concern. Table 11 outlines the number of species of plants and animals listed under provincial and federal jurisdictions in the CDFmm.

Table 11 Conservation Status of Plants and Animals Listed in the CDFmm

Rare Species Category	Vascular Plants	Non-vascular Plants	Vertebrate Animals	Non-vertebrate Animals
Number of Listed Individuals	127	3	53	35
Jurisdiction	Conservation Status		Number of Listed Individuals	
Provincial Rank	Blue-listed		101	
	Red-listed		117	
Federal Rank	COSEWIC (Endangered)		78 (45)	
	SARA Schedule 1		68	

The rank and status of species and ecosystems are based on several factors, and screened by committees of experts: rarity/abundance based on inventory data, connectivity based on spatial distribution, changes in abundance or distribution (e.g., fragmentation), areas with protected status, and external factors influencing the species or ecosystem (e.g., exotic pests or diseases, pending land use changes, etc.).

The CDFmm is home to a rich diversity and abundance of native species and habitats. Programs within the Ministry of Environment endeavour to conserve biodiversity, provide a variety of opportunities for the use and enjoyment of wildlife in B.C., and maintain a balance between the needs of wildlife and the needs of the people. The Species at Risk Coordination Office of the Integrated Land Management Bureau and the Ministry of the Environment are developing a provincial strategy for the conservation and recovery of species at risk in the province. The intention of the conservation framework will foster proactive, priority-driven conservation approaches, based on reliable science, which will lead to better outcomes for species at risk (ILMB, 2007)

6.0 RECOMMENDATIONS

The results of the project are limited by the fact that the mapping product that accompanies this report does not reflect an accurate measure of the urbanized, agricultural and logged area that has replaced mature and old forest since the date of many of the photos. To realize the full capacity of this project, and to fulfill the objectives and intent, we make the following recommendations.

- Conduct an analysis of the “depletion layer” using the TEM product provided. The TEM product can be updated by identifying the areas that are now logged using satellite imagery and updating those polygon labels to more accurately map the extent and impacts to ecosystems at risk.
- Update identified sensitive ecosystem inventory and quantify alterations to identified SEI polygons since the last SEI update (AXYS, 2005) where appropriate at this scale.
- Almagamate the TEM completed for the Gulf Islands National Park Reserve with this project.
- Conduct TEM on the CWHxm subzone, which also contains similar ecosystems and species that are listed provincially.
- Complete TEM for Vancouver Island. Many areas have already been mapped and this information can be integrated, gaps identified, and an integrated map of the entire Island can be used to provide detailed information for land use planning. Further, existing mapping could be updated fairly efficiently using satellite imagery where structural stages have changed.

7.0 CONCLUSION

The data collected and maps produced from this report represent a rich resource that can be used for many applications. One such application is as baseline inventory for a landscape management strategy across the CDFmm.

Data collected in this project was supported by a total of 1458 plots, as well as 331 background plots, resulting in 1234 field checked polygons which provided ecosystem information on geology, soils, site characteristics and vegetation for final ecosystem mapping. Level-five survey intensity was accomplished with 9% polygon inspection (including background plot data). The extensive background research, wide sampling framework, and rigorous internal and external quality assurance in place provide a high degree of confidence in the interpolations made during photo interpretation.

Mapping revealed that although the project area contains a diverse range of forested and non-forested ecosystems, including wetlands, and herbaceous meadows, most of them have been impacted by disturbance over the past century. A large proportion of the land area has been alienated by conversion to urban, rural, agricultural, and industrial use. Forest harvesting, mining, and associated activities have further restricted succession of ecosystems, maintaining a disproportionately high part of the landscape in early seral stages. Negligible mature and old forest remains within the CDFmm.

While some of these patterns are certainly associated with land tenure, an integrated framework for land use, such as the existing Vancouver Island Land Use Plan, can be used to guide priorities for regional outcomes. Considering the CDFmm as a single, integrated system may aid planners to better evaluate the potential impacts of resource and land use decisions across jurisdictions with respect to this restricted and high-value ecological zone.

8.0 REFERENCES CITED

- AXYS Environmental Consulting Ltd. 2005. Redigitizing of sensitive ecosystems inventory polygons to exclude disturbed areas. Report to Canadian Wildlife Service. Sidney, B.C. 27 pp.
- B.C. Conservation Data Centre (CDC). 2008. BC Species and Ecosystems Explorer. B.C. Min. Environ., Victoria, B.C. <http://srmapps.gov.bc.ca/apps/eswp/> (accessed May 31, 2008).
- B.C. Ministry of Environment. 2006. Standard for mapping ecosystems at risk in British Columbia: An approach to mapping ecosystems at risk and other sensitive ecosystems. Version 1. B.C. Min. Environ., Victoria, B.C. 98 pp.
- B.C. Ministry of Environment, Lands and Parks (BCMELP). 1999a. Ecosystems in British Columbia at Risk: Coastal Douglas-fir ecosystems. B.C. Ministry of Environment, Lands, and Parks. Victoria, B.C. 6 pp.
- B.C. Ministry of Environment, Lands and Parks (BCMELP). 1999b. Terrestrial Ecosystem Mapping of Jedediah Island Provincial Park. 1999. Prepared for BC Parks, Victoria, B.C. 87 pp.
- B.C. Ministry of Environment, Lands and Parks (BCMELP). 2001. Sooke Hills Wilderness and Mount Wells Regional Parks: Terrestrial Ecosystem Mapping. B.C. Ministry of Environment, Lands, and Parks. Victoria, B.C. 186 pp.
- B.C. Ministry of Water, Land and Air Protection (BCMWLAP). 2004. Douglas-fir/dull Oregon-grape (*Pseudotsuga menziesii*/*Mahonia nervosa*) in Accounts and Measures for Managing Identified Wildlife – Accounts V. 2004. B.C. Ministry of Water, Land and Air Protection, Victoria, B.C.
- Bichler, A.J., E.D. Brooks, and P.T. Bobrowsky. 2002. Sunshine Coast aggregate potential mapping project. In: B.C. Ministry of Energy and Mines, Geological Fieldwork 2001, Paper 2002-1, pp. 147-153.
- Blyth, H.E. and N.W. Rutter. 1992. Quaternary geology of Southeastern Vancouver Island and Gulf Islands (92B/5, 6, 11, 12, 13 and 14). In Geological Fieldwork 1992. B.C. Ministry of Energy, Mines and Petroleum Resources, Paper 1993-1 pp. 407-413.

- Blyth, H.E. and N.W. Rutter. 1993a. Surficial geology of the Victoria Area (NTS 92B/6). B.C. Ministry of Energy, Mines and Petroleum Resources, Open File 1993-23.
- Blyth, H.E. and N.W. Rutter. 1993b. Surficial geology of the Sidney Area (NTS 92B/11). B.C. Ministry of Energy, Mines and Petroleum Resources, Open File 1993-24.
- Blyth, H.E. and N.W. Rutter. 1993c. Surficial geology of the Duncan area (NTS 92B/13). B.C. Ministry of Energy, Mines and Petroleum Resources, Open File 1993-27.
- Blyth, H.E., N.W. Rutter, and L.M. Sankeralli 1993. Surficial geology of the Shawnigan area (NTS 92B/12). B.C. Ministry of Energy, Mines and Petroleum Resources, Open File 1993-26.
- Bobowsky, P.T. and J.J. Clague. 1995. Quaternary geology of southern Vancouver Island. B5 field trip guidebook. Geological Association of Canada, Mineralogical Association of Canada, joint annual meeting Victoria, B.C., May 17-19, 1995.
- Clague, J.L. 1994. Quaternary stratigraphy and history of south-coastal British Columbia. *In: Geology and geological hazards of the Vancouver region, southwestern British Columbia*, J.W.H. Monger (ed.) Geological Survey of Canada, Bull. 481:181-192.
- Dunster, K. and J. Booth. 2001. Terrestrial Ecosystem Mapping of Helliwell Provincial Park. 2001. Prepared for BC Parks, Strathcona District Office, B.C.
- Erickson, W.R. and D.V. Meidinger. 2007. Garry oak (*Quercus garryana*) plant communities in British Columbia: A guide to identification. B.C. Min. For., Res. Br., Victoria, B.C. Tech. Rep. 040.
- Gabrielse, H., J.W.H. Monger, J.O. Wheeler, and C.J. Yorath. 1991. Morphological belts, tectonic assemblages, and terrains. *In: Geology of the Cordilleran Orogen in Canada*. H. Gabrielse and C.J. Yorath (eds.); Geological Survey of Canada, Geology of Canada. 4(2/A):15-28.
- Green, R.N. and K. Klinka. 1994. A field guide to site identification and interpretation for the Vancouver Forest Region. Land Manage. Handb. 28. B.C. Min. For., Victoria, B.C.

- Holland, S.S. 1976. Landforms of British Columbia: A Physiographic Outline. Bulletin 48. Province of British Columbia.
- Howes, D.E. and E. Kenk. 1997. Terrain classification system for British Columbia. Revised edition. Manual 10. B.C. Ministry of Environment, Lands, and Parks. Victoria, B.C.
- Jungen, J.R. 1985. Soils of southern Vancouver Island. Report No. 44, B.C. Soil Survey, B.C. Ministry of Environment, Surveys and Mapping Branch. MoE Tech. Rep. 17.
- Jungen, J.R., P. Sanborn, and P.J. Christie (eds.). Soils of southeast Vancouver Island, Duncan-Nanaimo area. Co-published by Ministry of Agriculture and Food and Government of B.C., Victoria, B.C.
- Lea, T. 2006. Historical Garry oak ecosystems of Vancouver Island, British Columbia, pre-European contact to the present. *Davidsonia* 17(2):34–50.
- Madrone Environmental Services Ltd. 2002. Terrestrial Ecosystem Mapping and Management Recommendations for Fourteen Department of National Defence Properties CFB Esquimalt. Contract report to Department of Natural Resources and Department of National Defence. 67 pp.
- Madrone Environmental Services Ltd. 2003. Mount Maxwell Terrestrial Ecosystem Mapping and ecological assessment. Contract report to B.C. Ministry of Water, Land and Air Protection, Environmental Stewardship Division, Vancouver Region. Vancouver, B.C. 124 pp.
- Madrone Environmental Services Ltd. 2005. Environmental Assessment: Proposed Development Property, Cable Bay, B.C. Contract report to Cable Bay Lands Inc. 94 pp.
- Madrone Environmental Services Ltd. 2008a. Terrestrial Ecosystem Mapping of Saltspring Island (1:20,000). Contract report to Integrated Land Management Bureau. 103 pp.
- Madrone Environmental Services Ltd. 2008b. Sandstone Ecological Assessment: Proposed Development Property, South Nanaimo, B.C. Contract report to Northwest Properties. 89 pp.

- Mackenzie, W.H. and A. Banner. 2001. A Classification Framework for Wetlands and related Ecosystems in British Columbia: third approximation. <http://www.for.gov.bc.ca/research/becweb/> (accessed January, 2008).
- Mackenzie, W.H. and J.R. Moran. 2004. Wetlands of British Columbia: a guide to identification. Res. Br., B.C. Min. For., Victoria, BC. Land Manage. Handb. No. 52.
- Mathews, W.H. (compiler). 1986: Physiography of the Canadian Cordillera; Geological Survey of Canada, Map 1701A, scale 1: 5000 000.
- Monahan, P.A., V.M. Levson, P. Henderson, and A. Sy. 2000. Quaternary geological map of Greater Victoria TRIM sheets (92B043, 044, 053, 054). Geoscience Map 2000-2. B.C. Ministry of Energy and Mines, Victoria, B.C.
- Muller, J.E. 1980. Geology of Victoria. Map 1553A, Geological Survey of Canada.
- Nuszdorfer, F.C., K. Klinka, and D.A. Demarchi. 1990. Coastal Douglas-Fir zone. In Ecosystems of British Columbia. D. Meidinger and J. Pojar (compilers and editors). B.C. Min. For., Victoria, B.C. Special Rep. Ser. 6. pp. 81-93.
- Province of British Columbia. 2000. Vancouver Island summary land use plan. Province of B.C., Victoria, B.C.
- Resource Inventory Standards Committee (RISC). 1998a. Field manual for describing terrestrial ecosystems. Co-published by B.C. Min. For. and B.C. Min. Environ. Prov. of B.C., Victoria, BC.
- Resource Inventory Standards Committee (RISC). 2000. Standard for digital terrestrial ecosystem mapping (TEM) data capture in British Columbia. Ecosystem Technical Standards & Database Manual, Version 3.0. RISC, Ecosystems Working Group, Victoria, B.C.
- Resource Inventory Standards Committee (RISC). 2002. Digital terrestrial ecosystem mapping data capture (DC) user's guide. RISC, Ecosystems Working Group, Victoria, B.C.

- Resource Inventory Standards Committee (RISC). 2004. Standard for terrestrial ecosystem mapping (TEM) - digital data capture in British Columbia, Version 3.0, Errata No. 1.0. RISC, Ecosystems Working Group, Victoria, B.C.
- Resources Inventory Committee (RIC). 1996. Standards and guidelines for terrain mapping in British Columbia. RISC, Surficial Geology Task Group, Victoria, B.C.
- Resources Inventory Standards Committee (RISC). 1998b. Standard for terrestrial ecosystem mapping in British Columbia. Ecosystems Working Group, Resources Inventory Standards Committee. Victoria, B.C.
- Stanley-Jones, C.V. and W.A. Benson, editors. 1973. An inventory of land resources and resource potentials. Report to the Capital Regional District, Victoria, B.C.
- Stoffels, D. 2000. Natural disturbance and large scale vegetation succession scenarios for the Columbia Forest District Columbia Mountains Caribou Project [Appendix 9: fire return intervals by BEC zone]. Contract report to B.C. Min. For., Res. Br., Prince Rupert Regional Office, by Truffula Tree Ecological Services, Smithers, B.C.
- Yorath, C.J. and N.W. Nasmith. 1995. The geology of Southern Vancouver Island. Orca Book Publishers, Victoria, B.C.



APPENDIX I

Red- and Blue-listed Ecosystems and Species in the CDFmm

Appendix I. Red- and Blue-listed Ecosystems and Species in the CDFmm

Red- and blue-listed ecosystems in the CDFmm

Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit
<i>Abies grandis</i> / <i>Mahonia nervosa</i>	grand fir / dull Oregon-grape	G1	S1	Red	CDFmm/04
<i>Abies grandis</i> / <i>Tiarella trifoliata</i>	grand fir / three-leaved foamflower	G1	S1	Red	CDFmm/06
<i>Alnus rubra</i> / <i>Carex obnupta</i> [<i>Populus balsamifera</i> ssp. <i>trichocarpa</i>]	red alder / slough sedge [black cottonwood]	G1	S1	Red	CDFmm/14
<i>Festuca idahoensis</i> ssp. <i>roemerii</i> - <i>Koeleria macrantha</i>	Roemer's fescue - junegrass	G1	S1	Red	CDFmm/00; CWHxm1/00
<i>Pseudotsuga menziesii</i> / <i>Melica subulata</i>	Douglas-fir / Alaska oniongrass	G1	S1	Red	CDFmm/03
<i>Quercus garryana</i> - <i>Arbutus menziesii</i>	Garry oak - arbutus	G1	S1	Red	CDFmm/00
<i>Quercus garryana</i> / <i>Bromus carinatus</i>	Garry oak / California brome	G1	S1	Red	CDFmm/00
<i>Quercus garryana</i> / <i>Holodiscus discolor</i>	Garry oak / oceanspray	G1	S1	Red	CDFmm/00
<i>Thuja plicata</i> / <i>Achlys triphylla</i>	western redcedar / vanilla leaf	G1	S1	Red	CDFmm/12
<i>Thuja plicata</i> / <i>Oemleria cerasiformis</i>	western redcedar / Indian-plum	G1	S1	Red	CDFmm/13
<i>Carex macrocephala</i> Herbaceous Vegetation	large-headed sedge Herbaceous Vegetation	G1G2	S1S2	Red	CDFmm/00; CWHvh1/00; CWHwh1
<i>Populus tremuloides</i> / <i>Malus fusca</i> / <i>Carex obnupta</i>	trembling aspen / Pacific crab apple / slough sedge	G1G2	S1S2	Red	CDFmm/00
<i>Arbutus menziesii</i> / <i>Arctostaphylos columbiana</i>	arbutus / hairy manzanita	G2	S2	Red	CDFmm/00; CWHxm1/00
<i>Carex lasiocarpa</i> - <i>Rhynchospora alba</i>	slender sedge - white beak-rush	G2	S2	Red	CDFmm/Wf53; CWHmm1/Wf53; CWHmm2/Wf53; CWHxm1/Wf53; CWHxm2/Wf53
<i>Myosurus minimus</i> - <i>Montia</i> spp. - <i>Limnanthes macounii</i>	tiny mousetail - montias - Macoun's meadow-foam	G2	S1	Red	CDFmm/00
<i>Pseudotsuga menziesii</i> / <i>Mahonia nervosa</i>	Douglas-fir / dull Oregon-grape	G2	S2	Red	CDFmm/01
<i>Salix sitchensis</i> - <i>Salix lucida</i> ssp. <i>lasianдра</i> / <i>Lysichiton americanus</i>	Sitka willow - Pacific willow / skunk cabbage	G2	S2	Red	CDFmm/Ws51; CWH/Ws51; ICH/Ws51



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit
<i>Deschampsia cespitosa</i> ssp. <i>beringensis</i> - <i>Aster subspicatus</i>	tufted hairgrass - Douglas' aster	G3	S3	Blue	CDFmm/Ed02; CWH/Ed02
<i>Deschampsia cespitosa</i> ssp. <i>beringensis</i> - <i>Hordeum brachyantherum</i>	tufted hairgrass - meadow barley	G3	S3	Blue	CDFmm/Ed01
<i>Menyanthes trifoliata</i> - <i>Carex lasiocarpa</i>	buckbean - slender sedge	G3	S3	Blue	CDFmm/Wf06; CWHws1/Wf06; ICHwk1/Wf06; IDFdk2/Wf06; SBSdk/Wf06
<i>Myrica gale</i> / <i>Carex sitchensis</i>	sweet gale / Sitka sedge	G3	S2	Red	CDFmm/Wf52; CWHmm1/Wf52; CWHmm2/Wf52; CWHvh2/Wf52; CWHwm/Wf52; CWHxm1/Wf52; CWHxm2/Wf52
<i>Salicornia virginiana</i> - <i>Glaux maritima</i>	American glasswort - sea-milkwort	G3G4	S2	Red	CDFmm/Em02; CWH/Em02
<i>Typha latifolia</i> Marsh	common cattail Marsh	G5	S3	Blue	BGxh1/Wm05; BGxh2/Wm05; BGxw1/Wm05; CDFmm/Wm05; CWHdm/Wm05; CWHxm1/Wm05; CWHxm2/Wm05; IDFdk3/Wm05; IDFdm2/Wm05; PPxh1/Wm05
<i>Distichlis spicata</i> var. <i>spicata</i> Herbaceous Vegetation	seashore saltgrass Herbaceous Vegetation	G5	S1S2	Red	CDFmm/Em03
<i>Alnus rubra</i> / <i>Lysichiton americanus</i>	red alder / skunk cabbage	GNR	S2S3	Blue	CDFmm/11
<i>Carex lyngbyei</i> Herbaceous Vegetation	Lyngbye's sedge herbaceous vegetation	GNR	S3	Blue	CDFmm/Em05
<i>Eleocharis palustris</i> Herbaceous Vegetation	common spike-rush	GNR	S3	Blue	BGxw2/Wm04; CDFmm/Wm04; ESSFdv d/Wm04; ESSFdv/Wm04; IDfxm/Wm04; SBSdk/Wm04; SBSmk2/Wm04
<i>Artemisia campestris</i> - <i>Festuca rubra</i> / <i>Racomitrium canescens</i>	northern wormwood - red fescue / grey rock-moss	GNR	S1	Red	CDFmm/n/a
<i>Dulichium arundinaceum</i> Herbaceous Vegetation	three-way sedge	GNR	S2	Red	CDFmm/Wm51; CWHmm1/Wm51; CWHxm2/Wm51; ICHwk1/Wm51
<i>Juncus arcticus</i> - <i>Plantago macrocarpa</i>	arctic rush - Alaska plantain	GNR	S1	Red	CDFmm/Ed03; CWH/Ed03
<i>Pinus contorta</i> / <i>Sphagnum</i> spp. CDFmm	lodgepole pine / peat-mosses CDFmm	GNR	S1	Red	CDFmm/10
<i>Pseudotsuga menziesii</i> - <i>Arbutus menziesii</i>	Douglas-fir - arbutus	GNR	S2	Red	CDFmm/02
<i>Ruppia maritima</i> Herbaceous Vegetation	beaked ditch-grass Herbaceous Vegetation	GNR	S2	Red	CDFmm/Em01; CWH/Em01
<i>Thuja plicata</i> - <i>Pseudotsuga menziesii</i> / <i>Eurhynchium oreganum</i>	western redcedar - Douglas-fir / Oregon beaked-moss	GNR	S1	Red	CDFmm/05



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit
<i>Thuja plicata</i> / <i>Symphoricarpos albus</i>	western redcedar / common snowberry	GNR	S1	Red	CDFmm/07

Search Criteria

Ecological Communities

AND BC Conservation Status: Red (Extirpated, Endangered, or Threatened) OR Blue (Special Concern)

AND MOE Regions: 1- Vancouver Island, 2- Lower Mainland

AND BGC Zone, Subzone, Variant, Phase: CDFmm

Sort Order: Scientific Name Ascending

Accessed May 31 2008

Red- and blue-listed species in the CDFmm

Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
Fish						
<i>Acipenser medirostris</i>	Green Sturgeon	G3	S1N	Red	CDF; CWH	Estuarine; Marine; Riverine
<i>Acipenser transmontanus</i> pop. 4	White Sturgeon (Lower Fraser River population)	G4T2Q	S2	Red	CDF; CWH; IDF	Estuarine; Lacustrine; Marine; Riverine
<i>Oncorhynchus clarkii clarkii</i>	Cutthroat Trout, <i>clarkii</i> subspecies	G4T4	S3S4	Blue	BWBS; CDF; CWH; ICH; SBS	Estuarine; Lacustrine; Marine; Riverine
<i>Salvelinus malma</i>	Dolly Varden	G5	S3S4	Blue	BWBS; CDF; CWH; ESSF; ICH; MH; SBS	Estuarine; Lacustrine; Marine; Riverine
<i>Gasterosteus</i> sp. 16	Vananda Creek Limnetic Stickleback	G1	S1	Red	CDF	Lacustrine
<i>Gasterosteus</i> sp. 17	Vananda Creek Benthic Stickleback	G1	S1	Red	CDF	Lacustrine
<i>Gasterosteus</i> sp. 2	Enos Lake Limnetic Stickleback	G1	S1	Red	CDF	Lacustrine
<i>Gasterosteus</i> sp. 3	Enos Lake Benthic Stickleback	G1	S1	Red	CDF	Lacustrine
<i>Gasterosteus</i> sp. 4	Paxton Lake Limnetic Stickleback	G1	S1	Red	CDF	Lacustrine
<i>Gasterosteus</i> sp. 5	Paxton Lake Benthic Stickleback	G1	S1	Red	CDF	Lacustrine
Amphibians						
<i>Rana aurora</i>	Red-legged Frog	G4	S3S4	Blue	CDF; CWH	Lacustrine; Palustrine; Riverine; Terrestrial

Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Rana pipiens</i>	Northern Leopard Frog	G5	S1	Red	CDF; ICH; IDF; PP	Lacustrine; Palustrine; Riverine; Terrestrial
Reptiles						
<i>Chrysemys picta pop. 1</i>	Western Painted Turtle - Pacific Coast Population	G5TNR	S2	Red	CDF; CWH; MH	Lacustrine; Palustrine; Riverine
<i>Contia tenuis</i>	Sharp-tailed Snake	G5	S1	Red	CDF	Subterranean; Terrestrial
<i>Pituophis catenifer catenifer</i>	Gopher Snake, <i>catenifer</i> subspecies	G5T5	SX	Red	CDF; CWH	Palustrine; Terrestrial
Birds: Anseriformes						
<i>Branta canadensis occidentalis</i>	Canada Goose, <i>occidentalis</i> subspecies	G5T2T3	S1N	Blue	CDF; CWH	Lacustrine; Palustrine; Terrestrial
Birds: Pelecaniformes						
<i>Phalacrocorax auritus</i>	Double-crested Cormorant	G5	S3B	Blue	CDF; CWH; ICH; SBPS	Estuarine; Lacustrine; Marine; Palustrine; Riverine; Terrestrial
<i>Phalacrocorax pelagicus pelagicus</i>	Pelagic Cormorant, <i>pelagicus</i> subspecies	G5TU	S2B	Red	CDF; CWH	Estuarine; Marine; Terrestrial
<i>Phalacrocorax penicillatus</i>	Brandt's Cormorant	G5	S1B,S4N	Red	CDF; CWH	Estuarine; Marine
Birds: Ciconiiformes						
<i>Ardea herodias fannini</i>	Great Blue Heron, <i>fannini</i> subspecies	G5T4	S3B,S4N	Blue	CDF; CWH	Estuarine; Lacustrine; Palustrine; Riverine; Terrestrial
<i>Botaurus lentiginosus</i>	American Bittern	G4	S3B	Blue	BG; BWBS; CDF; CWH; ICH; IDF; PP; SBPS; SBS	Estuarine; Palustrine
<i>Butorides virescens</i>	Green Heron	G5	S3S4B	Blue	CDF; CWH	Estuarine; Lacustrine; Palustrine; Riverine
Birds: Faconiformes						
<i>Accipiter gentilis laingi</i>	Northern Goshawk, <i>laingi</i> subspecies	G5T2	S2B	Red	CDF; CWH; MH	Terrestrial
<i>Falco peregrinus anatum</i>	Peregrine Falcon, <i>anatum</i> subspecies	G4T4	S2B	Red	BG; BWBS; CDF; CWH; IDF; MS; PP; SBS	Estuarine; Terrestrial
<i>Falco peregrinus pealei</i>	Peregrine Falcon, <i>pealei</i> subspecies	G4T3	S3B	Blue	CDF; CWH	Estuarine; Lacustrine; Marine; Riverine; Terrestrial

Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
Birds: Charadriiformes						
<i>Hydroprogne caspia</i>	Caspian Tern	G5	S3B	Blue	CDF; CWH; IDF; SBS	Estuarine; Lacustrine; Marine; Palustrine; Riverine; Terrestrial
<i>Brachyramphus marmoratus</i>	Marbled Murrelet	G3G4	S2B,S4N	Red	CDF; CWH; MH	Estuarine; Lacustrine; Marine; Terrestrial
<i>Fratercula cirrhata</i>	Tufted Puffin	G5	S3B,S4N	Blue	CDF; CWH	Marine; Terrestrial
Birds: Colombiiformes						
<i>Patagioenas fasciata</i>	Band-tailed Pigeon	G4	S3S4B	Blue	CDF; CWH; ICH	Palustrine; Terrestrial
Birds: Cuculiformes						
<i>Coccyzus americanus</i>	Yellow-billed cuckoo	G5	SXB	Red	CDF; CWH	Estuarine; Palustrine; Terrestrial
Birds: Strigiformes						
<i>Tyto alba</i>	Barn Owl	G5	S3	Blue	BG; CDF; CWH	Palustrine; Terrestrial
<i>Asio flammeus</i>	Short-eared Owl	G5	S3B,S2N	Blue	BG; BWBS; CDF; CWH; ICH; IDF; PP; SBPS; SBS; SWB	Estuarine; Palustrine; Terrestrial
<i>Athene cunicularia</i>	Burrowing Owl	G4	S1B	Red	BG; CDF; CWH; IDF; PP	Terrestrial
<i>Glaucidium gnoma swarthi</i>	Northern Pygmy-Owl, <i>swarthi</i> subspecies	G5T3Q	S3	Blue	CDF; CWH; MH	Terrestrial
<i>Megascops kennicottii kennicottii</i>	Western Screech-Owl, <i>kennicottii</i> subspecies	G5T4	S3	Blue	CDF; CWH; IDF	Palustrine; Terrestrial
Birds: Piciformes						
<i>Melanerpes lewis</i>	Lewis's Woodpecker	G4	S2B	Red	BG; CDF; CWH; ICH; IDF; PP	Palustrine; Terrestrial
<i>Melanerpes lewis</i> pop. 1	Lewis's Woodpecker (Georgia Depression population)	G5TXQ	SXB	Red	CDF; CWH	Palustrine; Terrestrial
Birds: Passeriformes						
<i>Eremophila alpestris strigata</i>	Horned Lark, <i>strigata</i> subspecies	G5T2	SX	Red	CDF; CWH	Terrestrial
<i>Hirundo rustica</i>	Barn Swallow	G5	S3S4B	Blue	BAFA; BG; BWBS; CDF; CWH; ESSF; ICH; IDF; IMA; MH; MS; PP; SBPS; SWB	Estuarine; Lacustrine; Palustrine; Riverine; Terrestrial

Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Progne subis</i>	Purple Martin	G5	S2S3B	Blue	CDF; CWH; ICH	Estuarine; Lacustrine; Palustrine; Terrestrial
<i>Sialia mexicana</i> pop. 1	Western Bluebird (Georgia Depression population)	G5TNRQ	SHB	Red	CDF; CWH	Palustrine; Terrestrial
<i>Poocetes gramineus affinis</i>	Vesper Sparrow, <i>affinis</i> subspecies	G5T3	S1B	Red	CDF	Terrestrial
<i>Sturnella neglecta</i> pop. 1	Western Meadowlark (Georgia Depression population)	G5TNRQ	SXB	Red	CDF; CWH	Terrestrial
Mammals						
<i>Aplodontia rufa rufa</i>	Mountain Beaver, <i>rufa</i> subspecies	G5T4?	S3	Blue	CDF; CWH; MH	Terrestrial
<i>Myodes gapperi occidentalis</i>	Southern Red-backed Vole, <i>occidentalis</i> subspecies	G5T5	S1	Red	CDF; CWH	Palustrine; Terrestrial
<i>Lepus americanus washingtonii</i>	Snowshoe Hare, <i>washingtonii</i> subspecies	G5T3T5	S1	Red	CDF; CWH	Palustrine; Terrestrial
<i>Sorex palustris brooksi</i>	American Water Shrew, <i>brooksi</i> subspecies	G5T2	S2	Red	CDF; CWH	Lacustrine; Palustrine; Riverine
<i>Sorex rohweri</i>	Olympic Shrew	G4G5	S1	Red	CDF	Terrestrial
<i>Sorex trowbridgii</i>	Trowbridge's Shrew	G5	S3S4	Blue	CDF; CWH	Palustrine; Terrestrial
<i>Corynorhinus townsendii</i>	Townsend's Big-eared Bat	G4	S3	Blue	BG; CDF; CWH; ICH; IDF; PP	Palustrine; Subterranean; Terrestrial
<i>Martes pennanti</i>	Fisher	G5	S2S3	Blue	BAFA; BWBS; CDF; CMA; CWH; ESSF; ICH; IDF; IMA; MH; MS; PP; SBPS; SBS; SWB	Palustrine; Terrestrial
<i>Mustela erminea anguinae</i>	Ermine, <i>anguinae</i> subspecies	G5T3	S3	Blue	CDF; CWH; MH	Palustrine; Terrestrial
Marine Mammals						
<i>Eumetopias jubatus</i>	Steller Sea Lion	G3	S2S3B,S3N	Blue	CDF; CWH; MH	Estuarine; Marine
Insects						
<i>Erythemis collocata</i>	Western Pondhawk	G5	S3	Blue	BG; CDF; CWH; ESSF; PP	Lacustrine



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Sympetrum vicinum</i>	Autumn Meadowhawk	G5	S3S4	Blue	CDF; CWH	Lacustrine; Riverine
<i>Tramea lacerata</i>	Black Saddlebags	G5	S1	Red	CDF	
<i>Omus audouini</i>	Audouin's Night-stalking Tiger Beetle	G5	S1	Red	CDF; CWH	
<i>Epargyreus clarus</i>	Silver-spotted Skipper	G5	S3	Blue	CDF; CWH; ESSF; ICH; MH; MS; PP	Terrestrial
<i>Epargyreus clarus clarus</i>	Silver-spotted Skipper, <i>clarus</i> subspecies	G5T5	S3	Red	CDF; CWH	Terrestrial
<i>Erynnis propertius</i>	Propertius Duskywing	G5	S2S3	Blue	CDF; CWH	Terrestrial
<i>Euphyes vestris</i>	Dun Skipper	G5	S3	Blue	CDF; CWH; IDF	Palustrine; Terrestrial
<i>Parnassius smintheus olympiannus</i>	Rocky Mountain Parnassian, <i>olympiannus</i> subspecies	G5T4	S2S3	Blue	CDF; CWH	
<i>Euchloe ausonides insulanus</i>	Large Marble, <i>insulanus</i> subspecies	G5T1	SX	Red	CDF	Terrestrial
<i>Callophrys eryphon sheltonensis</i>	Western Pine Elfin, <i>sheltonensis</i> subspecies	G5TNR	S3	Blue	CDF; CWH	
<i>Callophrys mossii mossii</i>	Moss' Elfin, <i>mossii</i> subspecies	G4T4	S2S3	Blue	CDF; CWH	Terrestrial
<i>Plebejus saepiolus insulanus</i>	Greenish Blue, <i>insulanus</i> subspecies	G5TH	SH	Red	CDF; CWH	Terrestrial
<i>Cercyonis pegala incana</i>	Common Wood-nymph, <i>incana</i> subspecies	G5T4T5	S2	Red	CDF	Terrestrial
<i>Coenonympha tullia insulana</i>	Common Ringlet, <i>insulana</i> subspecies	G5T3T4	S1	Red	CDF; CWH	Terrestrial
<i>Danaus plexippus</i>	Monarch	G5	S3B	Blue	CDF; CWH; IDF; PP	Palustrine; Terrestrial
<i>Euphydryas editha taylori</i>	Edith's Checkerspot, <i>taylori</i> subspecies	G5T1	S1	Red	CDF	Terrestrial
<i>Copablepharon fuscum</i>	Sand-verbena Moth	G1G2	S1	Red	CDF	
Molluscs						
<i>Haliotis kamtschatkana</i>	Northern Abalone	G3G4	S2	Red	CDF; CWH	Marine
<i>Fossaria vancouverensis</i>		GHQ	SH	Red	CDF	Lacustrine



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Physella heterostropha</i>	Pewter Physa	G5Q	S1S3	Red	CDF	Lacustrine
<i>Physella integra</i>	Ashy Physa	G5	S1S3	Red	CDF	Lacustrine; Riverine
<i>Promenetus umbilicatellus</i>	Umbilicate Sprite	G4	S3S4	Blue	BG; CDF; IDF; PP	Riverine
<i>Carychium occidentale</i>	Western Thorn	G3G4	S2S3	Blue	CDF; CWH	Terrestrial
<i>Nearctula</i> sp. 1	Threaded Vertigo	G3G5	S2	Red	CDF; CWH	Terrestrial
<i>Vertigo andrusiana</i>	Pacific Vertigo	G2G3	S2	Red	CDF	Terrestrial
<i>Hemphillia dromedarius</i>	Dromedary Jumping-slug	G3G4	S2	Red	CDF; CWH	Terrestrial
<i>Hemphillia glandulosa</i>	Warty Jumping-slug	G3G4	S2S3	Blue	CDF; CWH	Palustrine; Terrestrial
<i>Prophysaon coeruleum</i>	Blue-grey Taildropper	G3G4	S1	Red	CDF; CWH	Terrestrial
<i>Prophysaon vanattae</i>	Scarletback Taildropper	G4	S3S4	Blue	CDF; CWH; MH	Terrestrial
<i>Pristiloma johnsoni</i>	Broadwhorl Tightcoil	G2G3	S2S3	Blue	CDF; CWH; MH	Terrestrial
<i>Zonitoides nitidus</i>	Black Gloss	G5	S3S4	Blue	CDF; CWH; IDF; PP	Palustrine
<i>Allogona townsendiana</i>	Oregon Forestsnail	G3G4	S1S2	Red	CDF; CWH	Terrestrial
<i>Cryptomastix devia</i>	Puget Oregonian	G3	SX	Red	CDF; CWH	Palustrine; Terrestrial
<i>Monadenia fidelis</i>	Pacific Sideband	G4G5	S3S4	Blue	CDF; CWH	Terrestrial
Ferns and allies						
<i>Botrychium simplex</i>	least moonwort	G5	S2S3	Blue	BWBSmw; CDFmm; CWHds; CWHxm; ICHmw; IDFxh; MSdk; SBPSxc	Palustrine; Riverine; Terrestrial
<i>Dryopteris arguta</i>	coastal wood fern	G5	S2S3	Blue	CDFmm	Palustrine; Terrestrial
<i>Isoetes nuttallii</i>	Nuttall's quillwort	G4?	S3	Blue	CDFmm; CWHxm	Palustrine; Riverine; Terrestrial
<i>Ophioglossum pusillum</i>	northern adder's-tongue	G5	S2S3	Blue	CDFmm; CWHvm; CWHxm; ICHmw; IDFxh	Lacustrine; Palustrine; Riverine; Terrestrial
<i>Woodwardia fimbriata</i>	giant chain fern	G5	S3	Blue	CDFmm; CWHxm	Palustrine; Riverine; Terrestrial
Dicotyledons						
<i>Abronia latifolia</i>	yellow sand-verbena	G5	S3	Blue	CDFmm; CWHvh; CWHxm	Terrestrial
<i>Anagallis minima</i>	chaffweed	G5	S2S3	Blue	CDFmm; CWHxm	Estuarine; Palustrine; Terrestrial
<i>Aster curtus</i>	white-top aster	G3	S3	Blue	CDFmm; CWHxm	Terrestrial
<i>Aster radulinus</i>	rough-leaved aster	G4G5	S1	Red	CDFmm; CWHxm	Terrestrial



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Balsamorhiza deltoidea</i>	deltoid balsamroot	G5	S1	Red	CDFmm; CWHxm	Terrestrial
<i>Bidens amplissima</i>	Vancouver Island beggarticks	G3	S3	Blue	CDFmm; CWHdm; CWHxm	Palustrine
<i>Callitriche heterophylla</i> ssp. <i>heterophylla</i>	two-edged water-starwort	G5T5	S2S3	Blue	CDFmm; CWHvm; CWHxm	Estuarine; Lacustrine; Palustrine
<i>Callitriche longipedunculata</i>	long-stalked water-starwort	G2G3	S1	Red	CDFmm	Palustrine; Terrestrial
<i>Caltha palustris</i> var. <i>palustris</i>	yellow marsh-marigold	G5T5	S2S3	Blue	CDFmm; CWHvm; CWHwm; CWHws	Estuarine; Palustrine
<i>Camissonia contorta</i>	contorted-pod evening-primrose	G5	S1	Red	CDFmm	Terrestrial
<i>Cardamine parviflora</i> var. <i>arenicola</i>	small-flowered bitter-cress	G5T5	S1	Red	CDFmm; CWHdm	Terrestrial
<i>Castilleja levisecta</i>	golden paintbrush	G1	S1	Red	CDFmm	Terrestrial
<i>Castilleja victoriae</i>	Victoria owl-clover	G1	S1	Red	CDFmm	Palustrine
<i>Centaureum muehlenbergii</i>	Muhlenberg's centaury	G5?	S1	Red	CDFmm; CWHdm	Palustrine
<i>Ceratophyllum echinatum</i>	spring hornwort	G4?	S3	Blue	CDFmm; CWHds; CWHxm	Lacustrine; Palustrine; Riverine
<i>Chamaesyce serpyllifolia</i> ssp. <i>serpyllifolia</i>	thyme-leaved spurge	G5T5	S2S3	Blue	BGxh; BGxw; CDFmm; CWHxm; IDFxh	Lacustrine; Palustrine; Terrestrial
<i>Clarkia amoena</i> var. <i>caurina</i>	farewell-to-spring	G5T5?	S3	Blue	CDFmm; CWHxm	Terrestrial
<i>Clarkia amoena</i> var. <i>lindleyi</i>	farewell-to-spring	G5T5	S3	Blue	CDFmm; CWHxm	Terrestrial
<i>Clarkia purpurea</i> ssp. <i>quadrivulnera</i>	small-flowered godetia	G5T5	S1	Red	CDFmm; CWHxm	
<i>Claytonia washingtoniana</i>	Washington springbeauty	G2G4	S2	Red	CDFmm; CWHdm; CWHxm; IDFWw	Riverine; Terrestrial
<i>Crassula aquatica</i>	pigmyweed	G5	S3	Blue	CDFmm; CWHds; ICHxw; IDFxh	Estuarine; Lacustrine; Palustrine
<i>Crassula connata</i> var. <i>connata</i>	erect pygmyweed	G5TNR	S2	Red	CDFmm	Palustrine; Riverine; Terrestrial
<i>Cuscuta campestris</i>	field dodder	G5	S2S3	Blue	BGxh; CDFmm; CWHdm; CWHxm; IDFxh	



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Elatine rubella</i>	three-flowered waterwort	G5	S2S3	Blue	BGxh; CDFmm; CWHdm; CWHxm; IDFxh	Estuarine; Lacustrine; Palustrine
<i>Epilobium ciliatum</i> ssp. <i>watsonii</i>	purple-leaved willowherb	G5T3T5	S2S3	Blue	CDFmm; CWHvh; CWHvm; CWHwh; CWHxm; ICHmw; IDFxh; SBSdh	Estuarine; Palustrine; Terrestrial
<i>Epilobium densiflorum</i>	dense spike-primrose	G5	S1	Red	CDFmm	Palustrine; Terrestrial
<i>Epilobium halleanum</i>	Hall's willowherb	G5	S2S3	Blue	BGxh; BGxw; CDFmm; ESSFdcp; ICHdw; ICHwk; SBSwk; SWBun	Lacustrine; Palustrine; Riverine; Terrestrial
<i>Epilobium torreyi</i>	brook spike-primrose	G5	SX	Red	CDFmm	Palustrine; Terrestrial
<i>Eupatorium maculatum</i> ssp. <i>bruneri</i>	Joe-pye weed	G5T4T5Q	S1	Red	CDFmm; ICHmw	
<i>Fraxinus latifolia</i>	Oregon ash	G5	S1	Red	CDFmm; CWHxm	Estuarine; Palustrine
<i>Githopsis specularioides</i>	common bluecup	G5	S2S3	Blue	CDFmm; CWHmm; CWHxm; MHmm	Palustrine; Riverine; Terrestrial
<i>Grindelia hirsutula</i> var. <i>hirsutula</i>	hairy gumweed	G5T3T4	S1	Red	CDFmm	Terrestrial
<i>Helenium autumnale</i> var. <i>grandiflorum</i>	mountain sneezeweed	G5T3T5	S2S3	Blue	BGxh; CDFmm; CWHdm; CWHxm; ICHmw; ICHxw; PPdh	Estuarine; Lacustrine; Palustrine; Terrestrial
<i>Heterocodon rariflorum</i>	heterocodon	G5	S3	Blue	CDFmm; CWHds; CWHxm; ICHdw; IDFd; IDFxh; PPdh	Palustrine; Riverine; Terrestrial
<i>Hippuris tetraphylla</i>	four-leaved mare's-tail	G5	S2S3	Blue	CDFmm; CWHds; CWHms	Estuarine; Palustrine
<i>Hutchinsia procumbens</i>	hutchinsia	G5	S1	Red	BGxh; BGxw; CDFmm; CWHxm; IDFxh	Lacustrine; Palustrine; Riverine; Terrestrial
<i>Hydrocotyle ranunculoides</i>	floating water pennywort	G5	SH	Red	CDFmm	Palustrine

Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Hydrophyllum tenuipes</i>	Pacific waterleaf	G4G5	S2	Red	CDFmm; CWHdm; CWHxm	Palustrine; Terrestrial
<i>Idahoia scapigera</i>	scalegod	G5	S2	Red	CDFmm; CWHxm; ICHxw; IDFww; PPxh	Palustrine; Riverine
<i>Jaumea carnosa</i>	fleshy jaumea	G4G5	S2S3	Blue	CDFmm; CWHvh; CWHvm; CWHxm	Estuarine
<i>Lepidium oxycarpum</i>	sharp-pod peppergrass	G4	SX	Red	CDFmm	
<i>Limnanthes macounii</i>	Macoun's meadow-foam	G2	S2	Red	CDFmm; CWHxm	Palustrine; Terrestrial
<i>Linaria canadensis</i> var. <i>texana</i>	blue toadflax	G5T4T5	S3	Blue	CDFmm	
<i>Lomatium dissectum</i> var. <i>dissectum</i>	fern-leaved desert-parsley	G4T4	S1	Red	CDFmm	Terrestrial
<i>Lomatium grayi</i>	Gray's desert-parsley	G5	S1	Red	CDFmm; CWHxm	Terrestrial
<i>Lotus formosissimus</i>	seaside birds-foot trefoil	G4	S1	Red	CDFmm	Terrestrial
<i>Lotus pinnatus</i>	bog bird's-foot trefoil	G4G5	S1	Red	CDFmm; CWHmm	Palustrine; Riverine; Terrestrial
<i>Lotus unifoliolatus</i> var. <i>unifoliolatus</i>	Spanish-clover	G5T5	S3	Blue	CDFmm; CWHxm	Terrestrial
<i>Lupinus densiflorus</i> var. <i>densiflorus</i>	dense-flowered lupine	G5T4	S1	Red	CDFmm	Riverine; Terrestrial
<i>Lupinus lepidus</i>	prairie lupine	G5	S1	Red	CDFmm; CWHmm; CWHxm	Terrestrial
<i>Lupinus oreganus</i> var. <i>kincaidii</i>	Kincaid's lupine	G5T2	SX	Red	CDFmm	Terrestrial
<i>Lupinus rivularis</i>	streambank lupine	G2G4	S1	Red	CDFmm; CWHdm; CWHxm	
<i>Marah oreganus</i>	manroot	G5	S1	Red	CDFmm; CWHxm	Terrestrial
<i>Meconella oregana</i>	white meconella	G2G3	S1	Red	CDFmm; CWHxm	Palustrine; Terrestrial
<i>Megalodonta beckii</i> var. <i>beckii</i>	water marigold	G4G5T4T5	S3	Blue	CDFmm; ICHdw; ICHmw; ICHxw; IDFdm; IDFun; SBSmk	Lacustrine; Palustrine; Riverine
<i>Microseris bigelovii</i>	coast microseris	G4	S1	Red	CDFmm	Palustrine; Terrestrial
<i>Microseris lindleyi</i>	Lindley's microseris	G5	S1	Red	CDFmm	Terrestrial
<i>Minuartia pusilla</i>	dwarf sandwort	G5	S1	Red	CDFmm	Terrestrial



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Myriophyllum quitense</i>	waterwort water-milfoil	G4?	S2S3	Blue	CDFmm; CWHvh; CWHwh; CWHxm	Lacustrine; Palustrine; Riverine
<i>Myriophyllum ussuriense</i>	Ussurian water-milfoil	G3	S3	Blue	CDFmm; CWHdm; CWHvh; ICHdw; IDFmw	Estuarine; Lacustrine; Palustrine; Riverine
<i>Navarretia intertexta</i>	needle-leaved navarretia	G5	S2	Red	CDFmm; CWHxm; ICHmk; IDFxh	Palustrine; Riverine; Terrestrial
<i>Orobancha pinorum</i>	pine broomrape	G4	S1	Red	CDFmm; CWHmm; CWHxm	Terrestrial
<i>Orthocarpus bracteosus</i>	rosy owl-clover	G3?	S1	Red	CDFmm	Palustrine; Terrestrial
<i>Plagiobothrys figuratus</i> ssp. <i>figuratus</i>	fragrant popcornflower	G4T4	S1	Red	CDFmm	Palustrine
<i>Plagiobothrys tenellus</i>	slender popcornflower	G4G5	S2	Red	CDFmm	Terrestrial
<i>Polygonum hydropiperoides</i>	water-pepper	G5	S2S3	Blue	CDFmm; CWHdm; CWHds; CWHxm	Lacustrine; Palustrine
<i>Potentilla gracilis</i> var. <i>gracilis</i>	graceful cinquefoil	G5T5	S2S3	Blue	CDFmm	
<i>Psilocarphus elatior</i>	tall woolly-heads	G4Q	S1	Red	CDFmm; CWHvh	Palustrine; Terrestrial
<i>Psilocarphus tenellus</i> var. <i>tenellus</i>	slender woolly-heads	G4T4	S3	Blue	CDFmm; CWHxm	Palustrine; Terrestrial
<i>Ranunculus alismifolius</i> var. <i>alismifolius</i>	water-plantain buttercup	G5T5	S1	Red	CDFmm	Palustrine; Terrestrial
<i>Ranunculus californicus</i>	California buttercup	G5	S1	Red	CDFmm	Terrestrial
<i>Ranunculus lobbii</i>	Lobb's water-buttercup	G4	SH	Red	CDFmm	Lacustrine; Palustrine
<i>Rubus lasiococcus</i>	dwarf bramble	G5	S2S3	Blue	CDFmm; CWHds; CWHmm; CWHxm; ESSFmw; MHmm	Terrestrial
<i>Rubus nivalis</i>	snow bramble	G4?	S2	Red	CDFmm; CWHmm; CWHvh; CWHvm; CWHxm; ICHmw; MHmm	Terrestrial
<i>Rupertia physodes</i>	California-tea	G4	S3	Blue	CDFmm; CWHmm; CWHxm; MHmm	Terrestrial



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Salix lemmonii</i>	Lemmon's willow	G5	S1	Red	CDFmm	Lacustrine; Palustrine
<i>Salix sessilifolia</i>	soft-leaved willow	G4	S2S3	Blue	CDFmm; CWHdm; CWHds; CWHvm; CWHxm	Estuarine; Lacustrine; Palustrine; Terrestrial
<i>Sanicula arctopoides</i>	snake-root sanicle	G5	S1	Red	CDFmm	
<i>Sanicula bipinnatifida</i>	purple sanicle	G5	S2	Red	CDFmm; CWHxm	Terrestrial
<i>Senecio macounii</i>	Macoun's groundsel	G5	S3	Blue	CDFmm; CWHmm; CWHxm	Estuarine; Terrestrial
<i>Sidalcea hendersonii</i>	Henderson's checker-mallow	G3	S3	Blue	CDFmm; CWHxm	Estuarine; Palustrine
<i>Silene scouleri</i> ssp. <i>grandis</i>	Scouler's catchfly	G5TNR	S1	Red	CDFmm	Terrestrial
<i>Thysanocarpus curvipes</i>	sand lacepod	G4G5	S3	Blue	CDFmm	
<i>Tonella tenella</i>	small-flowered tonella	G5	S1	Red	CDFmm	Terrestrial
<i>Toxicodendron diversilobum</i>	poison oak	G5	S2S3	Blue	CDFmm; CWHdm; CWHxm	Terrestrial
<i>Trifolium cyathiferum</i>	cup clover	G4	S1	Red	BGxh; CDFmm; CWHmm; CWHxm; ICHdw; ICHmw; IDFd; IDFxh; PPdh	Palustrine; Riverine; Terrestrial
<i>Trifolium depauperatum</i> var. <i>depauperatum</i>	poverty clover	G5T5?	S3	Blue	CDFmm; CWHxm	Palustrine; Riverine; Terrestrial
<i>Trifolium dichotomum</i>	Macrae's clover	G4?	S2S3	Blue	CDFmm; CWHxm	Terrestrial
<i>Triphysaria versicolor</i> ssp. <i>versicolor</i>	bearded owl-clover	G5T5	S1	Red	CDFmm	Palustrine; Terrestrial
<i>Utricularia ochroleuca</i>	ochroleucous bladderwort	G4?	S2S3	Blue	BWBSdk; CDFmm; ESSFmv; ICHmw	Lacustrine; Palustrine
<i>Viola howellii</i>	Howell's violet	G4	S2S3	Blue	CDFmm; CWHmm; CWHxm; MHmm	Palustrine; Riverine; Terrestrial
<i>Viola praemorsa</i> ssp. <i>praemorsa</i>	yellow montane violet	G5T3T5	S2	Red	CDFmm; CWHxm	Terrestrial
<i>Yabea microcarpa</i>	California hedge-parsley	G5?	S2	Red	CDFmm; CWHxm	Terrestrial
Monocotyledons						



Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Agrostis pallens</i>	dune bentgrass	G4G5	S3	Blue	CDFmm; CWHds; CWHmm; CWHvh; CWHvm; CWHxm	Palustrine; Terrestrial
<i>Allium amplexans</i>	slimleaf onion	G4	S3	Blue	CDFmm; CWHxm	Terrestrial
<i>Allium crenulatum</i>	Olympic onion	G4	S2	Red	CDFmm; CMA; CWHxm; MHmm	Terrestrial
<i>Allium geyeri</i> var. <i>tenerum</i>	Geyer's onion	G4G5T3T5	S2S3	Blue	BGxw; CDFmm; CWHxm; ESSFxc; IDFdk; IDFmw; IDFWw; IDFXm; MSdm	Estuarine; Palustrine; Terrestrial
<i>Alopecurus carolinianus</i>	Carolina meadow-foxtail	G5	S2	Red	CDFmm; CWHdm; CWHds; IDFXh	Palustrine; Terrestrial
<i>Bulbostylis capillaris</i>	densetuft hairsedge	G5	S1	Red	CDFmm	
<i>Carex feta</i>	green-sheathed sedge	G5	S2	Red	CDFmm; CWHxm	Palustrine; Riverine; Terrestrial
<i>Carex interrupta</i>	green-fruited sedge	G4	S2	Red	CDFmm; CWHdm	Lacustrine; Palustrine; Terrestrial
<i>Carex scoparia</i>	pointed broom sedge	G5	S2S3	Blue	BWBSmw; CDFmm; CWHdm; CWHvh; CWHxm; ICHdw; ICHwk; ICHxw; SBSvk	Lacustrine; Palustrine; Terrestrial
<i>Carex tumulicola</i>	foothill sedge	G4	S2	Red	CDFmm	Palustrine; Terrestrial
<i>Cephalanthera austinae</i>	phantom orchid	G4	S2	Red	CDFmm; CWHdm; CWHxm	Terrestrial
<i>Cyperus squarrosus</i>	awned cyperus	G5	S3	Blue	BGxh; CDFmm; CWHxm; IDFWw; IDFXh; PPxh	Lacustrine; Palustrine; Riverine; Terrestrial
<i>Eleocharis parvula</i>	small spike-rush	G5	S2S3	Blue	CDFmm; CWHvm; CWHwh; CWHxm	Estuarine; Lacustrine; Palustrine
<i>Eleocharis rostellata</i>	beaked spike-rush	G5	S2S3	Blue	CDFmm; CWHdm; CWHxm; ICHmw; IDFDm; IDFWw; MSdm; MSxk	Estuarine; Lacustrine; Palustrine; Riverine; Terrestrial

Scientific Name	English Name	Global Rank	Prov Rank	BC Status	BGC unit	Habitat Type
<i>Glyceria leptostachya</i>	slender-spiked mannagrass	G3	S2S3	Blue	CDFmm; CWHdm; CWHwh; CWHxm	Estuarine; Lacustrine; Palustrine; Terrestrial
<i>Juncus kelloggii</i>	Kellogg's rush	G3?	S1	Red	CDFmm	Palustrine
<i>Juncus occidentalis</i>	western rush	G5	S2S3	Blue	CDFmm	
<i>Juncus oxymeris</i>	pointed rush	G5	S2S3	Blue	CDFmm; CWHdm; CWHxm	Estuarine; Palustrine; Terrestrial
<i>Leymus triticoides</i>	creeping wildrye	G4G5	S1	Red	CDFmm	Terrestrial
<i>Lilaea scilloides</i>	flowering quillwort	G5?	S2S3	Blue	CDFmm; CWHdm; CWHvm; CWHxm	Estuarine; Palustrine
<i>Malaxis brachypoda</i>	white adder's-mouth orchid	G4Q	S2S3	Blue	BWBSdk; BWBSmw; CDFmm; CWHdm; CWHvm; CWHwh; CWHws; CWHxm; SBSvk	Estuarine; Lacustrine; Palustrine; Riverine; Terrestrial
<i>Melica harfordii</i>	Harford's melic	G5	S2S3	Blue	CDFmm	
<i>Piperia candida</i>	white-lip rein orchid	G3G4	S2	Red	CDFmm; CWHvh	Terrestrial
<i>Pleuropogon refractus</i>	nodding semaphoregrass	G4	S3	Blue	CDFmm; CWHdm; CWHms; CWHvh; CWHvm; CWHxm; MHmm	Palustrine; Terrestrial
<i>Potamogeton oakesianus</i>	Oakes' pondweed	G4	S2S3	Blue	CDFmm; CWHvm; ICHmw	Lacustrine
<i>Schoenoplectus americanus</i>	Olney's bulrush	G5	S1	Red	CDFmm; CWHds	Lacustrine; Palustrine; Riverine; Terrestrial
<i>Triglochin concinna</i>	graceful arrow-grass	G5	S2	Red	CDFmm; CWHvh; CWHwh	Estuarine
<i>Triteleia howellii</i>	Howell's triteleia	G3G4	S1	Red	CDFmm	Terrestrial
<i>Wolffia columbiana</i>	Columbian water-meal	G5	S1	Red	CDFmm	Lacustrine
Bryophytes						
<i>Bartramia stricta</i>	apple moss	GU	S1	Red	CDFmm	Terrestrial
<i>Entosthodon fascicularis</i>	banded cord-moss	G4G5	S2S3	Blue	CDFmm; CWHxm; ICHdm; IDFdw	Palustrine; Riverine; Terrestrial
<i>Syntrichia laevipila</i>	twisted oak moss	GNR	S2S3	Blue	CDFmm	

Search Criteria

Species Group:Plants & Animals

AND BC Conservation Status:Red (Extirpated, Endangered, or Threatened) OR Blue (Special Concern)

AND BGC Zone:CDF

Sort Order:Phylogenetic Ascending

Accessed May 31 2008





APPENDIX II

Air Photos Used for the Project

Appendix II. Air Photos Used for the Project

General Area	Flightline	Roll Number	Start Frame	End Frame	Total Photos	Flown	Scale	Source
Saltspring Island	SS-1	RC29	205	206	2	2005	1:16,500	McElhanney
Saltspring Island	SS-2	RC29	195	198	4	2005	1:16,500	McElhanney
Saltspring Island	SS-3	RC29	190	194	5	2005	1:16,500	McElhanney
Saltspring Island	SS-4	RC29	173	178	6	2005	1:16,500	McElhanney
Saltspring Island	SS-5	RC29	166	172	7	2005	1:16,500	McElhanney
Saltspring Island	SS-6	RC29	140	147	8	2005	1:16,500	McElhanney
Saltspring Island	SS-7	RC30	6	14	9	2005	1:16,500	McElhanney
Saltspring Island	SS-8	RC30	15	24	10	2005	1:16,500	McElhanney
Saltspring Island	SS-9	RC30	25	33	9	2005	1:16,500	McElhanney
Saltspring Island	SS-10	RC30	34	44	11	2005	1:16,500	McElhanney
Saltspring Island	SS-11	RC30	45	56	12	2005	1:16,500	McElhanney
Saltspring Island	SS-12	RC30	58	67	10	2005	1:16,500	McElhanney
Saltspring Island	SS-13	RC30	68	74	7	2005	1:16,500	McElhanney
Texada Island	TI-1	30BCC03037	208	216	9	2003	1:15000	ILMB
Texada Island	TI-2	30BCC03038	12	22	11	2003	1:15000	ILMB
Texada Island	TI-3	30BCC03038	24	32	9	2003	1:15000	ILMB
Texada Island	TI-4	30BCC03038	45	51	7	2003	1:15000	ILMB
Texada Island	TI-5	30BCC03038	52	58	7	2003	1:15000	ILMB
Texada Island	TI-6	30BCC03038	70	76	7	2003	1:15000	ILMB
Texada Island	TI-7	30BCC03038	77	81	5	2003	1:15000	ILMB
Texada Island	TI-8	30BCC03038	97	101	5	2003	1:15000	ILMB
Texada Island	TI-9	30BCC03038	104	108	5	2003	1:15000	ILMB
Texada Island	TI-10	30BCC03038	120	125	6	2003	1:15000	ILMB
Texada Island	TI-11	30BCC03038	134	139	6	2003	1:15000	ILMB
Texada Island	TI-12	30BCC03038	142	157	16	2003	1:15000	ILMB

Appendix II. Air Photos Used for the Project (continued)

General Area	Flightline	Roll Number	Start Frame	End Frame	Total Photos	Flown	Scale	Source
Texada Island	TI-13	30BCC03038	160	175	16	2003	1:15000	ILMB
Texada Island	TI-14	30BCC03038	176	189	14	2003	1:15000	ILMB
Texada Island	TI-15	30BCC03038	190	200	11	2003	1:15000	ILMB
Sunshine Coast	SC-1	30BCC03039	167	171	5	2003	1:15000	ILMB
Sunshine Coast	SC-2	30BCC03039	92	100	9	2003	1:15000	ILMB
Sunshine Coast	SC-3	30BCC03039	83	91	9	2003	1:15000	ILMB
Sunshine Coast	SC-4	30BCC03039	1	11	11	2003	1:15000	ILMB
Powell River	PR-1	30BCC03036	154	158	5	2003	1:15000	ILMB
Powell River	PR-2	30BCC03036	185	197	13	2003	1:15000	ILMB
Powell River	PR-3	30BCC03036	199	211	13	2003	1:15000	ILMB
Powell River	PR-4	30BCC03037	8	19	12	2003	1:15000	ILMB
Powell River	PR-5	30BCC03037	31	36	6	2003	1:15000	ILMB
Powell River	PR-6	30BCC03037	53	58	6	2003	1:15000	ILMB
Powell River	PR-7	30BCC03037	61	68	8	2003	1:15000	ILMB
Powell River	PR-8	30BCC03037	88	99	12	2003	1:15000	ILMB
Powell River	PR-9	30BCC03037	100	112	13	2003	1:15000	ILMB
Powell River	PR-10	30BCC03037	141	147	7	2003	1:15000	ILMB
Powell River	PR-11	30BCC03037	154	166	13	2003	1:15000	ILMB
Powell River	PR-12	30BCC03037	194	207	14	2003	1:15000	ILMB
Galiano Island	GI-1	RC29	202	204	3	2005	1:16,500	ILMB
Galiano Island	GI-2	RC29	199	201	3	2005	1:16,500	ILMB
Galiano Island	GI-3	RC29	185	189	5	2005	1:16,500	ILMB
Galiano Island	GI-4	RC29	179	184	6	2005	1:16,500	ILMB
Galiano Island	GI-5	RC29	157	165	9	2005	1:16,500	ILMB
Galiano Island	GI-6	RC29	148	153	6	2005	1:16,500	ILMB
Galiano Island	GI-7	RC29	134	139	6	2005	1:16,500	ILMB

Appendix II. Air Photos Used for the Project (continued)

General Area	Flightline	Roll Number	Start Frame	End Frame	Total Photos	Flown	Scale	Source
Denman Island	DI-1	354c 8	1	4	4	2001	1:20000	McElhanney
Denman Island	DI-2	354c 9	1	5	5	2001	1:20000	McElhanney
Denman Island	DI-3	354c 10	1	11	11	2001	1:20000	McElhanney
Denman Island	DI-4	354c 11	1	13	13	2001	1:20000	McElhanney
Denman Island	DI-5	354c 12	1	9	9	2001	1:20000	McElhanney
Northeast Van Isl	VI-6 #2	30BCC98008	117	119	3	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #3	30BCC98008	130	142	13	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #3	30BCC98037	91	93	3	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #4	30BCC98037	82	83	2	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #4	30BCC98008	11	29	19	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #5	30BCC98037	45	68	24	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #6	30BCC98037	112	128	17	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #7	30BCC98038	1	15	15	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #8	30BCC98038	91	99	9	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #8	30BCC98055	130	133	4	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #9	30BCC98038	100	115	16	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #10	30BCC98037	22	33	12	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #10	30BCC98038	154	158	5	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #10	30BCC98038	174	184	11	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #11	30BCC98037	13	21	9	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #11	30BCC98037	129	158	30	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #12	30BCC98036	134	157	24	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #12	30BCC98038	188	199	12	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #13	30BCC98036	107	133	27	1998	1:15000	ILMB
Northeast Van Isl	VI-6 #14	30BCC98036	87	106	20	1998	1:15000	ILMB

Appendix II. Air Photos Used for the Project (continued)

General Area	Flightline	Roll Number	Start Frame	End Frame	Total Photos	Flown	Scale	Source
Northeast Van Isl	VI-6 #15	30BCC98036	81	86	6	1998	1:15000	ILMB
Duncan/Cowichan	VI-4 #2	15BCB92131	80	95	16	1992	1:10000	ILMB
Duncan/Cowichan	VI-4 #3	15BCB92131	185	215	31	1992	1:10000	ILMB
Duncan/Cowichan	VI-4 #4	15BCB92133	165	201	37	1992	1:10000	ILMB
Duncan/Cowichan	VI-4 #5	15BCB92133	68	110	43	1992	1:10000	ILMB
Duncan/Cowichan	VI-4 #6	15BCB92132	30	83	54	1992	1:10000	ILMB
Duncan/Cowichan	VI-4 #7	15BCB92130	209	257	49	1992	1:10000	ILMB
Duncan/Cowichan	VI-4 #8	15BCB92132	84	125	42	1992	1:10000	ILMB
Duncan/Cowichan	VI-6 #7	30BCC98036	61	74	14	1998	1:15000	ILMB
Duncan/Cowichan	VI-6 #9	30BCC98038	149	153	5	1998	1:15000	ILMB
Duncan/Cowichan	VI-6 #10	30BCC98037	34	44	11	1998	1:15000	ILMB
Duncan/Cowichan	VI-6 #11	30BCC98037	1	12	12	1998	1:15000	ILMB
West Cowichan	VI-5 #2	30BCC98034	172	182	11	1998	1:15000	ILMB
West Cowichan	VI-5 #2	30BCC98034	152	158	7	1998	1:15000	ILMB
West Cowichan	VI-5 #3	30BCC98036	13	29	17	1998	1:15000	ILMB
West Cowichan	VI-5 #3	30BCC98055	155	162	8	1998	1:15000	ILMB
West Cowichan	VI-5 #4	30BCC98036	30	48	19	1998	1:15000	ILMB
Ladysmith	LS-1	15BCB86007	89	95	7	1980	1:20000	ILMB
Ladysmith	LS-2	15BCB86007	99	104	6	1980	1:20000	ILMB
Ladysmith	LS-3	15BCB86007	139	146	8	1980	1:20000	ILMB
Bamberton	BM-1	15BCB92140	215	220	6	1992	1:10000	ILMB
Bamberton	BM-2	15BCB92140	113	118	6	1992	1:10000	ILMB
Cowichan Bay	CB-1	15BCB93096	18	31	14	1992	1:10000	ILMB
Cowichan Bay	CB-2	15BCB93096	50	62	13	1992	1:10000	ILMB
Cowichan Bay	CB-3	15BCB93096	82	89	8	1992	1:10000	ILMB


Appendix II. Air Photos Used for the Project (continued)

General Area	Flightline	Roll Number	Start Frame	End Frame	Total Photos	Flown	Scale	Source
Gabriola, Thetis, Valdes, Kuper	GI-1	RC06	397	401	5	2007	1:16500	ILMB
Gabriola, Thetis, Valdes, Kuper	GI-2	RC06	390	396	7	2007	1:16500	ILMB
Gabriola, Thetis, Valdes, Kuper	GI-3	RC06	358	364	7	2007	1:16500	ILMB
Gabriola, Thetis, Valdes, Kuper	GI-4	RC06	351	357	7	2007	1:16500	ILMB
Gabriola, Thetis, Valdes, Kuper	GI-5	RC06	280	295	16	2007	1:16500	ILMB
Gabriola, Thetis, Valdes, Kuper	GI-6	RC06	262	279	18	2007	1:16500	ILMB
Gabriola, Thetis, Valdes, Kuper	GI-7	RC06	210	222	13	2007	1:16500	ILMB
Gabriola, Thetis, Valdes, Kuper	GI-8	RC06	203	209	7	2007	1:16500	ILMB
Saanich Peninsula/CRD	Saanich1	RC04	6	14	9	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich2	RC04	15	23	9	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich2	RC13	173	181	9	2005	1:8500	ILMB
Saanich Peninsula/CRD	Saanich2	RC13	3	7	5	2005	1:16500	ILMB
Saanich Peninsula/CRD	Saanich3	RC04	24	36	13	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich3	RC13	162	168	7	2005	1:16500	ILMB
Saanich Peninsula/CRD	Saanich4	RC04	37	56	20	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich5	RC04	57	75	19	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich6	RC04	76	97	22	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich7	RC04	98	106	9	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich8	RC04	107	114	8	2007	1:20000	ILMB
Saanich Peninsula/CRD	Saanich9	RC04	115	115	1	2007	1:20000	ILMB




APPENDIX III

Thematic Maps



TERRESTRIAL ECOSYSTEM MAP

POWELL RIVER NORTH
MAP 1 OF 9



Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone 1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDfmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDfmm.


The CDfmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDfmm zone.

MAP AND ECOSYSTEM LEGEND

OLD FOREST

Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Stags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.



MATURE FOREST

Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.



YOUNG FOREST

Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.



IMMATURE FOREST

Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.



GARRY OAK

Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.



NON-FORESTED

Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.



WETLAND

Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.



NON-NATURAL

Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.



BASE MAP FEATURES


- Protected Areas (Provincial & Municipal Parks)
- Crown Land (Municipal, Federal & Provincial)
- Major Roads
- CDfmm Boundary



KEY MAP 1:1,500,000



Lead Biologist: Wanda Miller, Michelle Trommelen, Pamela Williams
Lead Ecologists: Sid Tsang & Deepa Filatow
Project Managers: Jo-Anne Slacey, Carmin Cadran, Corey Erwin & Kim Everett
Data Entry and QA: Jane Thomson, Jackie Churchill & Tania Tripp
GIS/Map Production: Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Rezanoff
Contractor: Jane Thomson, Dana Luxmoore & Brett Korteling
Funding: Chartwell Consultants
Contract Monitor: ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre
Bill Zinovich, Planning Officer, ILMB



Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone 1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.


The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.

MAP AND ECOSYSTEM LEGEND

OLD FOREST

Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Snags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.



MATURE FOREST

Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.



YOUNG FOREST

Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.



IMMATURE FOREST

Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.



GARRY OAK

Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.



NON-FORESTED

Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.



WETLAND

Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.



NON-NATURAL

Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.



BASE MAP FEATURES

- Protected Areas (Provincial & Municipal Parks)
- Crown Land (Municipal, Federal & Provincial)
- Major Roads
- CDFmm Boundary



KEY MAP 1:1,500,000





Lead Biologist: Wanda Miller, Michelle Trommlen, Pamela Williams
Lead Ecologists: Sid Tsang & Deepa Filatow
Ecosystem Q/A: Jo-Anne Slacey, Carmin Cadinn, Corey Ervin & Kim Everett
Project Managers: Jane Thomson, Jackie Churchill & Tania Tripp
Data Entry and QA: Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Rezanoff
GIS/Map Production: Jane Thomson, Dana Luxmoore & Brett Korteling
Chartwell Consultants
Funding: ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre
Contract Monitor: Bill Zinovich, Planning Officer, ILMB



TERRESTRIAL ECOSYSTEM MAP

TEXADA ISLAND
MAP 2 OF 9





Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone 1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.


The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechart, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.

MAP AND ECOSYSTEM LEGEND

OLD FOREST

Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Snags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.



MATURE FOREST

Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.



YOUNG FOREST

Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.



IMMATURE FOREST

Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.



GARRY OAK

Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.



NON-FORESTED

Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.



WETLAND

Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.



NON-NATURAL

Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.



BASE MAP FEATURES

- Protected Areas (Provincial & Municipal Parks)
- Crown Land (Municipal, Federal & Provincial)
- Major Roads
- CDFmm Boundary



KEY MAP 1:1,500,000





Lead Biologist: Wanda Miller, Michelle Trommelen, Pamela Williams
Biologist Q/A: Sid Tsang & Deepa Filatow
Lead Ecologists: Helen Reid, Claudia Houwers, Jodie Krakowski & Caroline Astley
Ecologist Q/A: Jo-Anne Slacey, Carmin Cadran, Corey Ervin & Kim Everett
Project Managers: Jane Thomson, Jackie Churchill & Tania Tripp
Data Entry and QA: Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Rezanoff
GIS/Map Production: Jane Thomson, Dana Luxmoore & Brett Korteling
Montreuil Consultants: Chartwell Consultants
Funding: ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre
Contract Monitor: Bill Zinovich, Planning Officer, ILMB



TERRESTRIAL ECOSYSTEM MAP


SOUTH TEXADA ISLAND, LASQUETI ISLAND & SECHART
MAP 3 OF 9





TERRESTRIAL ECOSYSTEM MAP

DENMAN ISLAND, HORNBY ISLAND & QUALICUM
MAP 4 OF 9



Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone 1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.

The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.


MAP AND ECOSYSTEM LEGEND

OLD FOREST Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Stags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.	
MATURE FOREST Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.	
YOUNG FOREST Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.	
IMMATURE FOREST Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.	
GARRY OAK Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.	
NON-FORESTED Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.	
WETLAND Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.	
NON-NATURAL Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.	

BASE MAP FEATURES


Protected Areas (Provincial & Municipal Parks)	Major Roads
Crown Land (Municipal, Federal & Provincial)	CDFmm Boundary

0 750 1,500 3,000 4,500 6,000 7,500 Meters




Lead Biologist: Bioterrain Q/A: Lead Ecologists: Ecosystem Q/A: Project Managers: Data Entry and QA: GIS/Map Production: Montreal Consultants: Funding: Contract Monitor:	Wanda Miller, Michelle Trommelen, Pamela Williams Sid Tsang & Deepa Filatow Helen Reid, Claudia Houwers, Jodie Krakowski & Caroline Astley Jo-Anne Stacey, Carmin Cadinn, Corey Erwin & Kim Everett Jane Thompson, Jackie Churchill & Tania Tripp Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Rezanoff Jane Thompson, Dana Luxmoore & Brett Korteling Chartwell Consultants ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre Bill Zinovich, Planning Officer, ILMB
--	--

Map Revision Date: June 3, 2008



TERRESTRIAL ECOSYSTEM MAP

PARKSVILLE & NANOOSE BAY
MAP 5 OF 9



Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone 1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.

The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.


MAP AND ECOSYSTEM LEGEND

OLD FOREST Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Snags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.	
MATURE FOREST Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.	
YOUNG FOREST Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.	
IMMATURE FOREST Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.	
GARRY OAK Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.	
NON-FORESTED Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.	
WETLAND Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.	
NON-NATURAL Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.	

BASE MAP FEATURES


Protected Areas (Provincial & Municipal Parks)	Major Roads
Crown Land (Municipal, Federal & Provincial)	CDFmm Boundary

0 750 1,500 3,000 4,500 6,000 7,500 Meters



Lead Bioregion: Bioregion Q/A: Lead Ecologists: Ecosystem Q/A: Project Managers: Data Entry and QA: GIS/Map Production: Montreuil Consultants: Funding: Contract Monitor:	Wanda Miller, Michelle Trommelen, Pamela Williams Sid Tsang & Deepa Filatow Helen Reid, Claudia Houwers, Jodie Krakowski & Caroline Astley Jo-Anne Slacey, Carmin Cadim, Corey Ervin & Kim Everett Jane Thomson, Jackie Churchill & Tania Tripp Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Rezanoff Jane Thomson, Dana Luxmoore & Brett Korteling Chartwell Consultants ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre Bill Zinovich, Planning Officer, ILMB
--	---

Map Revision Date: June 3, 2008



Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone
1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.


The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.

MAP AND ECOSYSTEM LEGEND

OLD FOREST

Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Snags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.



MATURE FOREST

Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.



YOUNG FOREST

Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.



IMMATURE FOREST

Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.



GARRY OAK

Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.



NON-FORESTED

Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.



WETLAND

Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.



NON-NATURAL

Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.



BASE MAP FEATURES

- Protected Areas (Provincial & Municipal Parks)
- Crown Land (Municipal, Federal & Provincial)
- Major Roads
- CDFmm Boundary



KEY MAP 1:1,500,000





Lead Biologist: Wanda Miller, Michelle Trommlen, Pamela Williams
Lead Ecologists: Sid Tsang & Deepa Filatow
Ecosystem Q/A: Helen Reid, Claudia Houwers, Jodie Krakowski & Caroline Astley
Project Managers: Jo-Anne Slacey, Carmin Cadran, Corey Erwin & Kim Everett
Data Entry and QA: Jane Thomson, Jackie Churchill & Tania Tripp
GIS/Map Production: Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Rezanoff
Map Production: Jane Thomson, Dana Luxmoore & Brett Korteling
Chartwell Consultants
Funding: LMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre
Contract Monitor: Bill Zinovich, Planning Officer, LMB



TERRESTRIAL ECOSYSTEM MAP

NANAIMO, LADYSMITH & GABRIOLA ISLAND
MAP 6 OF 9





Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone 1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.


The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.

MAP AND ECOSYSTEM LEGEND

OLD FOREST

Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Stags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.



MATURE FOREST

Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.




YOUNG FOREST

Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.



IMMATURE FOREST

Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.



GARRY OAK

Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.



NON-FORESTED

Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.



WETLAND

Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.



NON-NATURAL

Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.



BASE MAP FEATURES

- Protected Areas (Provincial & Municipal Parks)
- Crown Land (Municipal, Federal & Provincial)
- Major Roads
- CDFmm Boundary



KEY MAP 1:1,500,000




Lead Biologist: Bioterrain Q/A: Madrone Environmental Services Ltd.
Lead Ecologists: Sid Tsang & Deepa Filatow
Ecosystem Q/A: Jo-Anne Stacey, Carmin Cadran, Corey Erwin & Kim Everett
Project Managers: Jane Thomson, Jackie Churchill & Tania Tripp
Data Entry and QA: Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Rezanoff
GIS/Map Production: Jane Thomson, Dana Luxmoore & Brett Korteling
Chartwell Consultants
Funding: ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre
Contract Monitor: Bill Zinovich, Planning Officer, ILMB



TERRESTRIAL ECOSYSTEM MAP


NORTH COWICHAN & SOUTHERN GULF ISLANDS MAP 7 OF 9





TERRESTRIAL ECOSYSTEM MAP

SOUTH COWICHAN & SAANICH PENINSULA MAP 8 OF 9



Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone
1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.

The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.

MAP AND ECOSYSTEM LEGEND

OLD FOREST
Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Snags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.

MATURE FOREST
Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.

YOUNG FOREST
Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.



IMMATURE FOREST
Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.

GARRY OAK
Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.

NON-FORESTED
Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.


WETLAND
Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.

NON-NATURAL
Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.



Lead Biologist: Wanda Miller, Michelle Trommlen, Pamela Williams
Lead Ecologists: Sid Tsang & Deepa Flatau
Ecologists: Helen Reid, Claudia Houwers, Jodie Krakowski & Caroline Astley
Project Managers: Jo-Anne Stacey, Carmin Cadim, Corey Ervin & Kim Everett
Data Entry and QA: Jane Thomson, Jackie Churchill & Tania Tripp
GIS/Map Production: Julie Cowie, Jackie Churchill, Sonia Meili, Kyle Reznoroff
Map Production: Jane Thomson, Dana Luxmoore & Brett Korteling
Chartwell Consultants
Funding: Chartwell Consultants
Contract Monitor: ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre
Bill Zinovich, Planning Officer, ILMB

Map Revision Date: June 3, 2008



Terrestrial Ecosystem Mapping of the Coastal Douglas-fir Zone 1:50,000

The Integrated Land Management Bureau (ILMB) retained Madrone Environmental Services Ltd. (Madrone) of Duncan, B.C. to conduct Terrestrial Ecosystem Mapping (TEM) for the Coastal Douglas-fir moist maritime (CDFmm) biogeoclimatic subzone. TEM consists of the stratification of landscape features into biophysical and ecological map units reflecting climate, physiography, surficial material, bedrock geology, soil, vegetation, and disturbance.

The objective of this project is to provide baseline data that can be used to support planners, scientists, and other decision makers within local, regional and provincial governments to coordinate conservation efforts, prepare sustainable growth strategies, and promote science-based, ecologically sensitive decision making around land use within the CDFmm.


The CDFmm covers approximately 252,000 hectares stretching along the east coast of Vancouver Island, the Sunshine Coast, the Gulf Islands and the Lower Mainland. The study area for this project did not include the Lower Mainland. A total of 200,000 hectares was mapped including the communities of: Powell River, Sechelt, Bowser, Qualicum, Parksville, Nanaimo, Ladysmith, Chemainus, Crofton, Cowichan, Greater Victoria and the Saanich Peninsula; the Gulf Islands: Thormanby, Savary, Hernando, Harwood, Texada, Lasqueti, Hornby, Denman, Gabriola, Valdes, Thetis, Kuper, Galiano and Salt Spring.

The thematic map displayed at left is the result of a number of queries performed on the final TEM database to classify the mapped ecosystems into eight broad categories: old forest, mature forest, young forest, immature forest, garry oak, non-forested, wetland and anthropogenic. These categories illustrate an overview perspective of the natural and non-natural ecosystems occurring within the CDFmm zone.

MAP AND ECOSYSTEM LEGEND

OLD FOREST

Old forest ecosystems are structurally complex stands comprised mainly of shade-tolerant and regenerating tree species. Snags and coarse woody debris, in all stages of decomposition, and patchy understories are common. These forests are categorized as structural stage 7 and are generally greater than 250 years in age.



MATURE FOREST

Mature forests are made up of trees that have matured since the last disturbance and are typically 80 to 250 years of age. These forests are structural stage 6 and are characterized by well developed understories and a canopy differentiated into distinct layers. A second cycle of shade tolerant trees may become established in mature forests.




YOUNG FOREST

Young forest ecosystems can begin as early as 30 years of age but typically range from 40-80 years old. These forests are structural stage 5 with self thinning evident and a forest canopy which has differentiation into distinct layers. Young forests have vigorous growth and are more open than immature forests.



IMMATURE FOREST

Immature forests are structural stage 4 and consist of trees taller than 10 m. These forests are typically dense and less than 40 years of age. Self thinning and vertical canopy structure is not evident in these forests. This category also includes recently harvested areas and other disturbed forested ecosystems.



GARRY OAK

Garry oak ecosystems include mosaics of woodlands, meadows, grasslands, scattered Douglas-fir stands, and open rocky areas. In British Columbia these ecosystems are red-listed and support high numbers of rare and endangered species of flora and fauna.



NON-FORESTED

Non-forested ecosystems are made up of herbaceous, shrubby, bryoid and sparsely vegetated units with less than 10% tree cover. They range from structural stage 1 to 3 and are generally maintained by environmental conditions or disturbance. Some of the typical non-forested units are cliffs, beaches, gravel bars, and rock outcrops.



WETLAND

Wetlands are characterized with a water table at, near, or above the surface, daily, seasonally or year-round. Soils in wetlands are water-saturated for enough periods that excess water and low soil oxygen concentrations create conditions necessary for water-tolerant plants to dominate. This unit includes fens, swamps, marshes, bogs, shallow open water and estuaries.



NON-NATURAL

Areas that have been cleared or altered permanently for human settlement or industry. These are anthropogenic units such as cultivated fields, gravel pits, mines, golf courses, orchards, vineyards, road and railway surfaces as well as rural and urban areas.



BASE MAP FEATURES

- Protected Areas (Provincial & Municipal Parks)
- Crown Land (Municipal, Federal & Provincial)
- Major Roads
- CDFmm Boundary



KEY MAP 1:1,500,000




Lead Biologist: Wanda Miller, Michelle Trommelen, Pamela Williams
Lead Ecologists: Sid Tsang & Deepa Filatow
Project Managers: Helen Reid, Claudia Houwers, Jodie Krakowski & Caroline Astley
Data Entry and QA: Jo-Anne Slacey, Carmin Cadim, Corey Erwin & Kim Everett
GIS/Map Production: Jane Thomson, Jackie Churchill & Tania Tripp
Chartwell Consultants: Jane Thomson, Dana Luxmoore & Brett Korteling
Funding: Chartwell Consultants
Contract Monitor: ILMB, The Islands Trust Fund, Ministry of Environment, The Bulkley Valley Centre
Bill Zinovich, Planning Officer, ILMB



TERRESTRIAL ECOSYSTEM MAP

SAANICH PENINSULA AND GREATER VICTORIA MAP 9 OF 9





APPENDIX IV

TEM Map Legend

TERRESTRIAL ECOSYSTEM MAPPING OF THE CDFmm SUBZONE

Map sheets: 92B/023, 032, 033, 034, 043, 044, 053, 054, 062, 063, 064, 071, 072, 073, 074, 081, 082, 083, 084, 091, 092, 093, 094, 92F/020, 028, 029, 030, 037, 038, 039, 040, 047, 048, 049, 050, 056, 057, 058, 059, 060, 066, 067, 068, 069, 077, 078, 087, 088, 096, 097, 098, 92G/001, 002, 003, 011, 012, 021, 041, 051

Scale 1: 20,000


June 2008

INTRODUCTION


This project synthesizes results of bioterrain and terrestrial ecosystem mapping of the CDFmm biogeoclimatic subzone. The CDFmm occurs in south eastern BC, covering ecosystems along the eastern coastline of Vancouver Island, the southern Gulf Islands, parts of the Sunshine Coast and a portion of the Fraser Valley. On Vancouver Island, Deep Bay marks the northern extent of the CDFmm; Metchosin marks the southern boundary. From Deep Bay moving south, the subzone extends along the Strait of Georgia from sea level to an approximate elevation of 150m above sea level (asl) and includes the major centres of Nanaimo, Duncan and Victoria. The CDFmm covers or partially covers all of the Gulf Islands south of Cortes Island; including: Texada, Hornby, Denman, Lasqueti, Gabriola Galiano, Thetis, Kuper, Saltspring, North Pender, South Pender, Mayne, Saturna, Sidney and several smaller islets in between. Across the Strait of Georgia, the CDFmm covers portions of Lund, Powell River, Sechelt and the Fraser Valley for a total area of approximately 252,000 hectares.

Digital maps will aid interpretation for resource management and land use planning; identified wildlife habitat capability and suitability; and to collate a comprehensive baseline data set of attributes of interest for the CDFmm. A seamless database of polygon attributes and the associated bioterrain and ecosystem data, as well as other features and parameters of interest accompanies this legend. Mapping was completed following the methods outlined in Standard for Terrestrial Ecosystem Mapping in British Columbia¹. Field work was completed in 2007 and 2008 at a modified survey intensity level 5.

ECOSECTION & BIOGEOCLIMATIC UNITS



ECOSYSTEM UNIT LABEL



MAP SYMBOLS

Ecosection ---

Biogeoclimatic Unit ---

Ecosystem Unit ---

Study Area Boundary ---

Plot Location •

Ecosections

SGI: Southern Gulf Islands
SGO: Strait of Georgia
NAL: Nanaimo Lowland
GEL: Georgia Lowland

Biogeoclimatic Units:

CDFmm: Coastal Douglas-Fir zone, moist maritime subzone
CWHxm1: Coastal Western Hemlock zone, very dry maritime subzone, eastern variant
CWHxm2: Coastal Western Hemlock zone, very dry maritime subzone, western variant

ECOSYSTEM UNITS

CDFmm

Site Code	Description	Site Series	Assumed Modifiers	Soil Moisture Regime	Mapped Modifiers
AS	Trembling aspen - Slough sedge	00	j, m	subhygric - subhydic	s
CD	Act - Red-osier dogwood	08	a, d, j, m	subhygric - hygric	n, s, t
CS	Cw - Slough sedge	14	d, j, m	subhydic	c, n, p, s, t, w
CW	Act - Willow	09	a, c, d, j	subhygric - hygric	

ECOSYSTEM UNITS					
CDFmm (continued)					
Site Code	Description	Site Series	Assumed Modifiers	Soil Moisture Regime	Mapped Modifiers
DA	FdPl - Arbutus	02	j, d, m, r	xeric	c, h, k, q, s, v, w, x, z
DG	FdBg - Oregon grape	04	d, j, m	subxeric - mesic	c, f, g, h, k, q, r, s, t, v, w, x, z
DO	Fd - Oniongrass	03	d, m, r	xeric	h, k, s, v, w, x
			d, j, m	subxeric - mesic	c, f, g, h, k, n, q, r, s, t, v, w, x, z
DS	Fd - Salal	01	j, m, s		c, d, h, k, n, v, w
FC	Fescue - Camas	00	j, m, r	subxeric	
GO	Garry Oak - Ocean Spray	00	j, m, v	xeric - submesic	s, v, w, z
LM	Dunegrass - Beach pea	00		submesic	c, w
LS	Pl - Sphagnum	10	d, j, p	subhydryc	
OM	Garry Oak - Moss	00	j, v	xeric	k, v, w
OR	Oceanspray - Rose	00	m, s	mesic	w
			j, m, r	xeric - submesic	h, k, s, v, w, z
QB	Garry Oak - Brome/mixed grasses	00		subhydryc	
RA	Nootka Rose - Pacific Crab Apple	00		subhydryc	
RC	Cw - Skunk cabbage	11	d, j, m		c, f, n, p, s, t
			d, j, m	subhygric - hygric	a, c, f, g, h, k, n, p, q, s, t, v, w, z
RF	CwBg - Foamflower	06	d, j, m	subhygric - hygric	c, g, h, k, n, r, s, t, v, w, x, y, z
RK	CwFd - Kindbergia	05	d, j, m		c, h, k, n, p, s, t, w
RP	Cw - Indian-plum	13	a, d, j, m	hygric	
RS	Cw - Snowberry	07		subhygric - hygric	g, k, s, t, w
RV	Cw - Vanilla-leaf	12	d, j, m	subhygric	c, h, n, p, s, v
SC	Cladina - Wallace's selaginella	00	j, m, r, v	subxeric	h, k, q, s, w, z
Ed01	Tufted hairgrass - Meadow barley	Ed01		subhydryc - hydric	
Ed03	Arctic rush - Alaska plantain	Ed03		hydric	
Em01	Widgeon-grass	Em01		subhydryc - hydric	
Em02	Glasswort - Sea milkwort	Em02		subhydryc - hydric	
Em03	Seashore saltgrass	Em03		subhydryc - hydric	
Em05	Lyngbye's sedge	Em05		subhydryc - hydric	
Wb50	Labrador tea - Bog-laurel - Peat-moss	Wb50		subhydryc	
Wf51	Sitka sedge - Peat-moss	Wf51		subhydryc	p
Wf52	Sweet gale - Sitka sedge	Wf52		subhydryc - hydric	p
Wf53	Slender sedge - White beak-rush	Wf53		hydric	d, s
Wm05	Cattail	Wm05		hydric	p
Wm06	Great bulrush	Wm06		subhydryc - hydric	
Wm50	Sitka sedge - Hemlock -parsley	Wm50		hydric	
Wm51	Three-way sedge	Wm51		subhydryc - hydric	
Ws50	Hardhack (pink spirea) - Sitka sedge	Ws50		subhydryc - hydric	p, s
Ws51	Sitka willow - Pacific willow -Skunk cabbage	Ws51		subhydryc - hydric	p
Ws52	Red alder - Skunk cabbage	Ws52		subhydryc - hydric	

CWHxm					
Site Code	Description	Site Series	Assumed Modifiers	Soil Moisture Regime	Mapped Modifiers
AM	Arbutus-Hairy manzanita	00	j, r, s	xeric	v, w, z
CS	Cw-Slough sedge	15	d, j, m a, c, d, j	subhydryc subhygric	
CW	Act-Willow (Fl50-Sitka willow-False lily-of-the-valley)	10		- hygric	
DC	FdPl-Cladina	02	j, m, r, s d, j, m	very xeric	h, k, s, v, w, z
DF	Fd-Sword fern	04		xeric - subxeric	h, k, q, s, v, w, z
DS	FdHw-Salal	03	d, m, w	xeric - subxeric	h, j, k, q, r, s, v, x, y, z
FC	Fescue-Common camas	00	j, r, s d, j, m	xeric subhygric	
HD	HwCw-Deer fern	06		- hygric	h, k, s, v, w
HK	HwFd-Kindbergia	01	d, j, m	submesic	h, k, q, r, s, v, w, x, z
QB	Garry Oak - Brome/mixed grasses	00	j, m, r	- mesic	
RB	Cw-Salmonberry	13	d, j, m	xeric - submesic	
RC	CwSs-Skunk cabbage (Ws53-Cw-Sword fern-Skunk cabbage)	12	d, j, m d, j, m	subhydryc subhygric	p, s
RF	Cw-Foamflower	07		- hygric	h, k, s, w
RS	Cw-Sword fern	05	d, m	submesic	h, j, k, q, s, v, w, x
RT	Cw-Black twinberry	14	d, j, m	- mesic	s
SC	Cladina - Wallace's selaginella	00	j, m, r, v a, d, j, m	hygric subxeric	
SS	Ss-Salmonberry	08		subhydryc	
Wb50	Labrador tea - Bog-laurel -Peat-moss	Wb50		- hygric subhydryc	
Wf53	Slender sedge - White beak-rush	Wf53		subhydryc	
Ws50	Hardhack (pink spirea) - Sitka sedge	Ws50		- hydric subhydryc	
Ws51	Sitka willow-Pacific willow-Skunk cabbage	Ws51		- hydric subhydryc	

Non-Vegetated / Sparsely Vegetated / Anthropogenic	
Site Code	Description
BE	Beach
CF	Cultivated Field
CL	Cliff
CO	Cultivated Orchard
CV	Cultivated Vineyard
DM	Dam
ES	Exposed Soil
GB	Gravel Bar
GC	Golf Course
GP	Gravel Pit
IN	Industrial
LA	Lake
MI	Mine
MU	Mudflat Sediment
OW	Shallow Open Water
PD	Pond
RE	Reservoir
RI	River
RN	Railway Surface
RO	Rock Outcrop
RZ	Road Surface
RW	Rural
TA	Talus
TZ	Mine Tailings
UR	Urban/ Suburban

SITE MODIFIERS	
Code	Topography
a	active floodplain ¹ : level or very gently sloping area bordering a river that has been formed by river erosion and deposition, with evidence of active sedimentation and deposition
g	gullying ¹ : occurs within a gully, or with gullying throughout the delineated area
h	hummocky ¹ terrain: indicated by the terrain surface expression
j	gentle slope: < 35% in the CWH and CDF zones
k	cool aspect: occurs on aspects 285°–135°, on moderately steep slopes (35%–100% in the CWH and CDF)
n	fan ¹ : occurs on a fluvial fan or on a colluvial fan or cone
q	very steep cool aspect–very steep slopes (< 100%) with aspects 285°–135°
r	ridge ¹ : occurs throughout an area of ridged terrain, or on a ridge crest
t	terrace ¹ : occurs on a fluvial, glaciofluvial, lacustrine, or rock cut terrace
w	warm aspect: 135°–285°, on moderately steep slopes (35%–100% slope in the CWH and CDF zones)
z	very steep warm aspect –slopes > 100% on aspects 135°–285°
Code	Soil
x	drier than typical
y	moister than typical
c	coarse-textured soils ² : sand and loamy sand, and sandy loam, loam, and sandy clay loam with > 70% coarse fragment volume
d	deep soil: > 100 cm to bedrock
f	fine-textured soils ² : silt and silt loam with < 20% coarse fragment volume; and clay, silty clay, silty clay loam, clay loam, sandy clay, and heavy clay with < 35% coarse fragment volume
p	peaty: on deep organics or a peaty surface (15–60 cm) ³ over mineral materials
s	shallow soils: 20–100 cm to bedrock
v	very shallow soils: < 20 cm to bedrock
STRUCTURAL STAGE	
Code	Structural Stage ¹
1	Sparse (1a) bare rock or ground / bryoid (1b) bryophytes and lichens dominant, may reflect recent disturbance
2	Herb some invading or residual shrubs and trees may be present, may reflect recent disturbance
3	Forb-dominated (2a) / Graminoid-dominated (2b) / Aquatic (2c) / Dwarf shrub (2d) Early successional stage or maintained by environmental conditions or disturbance
4	Low shrub (3a) < 2 m tall / Tall shrub (3a) 2–10 m tall
5	Pole/Sapling Trees > 10 m tall, often densely stocked, no vertical canopy structure, typically < 40 years since disturbance
6	Young Forest Self-thinning and canopy differentiation initiated, typically 40–80 years since disturbance
7	Mature Forest Mature tree canopy, typically 80–250 years since disturbance
	Old Forest Structurally complex stands comprised mainly of shade-tolerant and regenerating tree species; snags and coarse woody debris and patchy understories, typically > 250 years since disturbance.

DISTURBANCE MODIFIERS			
B	Biotic Disturbances	F	Fire disturbances
b d w k p v	<ul style="list-style-type: none"> Beaver tree cutting Domestic grazing/browsing Wildlife grazing/browsing Insects <ul style="list-style-type: none"> ⇒ Insect kill ⇒ Infestation Disease Aggressive vegetation 	c g r s i l bb pb wb	<ul style="list-style-type: none"> overstorey crown fire light surface (ground) fire repeated light surface fires severe surface fire repeated severe surface fires slash burning <ul style="list-style-type: none"> ⇒ broadcast burn ⇒ piled and burned ⇒ burned windrows
L	Forest Harvesting	L	Forest Harvesting
a c d	<ul style="list-style-type: none"> patch cut system <ul style="list-style-type: none"> ⇒ with reserves clearcut system <ul style="list-style-type: none"> ⇒ with reserves (patch retention) seed tree system <ul style="list-style-type: none"> ⇒ uniform ⇒ grouped 	e gr si st l s un gr st ir	<ul style="list-style-type: none"> selection system <ul style="list-style-type: none"> ⇒ group selection ⇒ single tree ⇒ strip land clearing Shelterwood system <ul style="list-style-type: none"> ⇒ Uniform ⇒ Group ⇒ Strip ⇒ Irregular

DATA SOURCES

This mapping project is based on a mix of monochrome and colour stereo aerial photography provided by the ILMB and the Islands Trust Fund. Airphotos ranged in scale from 1:8500 to 1:25000; photo age ranged from 1980 to 2007. Base map data is from Terrain Resource Inventory Mapping (TRIM) and provided by the Integrated Land Management Bureau (ILMB). A total of 9% polygon inspection was achieved. 78 full plots, 399 ground inspections and 985 visual checks were completed.

CREDITS

Bioterrain Mappers:	Wanda Miller, Michelle Trommelen, Pamela Williams, Sonia Meili, Brian Roberts, Gordon Butt
Bioterrain Q/A:	Sid Tsang (MoE), Deepa Filatow (MoE)
Ecosystem Mapping:	Helen Reid, Claudia Houwers, Jodie Krakowski, Chris Clement, Caroline Astley
Ecosystem Q/A:	Jo-Anne Stacey, Corey Erwin, Carmen Cadrin, Ted Lea, Kim Everett (MoE)
Field Data Collection:	Wanda Miller, Michelle Trommelen, Pamela Williams, Sonia Meili, Brian Roberts, Gordon Butt, Helen Reid, Claudia Houwers, Jodie Krakowski, Caroline Astley, Jackie Churchill, Tania Tripp
Project Manager:	Jane Thomson, assisted by Tania Tripp, & Jackie Churchill
Monorestitution:	Chartwell Consultants Ltd., Vancouver BC.
GIS/Map Production:	Jane Thomson
Funding:	Integrated Land Management Bureau, The Islands Trust Fund, Bulkley Valley Centre

LITERATURE CITED

¹Resources Inventory Committee [RIC]. 1998. Standard for terrestrial ecosystem mapping in British Columbia. Ecosystems Working Group, Terrestrial Ecosystems Task Force, Resources Inventory Committee. Vancouver, B.C. 100 pp.

²Howes, D.E. and E. Kenk (contributing eds.). 1997. Terrain classification system for British Columbia. V.2. Resource Inventory Branch, Min. Env., Lands and Parks. MOE Manual 10. Victoria, B.C. 99 pp.

³Soil Classification Working Group. 1998. The Canadian System of Soil Classification. Agric. and Agri-Food Can. Publ. 1646 (Revised) 187 pp. NRC Research Press, Ottawa, Ont.



APPENDIX V

Terrain Legend

Appendix V. Terrain Legend

MINERAL SOIL TEXTURE

c clay (<0.002 mm)	g gravel (256-2 mm)
z silt (0.062-0.002 mm)	d diamict (mixed fragments)
s sand (2-0.062 mm)	

SURFICIAL MATERIALS

C - colluvium	F^G - glaciofluvial	O - organic
D - weathered bedrock	L - lacustrine	R - bedrock
E - eolian	L^G - glaciolacustrine	W - marine
F - fluvial	M - morainal	W^A - active marine
F^A - active fluvial	N - not mapped	W^G - glaciomarine

SURFACE EXPRESSION

Simple (unidirectional) slopes

p plain, less than 5%
j gentle slope(s), 6-27%
a moderate slope(s), 28-49%
k moderately steep slope(s), 50-70%
s steep slopes(s), > 70%

Material thickness

b blanket (greater than 1m)
v veneer (less than 1 m)
w variable thickness, (0-3m)
x thin veneer (2-20cm)

Complex slopes

m rolling
u undulating
h hummocky
r ridged

Shape

c cone (slope greater than 27%)
f fan (slope less than 27%)
t terrace
d depression

GEOMORPHOLOGICAL PROCESS

Code	Definition	Code	Definition
E	channeled by meltwater	R	rapid mass movement
F	slow mass movement	V	gully erosion
H	kettled	W	washing

Mass Movement Subclasses (used with process F and/or R)

Code	Definition
"	initiation zone
b	descent of masses of bedrock by falling, bouncing & rolling
f	descent of a mass of surficial material by falling, bouncing & rolling
r	descent of masses of disintegrating bedrock by sliding
s	debris slide
u	slump of surficial material along a slip plane that is concave upward or planar.
x	combined slump (upper part) and earthflow (lower part)

SOIL DRAINAGE

Code	Definition	Code	Definition	Drainage separation	Definition
i	imperfectly drained	r	rapidly drained	,	no intermediate classes
m	moderately well drained	w	well drained	-	all intermediate classes
p	poorly drained	x	very rapidly drained		





APPENDIX VI

Expanded TEM Legend


Complete accounts for each map unit found in the study area are presented below within the expanded legend (Appendix V). A description of each ecosystem includes a site description, assumed modifiers, site characteristics, photographs and plot reference numbers. These descriptions are provided in detail and are specific to the study area each map unit and for both the CDFmm and CWHxm. Descriptions of atypical conditions are given as well where applicable.

For all forested units in the CDFmm and for forested units with plot vegetation data in the CWHxm, characteristic plant species for each ecosystem are listed by structural stage. Species lists are presented based on plot data with the exception of the species list for mature, undisturbed forested ecosystems and those ecosystems with no plot data, which are based on background information from Vancouver Region Field Guide (Green and Klinka, 1994) as well as experience from site visits. Non-forested and sparsely vegetated ecosystems are described based on a combination of plot vegetation data and background information, but not separated by structural stage. Anthropogenic units are defined based on the Standards for Terrestrial Ecosystem Mapping in BC (RISC, 1998) in association with field observations specific to each unit in the study area. Background information for classifying ecosystems was based on the Vancouver Region Field Guide (Green and Klinka, 1994) and Wetlands of BC (MacKenzie and Moran, 2004).

It must be noted that characteristic species are provided which are not necessarily indicator species. Given the time of field work (October thru February); many indicator species were not present or identifiable. Also, some ecosystems had very few plots (e.g. < 3) which limit the accuracy of the ecosystem species lists. Dominant species using plot data were determined to be all species present in an ecosystem with greater than 60% frequency and 3% cover, on average. Associate species were determined to be all species present in an ecosystem with greater than 25% frequency and 1% cover, on average.

Ecosystems that occur in the CDFmm are accompanied by a distribution map. The distribution map shows the distribution of each ecosystem based on the dominant polygon component (i.e., first decile).

Forested Site Series in the CDFmm


TEM Map Code	Site Association	CDFmm Site Series	
DS	Douglas-fir—Salal	01	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Douglas-fir - Salal ecosystems represent the regional CDFmm climate. Typical site conditions for zonal DS forests are mid to upper slope positions and gentle topography. Soils are most often moderately-well to well drained and medium-textured loamy soils with less than 35% coarse fragments. Zonal forests are dominated by Douglas-fir, in association with western redcedar. The understory is primarily dense salal, dull Oregon-grape, sword fern, and Oregon beaked moss. Less frequent species include oceanspray, red huckleberry, bracken fern, western hemlock, grand fir, step moss, and electrified cat's-tail moss. Indicator species that were recorded include orange and hairy honeysuckle and oceanspray. Forests are generally open with 30-50% canopy closure, although historic gap-driven disturbances have been diminished with the advent of more intensive forestry and fire suppression leading to younger successional forests with denser canopies. Douglas-fir in these sites is at risk of laminated root rot (<i>Phellinus weirii</i>) and, to a lesser degree, <i>Armillaria</i>. This site series is commonly found throughout the entire CDFmm. It is commonly found in complex with DA and RO on shallow soils or upper slopes and with DO on medium slopes, with medium soils.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m): <400</p> <p>Slope (%): Variable</p> <p>Aspect (°): Variable</p> <p>Surficial material: M, W^G, C</p> <p>Drainage: w</p> <p>SMR: 2-4</p> <p>SNR: A-C</p>	
		<p>Plots: 4828, 4835, 4875, 6103, 6104, 6115, 6120, 6124, 6151, 6152, 6167, 6168, 6176, 00-01811, 00-01815, 00-01817, 00-01819, 01-07832, 01-07856, 02-3005, 98-02170, 98-02171, 968967, BB_HR_G13, BB_HR_G22, BB_HR_V4, BB_HR_V21, BBJK02, BBJKV03, CA001, CA002, CA003, CA004, CA005, CA007, CA009, CA011, CA022, CA024, CAG025, CA027, CA031, CAG032, CAG033, CA034, CA035, CA036, CA037, CA038, CAG040, CA041, CA043, CAG044, CAG045, CA051, CA057, CAV058, CA060, CAG061, CA076, CAG083, CA086, CAG087, CA093, JK143, CAG102, CA103, CA107, CA108, CA110, CA115, CA118, CA119, CHG001, CHG002, CHG008, CHG014, CHG023, CHG048, CHG067, CHG076, CHG079, CHG087, CHG088, CHG101, CHG113, CHG115, CHG118, CHG123, CHG132, CHG133, CHG136, CHG138, CHG180, CHG191, CHG192, CHG199, CHG208, CHG226b, CHG230, CHG235, CHG241, CHG246, CHG264, CHG270, CHG273, CHG274, CHG283, CHG293, CORG006, CPG003, CPG008, CPG009, CPG018, CXG001, CXG004, CXG011, CXG035, DHG001, DHG024, DHG037, DHV036, WM054, WM055, CA122, CA125, GB001, GB002, GB003, JK200, TT001, GBG001, GBG002, GBG004, GB007, GB008, GB009, GB011, JK211, GBG018, GB019, GB020, GBG021, GBG023, GBV024, GB025, GB056, B_HR_V22, GB057, GB058, GB059, GB062, HRG005, HRG024, HRG036, HRG060, HRG061, HRG070, HRG075, HRG079, HRG100,</p>	


Plots (continued): HRG115, HRG138, HRG145, HRG146, HRG148, HRG157, HRG183, HRG186, HRG188, HRG201, HRG212, HRG213, HRG221, HRG226, HRG230, HRG302, HRG303. HRV011, HRV013, HRV017, HRV020, HRV021, HRV023, HRV025, HRV026, HRV027, HRV029, HRV031, HRV032, HRV049, HRV050, HRV062, HRV065, HRV076, HRV085, HRV092, HRV108, HRV112, HRV118, HRV122, HRV127, HRV132, HRV140, HRV149, HRV172, HRV176, HRV192, HRV193, HRV204, HRV205, HRV323, JCG001, JCG003, JCG013, JCG014, JCG017, JCG025, JCG042, JCG043, JCG044, JCG052, JCG054, JCG056, JCG063, JCG064, JCG067, JCG072, JCG091, JCG093, JCG094, JCG095, JCG099, JCG103, JCG105, JCG114, JCG130, JCG131, JCG132, JCG133, JCG135, JCG136, JCG137, JCG140, JCG141, JCG142, JCG145, JCG148, JCG151, JCG152, JCG163, JCG202, JCG210, JCG212, JCG228, JCG300, JCG318, JCG319, JCG322, JCG323, JCG324, JCG326, JCV039, JCV061, JCV062, JCV063, JCV064, JCV065, JK003, JKG004, JK006, JK007, JKG008, JK009, JK011, JK008, JK019, JK106, JK108, JK110, JK111, JK116, JK118, JK122, JK133, JK142, JK146, JK150, JK155, JK201, JK202, JK203, JK204, JKG223, JKG300, JK301, JK302, MTV001, MT004, MT016, MT017, MT021, MTG022, MTV023, MT024, MT040, MT042, MT044, MT045, MT046, MT047, MT049, MT051, MT054, MT055, MT056, MT059, SS001, SS002, BBJK03, SS004, TTV16, TTG001, TT005, TTG006, TT007, TT008, TTG009, TT010, TTG012, TTV17, TTV18, TTV19, TTV20, TTV21, TTV22, WM001, WM002, WM003, WMG022, WM026, WM027, WM042, ERSK_GR15, WM043, ERSK_GR13, ERSK_GR14, WM052, WM053, WM064, WM067, WM069, WM070, WM072, WM075, WM076

TEM Map Code		Site Association		CDFmm Site Series	
DS		Douglas-fir—Salal		01	
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE					
DS3 (Shrub/Herb)	DS4 (Immature Forest)	DS5 (Young Forest)	DS6 (Mature Forest)	DS7 (Old Forest)	Mature Undisturbed DS6
Dominant Species Douglas-fir salal bracken fern western redcedar	Dominant Species Douglas-fir salal Oregon beaked-moss	Dominant Species Douglas-fir salal Oregon beaked-moss dull Oregon-grape western redcedar	Dominant Species Douglas-fir Oregon beaked-moss salal western redcedar dull Oregon-grape sword fern	Dominant Species Douglas-fir salal step moss western redcedar dull Oregon-grape Oregon beaked-moss sword fern	Tree Layer Douglas-fir western redcedar grand fir bigleaf maple western flowering dogwood arbutus
Associate Species red huckleberry Oregon beaked-moss Scotch broom dull Oregon-grape sword fern trailing blackberry oceanspray	Associate Species western redcedar red alder dull Oregon-grape sword fern step moss western hemlock arbutus trailing blackberry	Associate Species sword fern step moss red huckleberry oceanspray bracken fern trailing blackberry western hemlock arbutus	Associate Species oceanspray step moss red huckleberry grand fir bigleaf maple bracken fern western hemlock	Associate Species Oregon beaked-moss sword fern lodgepole pine red huckleberry palm tree moss grand fir oceanspray trailing blackberry twinflower electrified cat's-tail moss evergreen huckleberry prickly rose western hemlock	Shrub Layer salal dull Oregon-grape oceanspray baldhip rose snowberry western trumpet honeysuckle Herb Layer vanilla-leaf bracken fern sword fern Moss Layer Oregon beaked moss step moss electrified cat's tail moss

TEM Map Code		Site Association		CDFmm Site Series	
DS		Douglas-fir—Salal		01	
DSc	coarse-textured soil	DSH	hummocky	DSrs	ridge; shallow soil
DSch	coarse-textured soil; hummocky	DSHs	hummocky; shallow soil	DSrv	ridge; very shallow soil
DSck	coarse-textured soil; cool aspect	DSHv	hummocky; very shallow soil	DSs	shallow soil
DScn	coarse-textured soil; fan	DSk	cool aspect	DSsw	shallow soil; warm aspect
DScs	coarse-textured soil; shallow soil	DSks	cool aspect; shallow soil	DSsx	shallow soil; drier than typical
DSct	coarse-textured soil; terrace	DSkv	cool aspect; very shallow soil	DSsz	shallow soil; very steep warm aspect
DScv	coarse-textured soil; very shallow soil	DSkx	cool aspect; drier than typical	DSt	terrace
DScw	coarse textured soil; warm aspect	DSn	fan	DSv	very shallow soil
DSf	fine-textured soil	DSns	fan; shallow soil	DSvx	very shallow soil; drier than typical
DSfs	fine-textured soil; shallow soil	DSq	very steep cool aspect	DSvz	very shallow soil; very steep warm aspect
DSg	gullying	DSqs	very steep cool aspect; shallow soil	DSw	warm aspect
DSgk	gullying; cool aspect	DSqv	very steep cool aspect; very shallow soil	DSx	drier than typical
DSgs	gullying; shallow soil	DSr	ridge	DSz	very steep warm aspect
<p>Atypical Characteristics/Additional Comments:</p> <p>The Douglas-fir Salal site is one of the most diverse ecosystems in the CDFmm subzone. It can occur on a variety of slope positions and a range of soil moisture regimes. Drier, submesic (2/B) sites are characterized by Douglas-fir and arbutus, virtually no grand fir or western redcedar is present. The understory is not nearly as well developed as mesic sites; there is significantly less salal and more dull Oregon-grape. Douglas-fir-Salal ecosystems were also mapped with very shallow modifiers based on the bioterrain label. Zonal sites occurred on polygons mapped as Wx or Wv and received very shallow and shallow site modifiers, respectively.</p> <p>This unit was often found on level areas of with coarse soils over glaciofluvial deposits in the Bowser, Parksville and Qualicum area. These dry, poor soils (moisture/nutrient regime consistently (2/B) contributed to poor stands of slow growing Douglas-fir mixed with shore pine. The understory consisted primarily of short dense salal, and almost no herbs.</p>					

Distribution of Map Units Containing DS Units





TEM Map Code	Site Association	CDFmm Site Series	
DA	Douglas-fir—Shore pine—Arbutus	02	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Douglas-fir – Shore pine – Arbutus ecosystems are typically found on drier, poorer sites including crests, upper slopes, warm aspects, and sites with deeper medium-textured (fine sand to silty loam) soils. Some have only a thin veneer of soil derived from glaciomarine or till origin overlying bedrock. Drainage ranges from very rapid to well. Forested ecosystems are dominated by Douglas-fir and arbutus with pine occurring on the driest and poorest sites. Garry oak also occurs as an associate species, typically growing on rock outcrops including marine-influenced reefs and islands. Forest canopies are discontinuous with large gaps, and fires are a relatively frequent historic disturbance agent. The understory has a relatively diverse layer of drought-tolerant shrubs such as hairy honeysuckle, dull Oregon grape, oceanspray and salal. Herbs are relatively sparse in these forest types. Mosses cover much of the ground and woody debris (where present) featuring electrified cat's-tail moss, step moss, Oregon beaked moss and juniper haircap moss.</p> <p>This site series are most commonly located in the southern portion of the CDFmm, the Gulf Islands, Powell River and Sechelt Area. Lasqueti Island and South Thormanby Island, Jedediah Island and Gallianno Island have the highest concentration of this ecosystem type. On Vancouver Island very few DA polygons have been mapped north of Nanoose.</p> <p>The DA unit is commonly complexed with RO and DS. In the Saanich area it is also commonly complexed with Garry oak ecosystems, QB and moss covered rock outcrops, SC.</p> <p>Assumed modifiers: d, j, m, r</p>		<p>Elevation (m): 0-500</p> <p>Slope (%): 15-60</p> <p>Aspect (°): 135-270</p> <p>Surficial material: M, R</p> <p>Drainage: w-r</p> <p>SMR: 0-1</p> <p>SNR: A-C</p>	
		<p>Plots: 4814, 6116, BBJK14, CAG002, CAG003, CAG010, CAG013, CA016, CAG019, CA028, CAG036, CAG093, CA144, CA111, CAG112, CA113, CAG116, CAG120, CA123, CHG018, CHG075, CHG110, CHG193, CHG244, CHG251, CHG259, CHG260, CHG268, CHG269, CHG271, CHG272, CHG290, CHV016, CHV181, CPG010, CXG018, DHG002, DHG006, DHG026, DHG029, DHG040, GBG001, GBG008, GBG022, GB023, GB026, GBV004, HRV136, HRV210, HRV224, HRV301, JCG138, JCG146, JKG002, JKG006, JKG010, JK117, JK119, JKG120, JK121, JK154, JKG301, JKG304, MTG019, MT057, SSG004, TTG002, TTV007, TTV15, WM066</p>	

TEM Map Code	Site Association				CDFmm Site Series
DA	Douglas-fir—Shore pine—Arbutus				02
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE					
DA3 (Shrub/Herb)	DA4 (Immature Forest)	DA5 (Young Forest)	DA6 (Mature Forest)	DA7 (Old Forest)	Mature Undisturbed DA6
Dominant Species arbutus oceanspray heron's-bill moss <i>Cladina</i> lichens Associate Species rock-moss electrified cat's-tail moss hairy manzanita lodgepole pine Douglas-fir Wallace's selaginella common snowberry kinnikinnick juniper haircap moss	Dominant Species arbutus Douglas-fir salal dull Oregon-grape oceanspray Associate Species Oregon beaked-moss red alder Scotch broom step moss sword fern	Dominant Species Douglas-fir arbutus Oregon beaked-moss Associate Species oceanspray salal western redcedar curly heron's-bill moss dull Oregon-grape Garry oak electrified cat's-tail moss step moss juniper haircap moss	Dominant Species Douglas-fir salal arbutus Associate Species step moss Oregon beaked-moss dull Oregon-grape hairy honeysuckle western redcedar heron's-bill moss juniper haircap moss Garry oak electrified cat's-tail moss Wallace's selaginella oceanspray	Dominant Species Douglas-fir electrified cat's-tail moss step moss arbutus Oregon beaked-moss Associate Species oceanspray broom-moss heron's-bill moss grey rock-moss Wallace's selaginella false-polytrichum Garry oak Scotch broom lodgepole pine broad-leaved stonecrop	Tree Layer Douglas-fir arbutus Garry oak Shrub Layer oceanspray dull Oregon-grape baldhip rose western trumpet honeysuckle snowberry hairy honeysuckle falsebox salal Herb Layer Alaska oniongrass purple peavine sword fern big-leaved sandwort Pacific sanicle Moss Layer electrified cat's tail moss Oregon beaked moss step moss lichen.

TEM Map Code		Site Association		CDFmm Site Series	
DA		Douglas-fir—Shore pine—Arbutus		02	
DAcs	coarse-textured soil; shallow soil	DAk	cool aspect	DAsw	shallow soil; warm aspect
DAcv	coarse-textured soil; very shallow soil	DAks	cool aspect; shallow soil	DAsz	shallow soil; very steep warm aspect
DAcw	coarse textured soil; warm aspect	DAkv	cool aspect; very shallow soil	DAv	very shallow soil
DAh	Hummocky	DAq	very steep cool aspect	DAvw	very shallow soil; warm aspect
DAhs	hummocky; shallow soil	DAqs	very steep cool aspect; shallow soil	DAvz	very shallow soil; very steep warm aspect
DAhv	hummocky; very shallow soil	DAqv	very steep cool aspect; very shallow soil	DAw	warm aspect
DAhw	hummocky; warm aspect	DAs	shallow soil	DAX	drier than typical
Atypical Characteristics/Additional Comments: The DA unit is most commonly found on shallow to very shallow soils in the CDFmm. Soils are often coarse textured and well drained. These sites are often steep, in which case they are predominantly south-facing, although north-facing do also occur.					

Distribution of Map Units Containing DA Units




TEM Map Code	Site Association	CDFmm Site Series	
DO	Douglas-fir—Oniongrass	03	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Douglas-fir - Oniongrass ecosystems are predominate in rich to very rich xeric to subxeric sites (crests to upper slopes) where bedrock is near the surface or forms outcrops. They often contain or form mosaics with herbaceous meadows containing Garry oak where soils are deeper; sites that harbour high plant, vertebrate and invertebrate diversity. The tree layer is dominated by Douglas-fir; Garry oak is present on warm aspects with mull humus forms; arbutus is typically the only other tree species. The shrub layer is sparse with small amounts of oceanspray and hairy honeysuckle and regeneration of the overstory species. The herb layer is diverse with many annual and perennial species such as blue wildrye, orchard-grass, Pacific sanicle, quackgrass, and cleavers. The bryophyte layer is dominated by electrified cat's-tail moss and Oregon beaked moss.</p> <p>This unit is relatively rare in the CDF and distribution is primarily restricted to Saltspring Island, the Cowichan Valley, and Saanich.</p> <p>The DO ecosystem is typically mapped in complex with DS and DA.</p> <p>Assumed modifiers: d, m, r</p>		<p>Elevation (m) <550</p> <p>Slope (%) 15-50</p> <p>Aspect (°): 100-250</p> <p>Surficial material W^G, M</p> <p>Drainage r-w</p> <p>SMR 0-1</p> <p>SNR D-E</p>	
		<p>Plots: 4813, 6117, 9629191, CA010, CA049, CAG076, CA120, CHG038, CHG223, CHG245, GBG003, GB004, GB010, HRG182, HRG185, HRG228, JK207, JK208, JK209, JK210, JK217, JK218, MT003, MT041, MT043, QAG004, TTV003, WM063</p>	

TEM Map Code	Site Association				CDFmm Site Series
DO	Douglas-fir—Oniongrass				03
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE					
DO3 (Shrub/Herb)	DO4 (Immature Forest)	DO5 (Young Forest)	DO6 (Mature Forest)	DO7 (Old Forest)	Mature Undisturbed DO6
Dominant Species Scotch broom long rock-moss sidewalk moss arbutus	Dominant Species Douglas-fir Oregon beaked-moss curly heron's-bill moss	Dominant Species Douglas-fir arbutus electrified cat's-tail moss oceanspray curly heron's-bill moss Oregon beaked-moss	Dominant Species Douglas-fir electrified cat's-tail moss arbutus Garry oak blue wildrye heron's-bill moss hairy honeysuckle Oregon beaked-moss Scotch broom	Dominant Species Douglas-fir electrified cat's-tail moss Garry oak heron's-bill moss Oregon beaked-moss oceanspray Pacific sanicle quackgrass cleavers	Tree Layer Douglas-fir Garry oak Shrub Layer hairy honeysuckle snowberry Herb Layer Alaska oniongrass Pacific sanicle big-leaved sandwort broad-leaved shootingstar nodding trisetum Moss Layer electrified cat's tail moss Oregon beaked moss
Associate Species Douglas-fir dull Oregon-grape bigleaf maple dovefoot geranium hairy manzanita oceanspray	Associate Species western hemlock sweet vernalgrass Garry oak long-stoloned sedge oceanspray Alaska oniongrass gorse juniper haircap moss Scotch broom sedge starwort step moss bedstraw electrified cat's-tail moss	Associate Species dull Oregon-grape salal Garry oak bigleaf maple baldhip rose variable moss	Associate Species dull Oregon-grape oceanspray bigleaf maple kinnikinnick salal western redcedar grand fir western yew clover orchard-grass Kentucky bluegrass ragged-moss	Associate Species broad-leaved stonecrop clad lichens common snowberry blue wildrye dull Oregon-grape orchard-grass Scotch broom spurge-laurel sword fern	

TEM Map Code		Site Association		CDFmm Site Series	
DO		Douglas-fir—Oniongrass		03	
DOh	hummocky	DOkv	cool aspect; very shallow soil	DOvw	very shallow soil; warm aspect
DOhv	hummocky; very shallow soil	DOs	shallow soil	DOw	warm aspect
DOk	cool aspect	DOsw	shallow soil; warm aspect	DOwx	warm aspect; drier than typical
DOKs	cool aspect; shallow soil	DOv	very shallow soil		
Atypical Characteristics/Additional Comments: DO if often found on shallow to very shallow soils and typically on warmer aspects.					

Distribution of Map Units Containing DO Units





TEM Map Code	Site Association	CDFmm Site Series	
DG	Douglas-fir—Grand fir—Oregon grape	04	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Richer than zonal sites, Douglas-fir – Grand fir – Oregon grape ecosystems occur mostly on middle to upper sites on gently sloping ground with deep, medium-textured soil, indicated by the absence of arbutus which may be an incidental on some sites.</p> <p>Douglas-fir and western redcedar are the dominant tree species. Bigleaf maple, grand fir, western hemlock and red alder are generally present. In the understory, salal, dull Oregon-grape, sword fern and Oregon beaked moss are dominant.</p> <p>Younger stands have denser canopies due to the varying shade tolerance of the component species and sparse understories until canopy breakup occurs later in structural stage 5.</p> <p>DG is uncommon in the CDFmm and is mainly found on the Gulf Islands, including Denman, Hornby, Lasqueti, Texada, Gabriola, Valdez and Saltspring Islands. Several mapped occurrences are also located in Saanich, the Cowichan Valley, Ladysmith, Nanoose, Nanaimo and Bowser areas.</p> <p>DG is generally mapped in complex with DS or DA.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m)</p>	<150
		<p>Slope (%)</p>	0-20 (<35)
		<p>Aspect (°):</p>	Variable
		<p>Surficial material</p>	W ^G , M
		<p>Drainage</p>	w
		<p>SMR</p>	2-4
		<p>SNR</p>	D-E
		<p>Plots: 4833, 4873, 6101, 6106, 6150, 6190, 6191, 08094, 00-01874, 01-07855, 02-043, BB_HR_V12, G13, BBJKG02, BBJK10, BBJK12, CA013, CA014, CA015, CA033, CAG039, CA042, CA046, CA050, CA084, CA096, CA104, CAG105, CA106, CA117, CHG060, CHG062, CHG063, CHV061, CXG002, CXG029, DHG012, DHV027, WM057, GB005, GBG006, GB012, GB033, JK021, GBG041, GB043, GB045, BB_HR_V4, V5, GBG047, GB048, HRG004, HRG056, HRG184, HRG199, HRG202, HRV218, JCG078, JCG098, JCG100, JCG134, JCG168, JCG224, JCG231, JK001, JK004, JK023, JK101, JK114, JK124, JK132, WM077, JK139, JK140, JK149, JK205, JK221, JK223, JK224, MTG006, MTG009, MT010, MT018, MT033, MTV046, MT048, MT053, SS003, TTG12, WMG024, WMG026, WM068, WM071</p>	

TEM Map Code	Site Association		CDFmm Site Series
DG	Douglas-fir—Grand fir—Oregon grape		04
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE			
DG4 (Immature Forest)	DG5 (Young Forest)	DG6 (Mature Forest)	Mature Undisturbed DG6
Dominant Species Douglas-fir sword fern trailing blackberry dull Oregon-grape western redcedar salal	Dominant Species Douglas-fir sword fern Oregon beaked-moss western redcedar red alder bigleaf maple salal	Dominant Species Douglas-fir sword fern western redcedar salal dull Oregon-grape	Tree Layer Douglas-fir western redcedar grand fir bigleaf maple western flowering dogwood
Associate Species red alder salmonberry slender beaked-moss grand fir oceanspray western hemlock tangle moss pipecleaner moss Oregon beaked-moss	Associate Species dull Oregon-grape grand fir oceanspray trailing blackberry salmonberry western hemlock	Associate Species Oregon beaked-moss bigleaf maple red alder grand fir western hemlock	Shrub Layer salal dull Oregon-grape oceanspray baldhip rose
			Herb Layer sword fern vanilla-leaf bracken fern three-leaved foamflower
			Moss Layer Oregon beaked moss step moss palm tree moss electrified cat's tail moss

TEM Map Code		Site Association		CDFmm Site Series	
DG		Douglas-fir—Grand fir—Oregon grape		04	
DGc	coarse-textured soil	DGh	hummocky	DGsw	shallow soil; warm aspect
DGck	coarse-textured soil; cool aspect	DGhs	hummocky; shallow soil	DGsx	shallow soil; drier than typical
DGcs	coarse-textured soil; shallow soil	DGhv	hummocky; very shallow soil	DGt	terrace
DGcw	coarse textured soil; warm aspect	DGk	cool aspect	DGv	very shallow soil
DGf	fine-textured soil	DGks	cool aspect; shallow soil	DGvw	very shallow soil; warm aspect
DGg	gullying	DGkv	cool aspect; very shallow soil	DGw	warm aspect
DGgk	gullying; cool aspect	DGq	very steep cool aspect	DGwx	warm aspect; drier than typical
DGgs	gullying; shallow soil	DGrs	ridge; shallow soil	DGz	very steep warm aspect
DGgw	gullying; warm aspect	DGs	shallow soil		
Atypical Characteristics/Additional Comments: DG was frequently mapped on south-facing slopes, although north-facing slopes were also mapped. This ecosystem was commonly found on shallow to very shallow coarse soils, particularly on the Gulf Islands.					


Distribution of Map Units Containing DG Units




TEM Map Code	Site Association	CDFmm Site Series	
RK	Western redcedar—Douglas-fir—Oregon beaked moss	05	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar – Douglas-fir – Oregon beaked moss ecosystems have more available moisture than zonal sites (slightly dry to fresh), and are typically situated on lower gentle sloping (moisture receiving) sites. Soils are typically deep and often imperfectly to moderately-well drained.</p> <p>Douglas-fir and western redcedar are typically the most abundant, but bigleaf maple, red alder and western hemlock can also occur. The shrub layer is dominated by salal and dull Oregon grape, with trailing blackberry also occurring. Sword fern is dominant in the herb layer. Step moss and Oregon beaked moss are the major bryophytes.</p> <p>This unit is not common throughout the CDF. However, it does occur on Saltspring, Texada, and Thetis Islands, as well as in the Cowichan Valley, north to Ladysmith. RK ecosystems are also mapped in the Powell River, Lasqueti Island, Denman Island, Galiano Island, Bowser and on the Sannich Peninsula, but are less frequent in these areas.</p> <p>These moist forests are generally found complexed with zonal (DS) forest or moist richer forests RF.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m) <350</p> <p>Slope (%) 0-50</p> <p>Aspect (°) variable</p> <p>Surficial material M, W^G (C)</p> <p>Drainage w-m</p> <p>SMR 5-6</p> <p>SNR A-C</p>	
		<p>Plots: 4874, 6119, 6149, BB_HR_G11, BBJKV05, BBJKV07, BBJK08, BBJKG09, BBJK13, CAG001, CA047, CA048, CAG054, CA065, CA066, CAG068, CAG072, CA077, CA078, CA081, CA091, CA092, CAG095, CA097, CAG098, CA109, CHG094, CORG001, CPG015, CXG024, DHG003, DHG007, GB015, GB016, GB017, GB027, GB032, GB041, GB046, GBV049, GBG050, GBG051, BBJK06, GB060, GB061, GB063, GB064, HRG030, HRG081, HRV016, HRV151, JCG005, JKG002, JK126, JK134, JK144, JK152, JK156, JKG210, SS005, SS006, JK206, JK213, JKG215, MT006, MT019, MT020, MT025, MT030, MT031, MT032, MTG034, MT052, TTV015, WMG029, WM047, WM048, WM024, WM046, WM049, WM058, WM059, WM061, WM074</p>	

TEM Map Code	Site Association		CDFmm Site Series
RK	Western redcedar—Douglas-fir—Oregon beaked moss		05
RKc	coarse-textured soil	RKkn	cool aspect; fan
RKcw	coarse textured soil; warm aspect	RKks	cool aspect; shallow soil
RKg	gullying	RKkv	cool aspect; very shallow soil
RKgk	gullying; cool aspect	RKn	fan
RKgw	gullying; warm aspect	RKr	ridge
RKh	hummocky	RKs	shallow soil
RKhs	hummocky; shallow soil	RKsw	shallow soil; warm aspect
RKk	cool aspect	RKt	terrace
		RKv	very shallow soil
		RKvw	very shallow soil; warm aspect
		RKvx	very shallow soil; drier than typical
		RKvy	very shallow soil; moister than typical
		RKw	warm aspect
		RKz	very steep warm aspect
Atypical Characteristics/Additional Comments: RK forests are commonly found on shallow to very shallow coarse-textured soils. These sites can be found on slopes, steep in some cases, which have either cool or warm aspects.			

Distribution of Map Units Containing RK Units




TEM Map Code	Site Association	CDFmm Site Series	
RF	Western redcedar—Grand fir—Foamflower	06	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar – Grand fir - Foamflower ecosystems are found in similar terrain and slope positions to site series 05 (gentle lower slopes and moisture receiving sites), but soils tend to be deeper and richer, with a higher proportion of fines (e.g., loamy silt, silty clay loam).</p> <p>Dominant tree species include western redcedar, Douglas-fir, bigleaf maple, red alder and grand fir. The canopy tends to have more complete closure and ingress of western redcedar, with few, shade-tolerant shrubs and herbs persisting in the understory – primarily sword fern, red huckleberry, salal and dull Oregon grape. The most common mosses in these forest types are Oregon beaked moss and slender beaked-moss.</p> <p>This ecosystem is common throughout the CDFmm, with the exception of Saltspring Island and Gabriola Island, where notably fewer units are present. The highest concentration of the RF unit is on Lasqueti Island, Powell River, Sechelt and the western portion of the Cowichan Valley, north to Ladysmith.</p> <p>The RF unit is often found complexed with zonal (DS) forests and adjacent to streams and rivers (RI).</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m) 0-250</p> <p>Slope (%) 0-35</p> <p>Aspect (°) Variable</p> <p>Surficial material W^G, M</p> <p>Drainage m</p> <p>SMR 5-6</p> <p>SNR D-E</p>	
		<p>Plots: 4868, 4876, 6139, 6140, 6153, 6159, 6192, 00-01810, 00-01812, 00-01813, 00-0186, 00-01818, 00-01876, 02-3155, BB_HR_G7, BB_HR_V20, BBJKG04, GB053, CA026, CA032, CAG056, CA068, CA124, CHG019, CHG033, CHG039, CHG077, CHG083, CHG105, CHG122, CHG137, CHG207, CHG218, CHG228, CHG240, CHG252, CHG255, CHG294, CORG004, CXG037, DHG023, GBG009, GB014, GB036, GB039, GBG046, GBG053, GBV006, GBV014, GBV017, GBV048, HRG038, HRG047, HRG089, HRG139, HRG142, HRG143, HRV064, HRV082, HRV083, HRV091, HRV095a, HRV097, HRV147, HRV160, JCG011, JCG074, JCG139, JCG154, JCG214, JCG230, JK103, JK109, JK123, JK129, JK130, JK147, JK151, WM019, WM020, JK215, JK216, TT011, TTV013, WMG014, WMG015, WM018, WM040</p>	

TEM Map Code	Site Association			CDFmm Site Series
RF	Western redcedar—Grand fir—Foamflower			06
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE				
RF4 (Immature Forest)	RF5 (Young Forest)	RF6 (Mature Forest)	RF7 (Old Forest)	Mature Undisturbed RF6
Dominant Species red alder sword fern western redcedar Oregon beaked-moss	Dominant Species sword fern red alder western redcedar bigleaf maple	Dominant Species sword fern western redcedar red alder bigleaf maple Douglas-fir grand fir	Dominant Species western redcedar sword fern Douglas-fir palm tree moss bigleaf maple slender beaked-moss	Tree Layer Douglas-fir western redcedar grand fir western flowering dogwood bigleaf maple
Associate Species Douglas-fir coastal leafy moss western hemlock bigleaf maple salal dull Oregon-grape salmonberry	Associate Species Douglas-fir Oregon beaked-moss salmonberry lanky moss dull Oregon-grape red huckleberry salal grand fir western hemlock trailing blackberry	Associate Species Oregon beaked-moss dull Oregon-grape salal red huckleberry western hemlock salmonberry slender beaked-moss trailing blackberry leafy moss	Associate Species Oregon beaked-moss salal grand fir red alder leafy moss red huckleberry dull Oregon-grape one-leaved foamflower English holly salmonberry	Shrub Layer dull Oregon-grape salal
				Herb Layer three-leaved foamflower sword fern vanilla-leaf
				Moss Layer Oregon beaked moss step moss palm tree moss electrified cat's tail moss

TEM Map Code		Site Association		CDFmm Site Series	
RF		Western redcedar—Grand fir—Foamflower		06	
RFa	active floodplain	RFgs	gullying; shallow soil	RFq	very steep cool aspect
RFc	coarse-textured soil	RFgw	gullying; warm aspect	RFs	shallow soil
RFcn	coarse-textured soil; fan	RFh	hummocky	RFst	shallow soil; terrace
RFcs	coarse-textured soil; shallow soil	RFhs	hummocky; shallow soil	RFsw	shallow soil; warm aspect
RFct	coarse-textured soil; terrace	RFk	cool aspect	RFt	terrace
RFcw	coarse textured soil; warm aspect	RFks	cool aspect; shallow soil	RFvw	very shallow soil; warm aspect
RFf	fine-textured soil	RFn	fan	RFw	warm aspect
RFfs	fine-textured soil; shallow soil	RFns	fan, shallow soil	RFz	very steep warm aspect
RFg	gullying	RFp	peaty material		
RFgk	gullying; cool aspect	RFps	peaty material; shallow soil		
<p>Atypical Characteristics/Additional Comments:</p> <p>This unit is commonly found on shallow, coarse-textured soils. The RF unit is associated with streams and rivers and is commonly found on the slopes on both sides of the water courses. These slopes are often greater than 35% resulting in aspect modifier “k or w” being applied to the ecosystem unit. These water courses are often gullied and the modifier, “g” is commonly applied to these units.</p> <p>Several RF units in the Bowser area and on Texada Island were found on shallow to deep, organic, peaty soils “p”. These sites are generally hummocky and the herbaceous layer is generally sparse, except in wetter pockets where skunk cabbage and lady fern are found.</p> <p>Young ecosystems often contain a high percentage of red alder in the canopy and have a dense shrub layer of moisture loving species such as salmonberry.</p>					

Distribution of Map Units Containing RF Units




TEM Map Code	Site Association	CDFmm Site Series	
RS	Western redcedar—Snowberry High Bench Floodplain	07	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar – Snowberry ecosystems occur on high bench floodplains. The moderately well-sorted deep fluvial deposits upon these sites are moderately well to well drained. Forests are infrequently inundated and support species adapted to periodic flooding or high water tables.</p> <p>Western redcedar is the most frequent conifer, red alder and bigleaf maple frequent components, and black cottonwood less frequent. The understory is dominated by salmonberry, stink currant, piggy-back plant, lady fern and Cooley's hedge-nettle. The moss layer is dominated by slender beaked-moss.</p> <p>This ecosystem is rare in the CDF and distribution is restricted to several sites in Powell River, Saltspring Island and from Parksville to the Saanich Peninsula. The Cowichan and Chemainus Rivers in the Cowichan Valley had the most extensive high bench floodplain sites.</p> <p>The RS unit is adjacent to rivers and streams and is most commonly complexed with the moist rich forest RF ecosystem as well as the medium and low bench floodplain units, CD and CW, respectively.</p> <p>Assumed modifiers: a, d, j, m</p>		Elevation (m)	50-150
		Slope (%)	0-35
		Aspect (°)	variable
		Surficial material	F ^A p, W ^G b
		Drainage	m-w
		SMR	5-7
		SNR	D-E
		Plots: 00-01807, CORG008, CHV130, CHV164	

TEM Map Code	Site Association	CDFmm Site Series
RS	Western redcedar—Snowberry High Bench Floodplain	07
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE		
RS5 (Young Forest)		Mature Undisturbed RS6
Dominant Species red alder salmonberry western redcedar piggy-back plant bigleaf maple lady fern Cooley's hedge-nettle Associate Species blue wildrye stink currant slender beaked-moss Robert's geranium sword fern stinging nettle skunk cabbage sedge Douglas-fir		Tree Layer red alder bigleaf maple western redcedar black cottonwood Shrub Layer salmonberry red elderberry devil's club stink currant Herb Layer piggy-back plant sword fern vanilla-leaf star-flowered false Soloman's-seal blue wildrye lady fern three-leaved foamflower false lily-of-the-valley mountain sweet-cicely Moss Layer coastal leafy moss palm tree moss slender beaked moss

TEM Map Code	Site Association	CDFmm Site Series
RS	Western redcedar—Snowberry High Bench Floodplain	07
RSg gullying RSgw gullying; warm aspect RSk cool aspect RSs shallow soil RSt terrace Atypical Characteristics/Additional Comments: Rivers and streams are often associated with gullying. Therefore, the RS ecosystems adjacent to these waterbodies are often mapped within steep-sided gullies.		

Distribution of Map Units Containing RS Units




TEM Map Code	Site Association	CDFmm Site Series	
CD	Black cottonwood—Red-osier dogwood Medium Bench Floodplain	08	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Black cottonwood – Red-osier dogwood ecosystem occurs on mid bench active floodplain sites. This deciduous forest has an open canopy with a well-developed shrub understory. Red alder, bigleaf maple, western redcedar, and Douglas-fir are the dominant tree species. The understory is dominated by shrubs, including a high cover of red-osier dogwood, salmonberry and common snowberry. The moderately well-developed herb layer is often characterized by sword fern, three-leaved foamflower, and lady fern. The moderately well-developed but floristically simple moss layer is dominated by high cover of coastal leafy moss and palm tree moss.</p> <p>This floodplain unit is very rare the CDFmm but is found on the Vancouver Island portion of the study area, including the Qualicum/Parksville area, Cowichan Valley and Saanich. Most of the CD units are found on the Qualicum, Englishman, Chemainus and Cowichan Rivers.</p> <p>The CD unit is adjacent to rivers and streams and is most commonly complexed with the high bench floodplain unit, RS.</p> <p>Assumed modifiers: a, d, j, m</p>		Elevation (m) 5-100 Slope (%) 0 Aspect (°) 999 Surficial material F ^A p, Fv Drainage m-w SMR 5-6 SNR D	
		Plots: HRG055, HRG187, HRV189, CHG032, CHG034, CXG012	

TEM Map Code	Site Association			CDFmm Site Series
CD	Black cottonwood—Red-osier dogwood Medium Bench Floodplain			08
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE				
CD3 (Shrub/Herb)	CD5 (Young Forest)	CD6 (Mature Forest)	Mature Undisturbed CD6	
Dominant Species red alder red-osier dogwood reed canarygrass willow Nootka rose Himalayan blackberry	Dominant Species red alder trembling aspen bigleaf maple salmonberry	Dominant Species palm tree moss salmonberry bigleaf maple Douglas-fir lady fern western redcedar creeping buttercup piggy-back plant red alder sedge sword fern three-leaved foamflower common horsetail	Tree Layer red alder bigleaf maple western redcedar grand fir Douglas-fir Shrub Layer indian-plum salmonberry red elderberry common snowberry Herb Layer sword fern three-leaved foamflower	
Associate Species black hawthorn western redcedar	Associate Species red-osier dogwood bluejoint reedgrass western hemlock common snowberry narrow-leaved everlasting peavine Nootka rose western redcedar sword fern common burdock	Associate Species coastal leafy moss Cooley's hedge-nettle deer fern northern maiden-hair oceanspray red huckleberry	Moss Layer slender beaked moss coastal leafy moss palm tree moss	

TEM Map Code	Site Association	CDFmm Site Series
CD	Black cottonwood—Red-osier dogwood Medium Bench Floodplain	08
CDn fan CDs shallow soil CDt terrace Atypical Characteristics/Additional Comments: N/A		

Distribution of Map Units Containing CD Units




TEM Map Code	Site Association	CDFmm Site Series	
CW	Black cottonwood—Willow Low Bench Floodplain	09	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Black cottonwood-Willow ecosystem occurs on low bench active floodplain sites. It typically features deep coarse-textured gravelly sandy soils with relatively high coarse fragment content. Frequent flooding events and inundation supports species such as red alder, willows and black cottonwood, with associated flood-tolerant shrubs such as salmonberry. Herbs and mosses are infrequent or absent, with blue wildrye most common. Soils are typically coarse and well drained however frequent flooding action precludes the development of soil and humus development.</p> <p>These low bench floodplain sites are rare in the CDF. They are mapped on Texada and Saltspring Islands as well as in South Nanaimo on the Nanaimo River, in the Cowichan Valley on the Chemainus and Cowichan Rivers and in Saanich on Goldstream.</p> <p>The CW unit is found adjacent to streams and rivers and is most commonly complexed with the floodplain unit, RS.</p> <p>Assumed modifiers: a, c, d, j</p>		<p>Elevation (m)</p>	<150
		<p>Slope (%)</p>	<15
		<p>Aspect (°)</p>	variable
		<p>Surficial material</p>	F ^A
		<p>Drainage</p>	m-w
		<p>SMR</p>	5-6
		<p>SNR</p>	D
		Plots: GBG005, JK136	

TEM Map Code	Site Association	CDFmm Site Series
CW	Black cottonwood—Willow Low Bench Floodplain	09
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE		
CW4 (Immature Forest)		Mature Undisturbed CW6
Dominant Species Douglas-fir grand fir dull Oregon-grape Oregon beaked-moss Associate Species oceanspray trailing blackberry bracken fern Indian-plum red huckleberry salmonberry saskatoon sword fern vanilla-leaf	Tree Layer red Alder black cottonwood Shrub Layer willow sp. salmonberry red elderberry red-osier dogwood Herb Layer blue wildrye scouring-rush common horsetail Moss Layer electrified cat's-tail moss	

Distribution of Map Units Containing CW Units




TEM Map Code	Site Association	CDFmm Site Series	
LS	Shore pine—Sphagnum	10	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>This Shorepine-Sphagnum ecosystem is usually situated in basin depressions where the water table is high. Water flow is virtually stagnant creating nutrient-poor conditions. The LS ecosystem is rare in the CDFmm. The tree canopy is typically very open with scattered shore pine. The shrub understory is dense, but dominated mainly by Labrador tea and salal. The moss layer is dominated by step moss, red-stemmed feathermoss, peat-moss and Oregon beaked-moss.</p> <p>This unit is not common in the CDFmm. It is most abundant in the Bowser/Qualicum/Parksville area and on Lasqueti Island. Several sites are mapped in the Powell River area and associated Islands, Sechelt, Nanaimo, as well as Gabriola, Valdez, Galiano and Saltspring Islands.</p> <p>The LS unit is most commonly complexed with the hardhack dominated shrub unit, Ws50. Western redcedar swamp forests (RC) commonly occur adjacent to these depressions, and are often complexed with LS.</p> <p>Assumed modifiers: d, j, p</p>		<p>Elevation (m)</p>	<150
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Ovb
		Drainage	p
		SMR	7
		SNR	C-E
		Plots: DHG035, HRG037	

TEM Map Code	Site Association	CDFmm Site Series
LS	Shore pine—Sphagnum	10
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE		
LS3 (Shrub/Herb)		Mature Undisturbed LS6
Dominant Species Labrador tea salal lodgepole pine western redcedar western hemlock Associate Species sedge step moss hardhack red-stemmed feathermoss western bog-laurel peat-moss Oregon beaked-moss skunk cabbage		Tree Layer shore pine Shrub Layer Labrador tea salal Herb Layer n/a Moss Layer sphagnum moss step moss electrified cat's tail moss

Distribution of Map Units Containing LS Units



TEM Map Code	Site Association	CDFmm Site Series	
RC	Western redcedar—Skunk cabbage	11	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar—Skunk cabbage forests are nutrient-medium to nutrient-rich swamps with poor drainage. Soils often include a major component of organic material intermixed with deep, medium textured soils such as silty loam or silty clay loam. These forests inhabit level sites to depressions. Western redcedar, red alder, bigleaf maple and western hemlock are the dominant trees. The shrub layer has moderately high cover but low diversity, comprising of salal and salmonberry. The herb layer is often relatively diverse with many species, but the major components are lady fern, skunk cabbage, three-leaved foamflower and lady fern. The moss layer often has a high cover of Oregon beaked-moss, palm tree moss, slender beaked-moss and leafy moss.</p> <p>This unit is well represented in some areas of the CDFmm, but is not very common. Its distribution is very restricted in the Bowser area, south to the Cowichan Valley. The highest concentrations of RC units are found in the Powell River area and associated Islands, Sechelt, the Cowichan Valley and Texada and Denman Islands. Several units are mapped in Saanich and on Galiano and Saltspring Islands.</p> <p>These swamp forest are often adjacent to wetlands and are commonly complexed with LS, Ws50, Ws51, and Wf50. RC sites are also found near water courses and on terraces above water courses, where the it is commonly complexed with RF ecosystems.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m)</p>	<150
		Slope (%)	<20
		Aspect (°)	variable
		Surficial material	Ov, M
		Drainage	p
		SMR	7
		SNR	C-E
		Plots: 00-01808, BB_HR_G11, CHG078, HRG158, HRV001, HRV002, JCG019, JKV001, JKV003, JK127, MT014	


TEM Map Code	Site Association		CDFmm Site Series
RC	Western redcedar—Skunk cabbage		11
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE			
RC3 (Shrub/Herb)	RC5 (Young Forest)	RC6 (Mature Forest)	Mature Undisturbed RC6
Dominant Species red alder common horsetail Nootka rose skunk cabbage salmonberry western redcedar Canada thistle common rush meadow buttercup Cooley's hedge-nettle enchanter's-nightshade Associate Species sword fern	Dominant Species western redcedar red alder skunk cabbage salal Associate Species western hemlock leafy moss sword fern Oregon beaked-moss	Dominant Species western redcedar lady fern palm tree moss red alder Associate Species slender beaked-moss Oregon beaked-moss deer fern western hemlock piggy-back plant Sitka spruce salmonberry skunk cabbage large leafy moss leafy moss bigleaf maple three-leaved foamflower flat-moss spiny wood fern stink currant sword fern	Tree Layer western redcedar bigleaf maple Shrub Layer salmonberry red elderberry indian plum salal Herb Layer false lily-of-the-valley skunk cabbage lady fern three-leaved foamflower Moss Layer slender beaked moss palm tree moss

TEM Map Code	Site Association		CDFmm Site Series
RC	Western redcedar—Skunk cabbage		11
RCc	coarse-textured soil	RCp	peaty material
RCf	fine-textured soil	RCs	shallow soil
RCfs	fine-textured soil; shallow soil	RCt	terrace
RCn	fan		
Atypical Characteristics/Additional Comments: RC units are commonly found on peaty material in the CDFmm and on coarse-textured soils with a duric horizon below. Younger forests are typically dominated by red alder, with scattered western redcedar.			

Distribution of Map Units Containing RC Units




Forested Fluctuating Water Table Site Series in the CDFmm

TEM Map Code	Site Association	CDFmm Site Series	
RV	Western redcedar—Vanilla-leaf	12	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar-Vanilla-leaf ecosystems are nutrient-rich and are located on lower gentle slopes and often have fluctuating water tables. Soils are deep and medium textured, but could have a restricting layer or be adjacent to a seepage or riparian site. The soil texture and seasonally high water table may result in susceptibility to disturbance via soil compaction.</p> <p>The tree canopy is dense with gaps, including major components of grand fir, Douglas-fir, bigleaf maple, red alder, and western redcedar. Shrubs such as salal, dull Oregon-grape, salmonberry and oceanspray also occur. Sword fern is frequently abundant. The bryophyte layer features Oregon beaked moss and step moss.</p> <p>RV units are fairly common in the CDFmm, in the Bowser/Parksvill/Qualicum area, Ladysmith and on Gabriola Island.</p> <p>This unit is the driest of the three fluctuating water table units and is often complexed with either of the other two units, RP and CS. It is also complexed with the zonal unit DS and the rich, moist unit RF.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m) 20-200</p> <p>Slope (%) <35</p> <p>Aspect (°) variable</p> <p>Surficial material W^G, M (F)</p> <p>Drainage m</p> <p>SMR 5</p> <p>SNR C</p>	
		<p>Plots: 00-01875, HRG046, HRG150, HRG154, HRG180, HRG181, HRG191, HRV110, HRV111, HRV119, HRV123, HRV124, HRV125, HRV206, HRV216, JCG149, JCG161, JCG164</p>	

TEM Map Code	Site Association		CDFmm Site Series
RV	Western redcedar—Vanilla-leaf		12
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE			
RV3 (Shrub/Herb)	RV5 (Young Forest)	RV6 (Mature Forest)	Mature Undisturbed RV6
Dominant Species sword fern salmonberry red alder Douglas-fir	Dominant Species Douglas-fir salal western redcedar red alder sword fern	Dominant Species sword fern Douglas-fir western redcedar salal red alder Oregon beaked-moss grand fir step moss salmonberry	Tree Layer red alder grand fir Douglas-fir bigleaf maple western flowering dogwood
Associate Species salal black gooseberry western redcedar Oregon beaked-moss western hemlock western white pine bigleaf maple	Associate Species dull Oregon-grape Oregon beaked-moss bigleaf maple trailing blackberry salmonberry Sitka spruce bracken fern oceanspray	Associate Species red huckleberry dull Oregon-grape lanky moss bigleaf maple western hemlock oceanspray	Shrub Layer salal indian-plum oceanspray dull Oregon-grape
			Herb Layer sword fern vanilla-leaf three-leaved foamflower wall-lettuce
			Moss Layer Oregon beaked moss slender beaked moss palm tree moss coastal leafy moss

TEM Map Code	Site Association		CDFmm Site Series
RV	Western redcedar—Vanilla-leaf		12
RVc	coarse-textured soil	RVhs	hummocky; shallow soil
RVcn	coarse-textured soil; fan	RVp	peaty material
RVcs	coarse-textured soil; shallow soil	RVs	shallow soil
RVh	hummocky	RVv	very shallow soil
Atypical Characteristics/Additional Comments: Most RV ecosystems in the CDFmm occur on glaciomarine or glaciofluvial deposits where soils are generally deep and coarse-textured. Several of these sites occur on peaty soils, in the Bowser/Qualicum area and on Texada Island. These can be hummocky, with wet areas where the more moist fluctuation water table ecosystems, RP and CS, occur.			

Distribution of Map Units Containing RV Units



TEM Map Code	Site Association	CDFmm Site Series	
RP	Western redcedar—Indian-plum	13	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Western redcedar-Indian plum forest occurs on fluctuating water tables and is similar to RV ecosystems in terms of soil nutrient status and slope position. In comparison, RC ecosystems are wetter, with less cover of Douglas-fir and more red alder, especially in disturbed sites. The increased presence of red alder may reflect the gap dynamics of these stand types as the high water tables limit rooting depth and soil bearing strength, causing individual trees to fall over and expose mineral soil seedbeds for red alder which requires both light and mineral substrate. Western redcedar, bigleaf maple, grand fir and western hemlock also occur.</p> <p>RP ecosystems have more berry-producing shrubs which indicate richer sites (e.g., trailing blackberry, salmonberry) as well as salal and oceanspray. The herb layer is relatively sparse as understory light is limited by the dense forest canopy; shade-tolerant sword fern is the most common species. The herb layer is poorly developed while moss cover is relatively high, dominated by Oregon beaked-moss, step moss and palm tree moss.</p> <p>The RP unit is often occurs in drainage channels and creeks running through cultivated fields. It was the predominant wet forest associated with cultivated fields and rural creeks on level ground with fluctuating water tables.</p> <p>This unit is common within the CDFmm and has the greatest representation of the three fluctuating water table ecosystems. It is distributed throughout the CDFmm, with the exception of Powell River and associated islands, Texada, Denman, Galiano, and Saltspring Islands. RP sites are most commonly found in the Bowser/Parksville/Qualicum area, Sechelt, Saanich Peninsula and on Thormanby and Gabriola Islands.</p> <p>RP ecosystems are often complexed with the other two fluctuating water tables, RV and CS. It often occurs within large DS units where small streams and creeks are present.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m)</p>	350
		Slope (%)	<35
		Aspect (°)	variable
		Surficial material	M, W ^G
		Drainage	m - i
		SMR	6
		SNR	D





Plots: 3977, 6114, 08078, 98-02172, CHG106, CHG253, DHG014, HRG041, HRG044, HRG152, HRG156, HRG214, HRG220, HRV043, HRV084, HRV096, HRV120, HRV170, HRV175, HRV198, HRV208, JCG217, MT012

TEM Map Code	Site Association				CDFmm Site Series
RP	Western redcedar—Indian-plum				13
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE					
RP3 (Shrub/Herb)	RP4 (Immature Forest)	RP5 (Young Forest)	RP6 (Mature Forest)	Mature Undisturbed RP6	
Dominant Species sword fern red alder salmonberry Douglas-fir	Dominant Species red alder salmonberry sedge bracken fern salal common horsetail	Dominant Species red alder salmonberry sword fern western redcedar Douglas-fir	Dominant Species Douglas-fir red alder sword fern grand fir western redcedar salal salmonberry	Tree Layer red alder grand fir bigleaf maple Douglas-fir	
Associate Species trailing blackberry reed canarygrass bigleaf maple vanilla-leaf	Associate Species western redcedar western white pine	Associate Species Oregon beaked-moss salal grand fir western hemlock bigleaf maple step moss dull Oregon-grape	Associate Species Oregon beaked-moss trailing blackberry red huckleberry western hemlock bigleaf maple step moss palm tree moss oceanspray	Shrub Layer indian-plum trailing blackberry common snowberry salmonberry red elderberry oceanspray	
				Herb Layer sword fern three-leaved foamflower	
				Moss Layer slender beaked moss Oregon beaked moss palm tree moss coastal leafy moss	

TEM Map Code	Site Association		CDFmm Site Series
RP	Western redcedar—Indian-plum		13
RPC	coarse-textured soil	RPn	fan
RPcn	coarse-textured soil; fan	RPp	peaty material
RPCs	coarse-textured soil; shallow soil	RPs	shallow soil
RPct	coarse-textured soil; terrace	RPt	terrace
RPh	hummocky	RPw	warm aspect
RPks	cool aspect; shallow soil		
Atypical Characteristics/Additional Comments: These sites are common on coarse-textured soils over glaciomarine deposits and have been observed on peaty soils in the Bowser area. They are also commonly found in large cultivated fields in the Qualicum/Parksville area and on the Saanich Peninsula, where small streams and drainage channels occur. On these sites, a couch grass, reed canary grass and a variety of sedges and rushes are often growing. Species composition in young stands generally consists of dense red alder with some scattered western redcedar. The understory in these young stands is dominated by salmonberry, sedges and horsetails.			

Distribution of Map Units Containing RP Units




TEM Map Code	Site Association	CDFmm Site Series	
CS	Western redcedar—Slough sedge	14	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Western redcedar-Slough sedge swamp forest occurs on fluctuating water tables on poorly drained level sites and depressions. Western redcedar and sword fern may dominate elevated microsites, while sedges, lady fern, and horsetails occupy hollows with occasional skunk cabbage. Soils are moderately deep to deep (0.5-1+ m) with medium texture, typically gleyed, with fluctuating water tables.</p> <p>Tree species are limited to shade- and moisture-tolerant trees with relatively shallow roots: western redcedar, grand fir on margins, as well as red alder. Shrubs in CS sites are diverse, with salmonberry, salal, trailing blackberry, red-osier dogwood and common snowberry. Herbaceous species are variable, with slough sedge the most common component of the herb layer, and sword fern also common. Moss tends to occupy little of the substrate with Oregon beaked-moss, step moss, slender beaked-moss and palm tree moss present.</p> <p>This ecosystem is fairly common and is distributed throughout the CDFmm, with the exceptions of Powell River area, Denman, Galiano, Gabriola Islands The greatest concentration of this unit is mapped in the Bowser/Qualicum /Parksville area and Lasqueti Island.</p> <p>The CS unit commonly occurs in small pockets complexed with other wet forest ecosystems such as the RP, RF and RC. Wetlands, Wm50 and Wm51 also commonly occur adjacent to CS units.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m) 0-150 m</p> <p>Slope (%) <35</p> <p>Aspect (°) variable</p> <p>Surficial material O, M, L, W^G</p> <p>Drainage m - p</p> <p>SMR 6-7</p> <p>SNR C-E</p>	
		<p>Plots: 00-01806, 00-01809, CA008, CA039, CHG120, CXG007, DHG004, , HRG020, HRG096, HRG133, HRG153, HRG161, HRV003, HRV042, HRV066, HRV121, HRV159, MT036</p>	

TEM Map Code	Site Association				CDFmm Site Series
CS	Western redcedar—Slough sedge				14
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE					
CS3 (Shrub/Herb)	CS4 (Immature Forest)	CS5 (Young Forest)	CS6 (Mature Forest)	Mature Undisturbed CS6	
Dominant Species n/a	Dominant Species salmonberry red alder lady fern slough sedge western redcedar	Dominant Species red alder sword fern slough sedge western redcedar salmonberry Douglas-fir	Dominant Species slough sedge red alder western redcedar salmonberry Oregon beaked-moss sword fern salal	Tree Layer black cottonwood red alder grand fir	
Associate Species salmonberry Pacific water-parsley slough sedge red alder brome little buttercup purple-leaved willowherb wildrye Pacific crab apple reed canarygrass skunk cabbage western hemlock grand fir Cooley's hedge-nettle giant horsetail red huckleberry salal western redcedar	Associate Species common horsetail red huckleberry Sitka spruce sword fern deer fern salal	Associate Species grand fir salal trailing blackberry red-osier dogwood western hemlock step moss slender beaked-moss Oregon beaked-moss lanky moss	Associate Species trailing blackberry western hemlock grand fir slender beaked-moss step moss cascara red-osier dogwood lady fern Sitka spruce lodgepole pine bigleaf maple Douglas-fir creeping buttercup palm tree moss common snowberry	Shrub Layer common snowberry red-osier dogwood indian-plum Pacific crab apple black twinberry trailing blackberry salmonberry red elderberry Pacific ninebark	
				Herb Layer slough sedge sword fern false lily-of-the-valley water-parsley Cooley's hedge-nettle	
				Moss Layer slender beaked moss palm tree moss	

TEM Map Code	Site Association		CDFmm Site Series
CS	Western redcedar—Slough sedge		14
CSc	coarse-textured soil	CSs	shallow soil
CScn	coarse-textured soil; fan	CSt	terrace
CSct	coarse-textured soil; terrace	CSw	warm aspect
CSp	peaty material		
Atypical Characteristics/Additional Comments: These sites are generally associated with coarse-textured soils over glaciomarine deposits. Several of these wet forested ecosystems were found on peaty soils over glaciomarine deposits. Younger stands are often dominated by red alder, with salmonberry, slough-sedge and lady fern in the understory.			

Distribution of Map Units Containing CS Units




Non-Forested and Sparsely Vegetated Ecosystems of the CDFmm

TEM Map Code	Site Association	CDFmm Site Series	
AS	Aspen—Slough sedge	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Aspen – Slough sedge ecosystem is extremely rare and no plot data was collected. It was mapped as structural stage 3, 4 and 5. Trembling aspen, very uncommon in the CDFmm and Saltspring Island, is known to occur in localized treed swamp sites in the study area from the Sensitive Ecosystem Inventory (Ward et al. 1997). AS ecosystems provide a nutrient-rich environment, with contiguous understory cover of reeds, sedges, and other species tolerant of seasonal flooding and high water table. This may be a successional phase that would, if left undisturbed, develop into a conifer-dominated forest type such as site series RP or CS.</p> <p>These ecosystems are small, and difficult to identify on air photos. Sites that are mapped were either sampled or known by the mapper. These mapped locations are on southern Hornby Island, south Nanaimo and south Victoria area.</p> <p>Assumed modifiers: j, m</p>		Elevation (m)	60-200 m
		Slope (%)	0
		Aspect (°)	n/a
		Surficial material	M, L
		Drainage	i - p
		SMR	4-5
		SNR	D
No photo available		Plots: N/A	

TEM Map Code	Site Association	CDFmm Site Series
AS	Aspen—Slough sedge	00
ASs shallow soil Atypical Characteristics/Additional Comments: N/A		

Distribution of Map Units Containing AS Units



TEM Map Code	Site Association	CDFmm & CWHxm Site Series	
FC	Fescue—Camas	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Fescue-Camas ecosystems are common, typically occurring on gentle to moderately sloping, warm aspects with deep but occasionally shallow soils. FC sites were predominantly mapped as grasslands (structural stage 2b), however the ratio of Graminoids to forbs varied considerably depending on site conditions.</p> <p>This non-forested unit is characterized by the presences of graminoids, forbs and bryophytes; shrubs are infrequent. Common snowberry, Nootka rose, and invasives (e.g., Scotch broom) are the most common shrub species. Trees are sparse and shrubby in habit, including Garry oak, arbutus, and Douglas-fir which may also be present as scattered veterans on site. Herbaceous species are somewhat variable, and depending on distance from disturbance, may contain a high proportion of invasive perennial grasses such as early hairgrass, orchard grass, Kentucky bluegrass, and many others.</p> <p>Native herbaceous plants include camas species, western and Roemer's fescue, Pacific sanicle, Alaska oniongrass, <i>Brodiaea</i> species., <i>Lomatium</i> species, and others which varied by season and presence among sites. Bryophytes and lichens were not abundant; rock moss and Wallace's selaginella occasionally colonized bedrock outcrops.</p> <p>FC is rare in the CDF and is mapped on Vancouver Island south of Nanoose. Most units are mapped in the Cowichan valley and on Saltspring Island. Several small areas are mapped in Saanich and Nanaimo.</p> <p>Assumed modifiers: j, m, s</p>		<p>Elevation (m)</p> <p>0-550</p>	
		<p>Slope (%)</p> <p>15-55</p>	
		<p>Aspect (°)</p> <p>135-275</p>	
		<p>Surficial material</p> <p>C, M, W^G</p>	
		<p>Drainage</p> <p>r - x</p>	
		<p>SMR</p> <p>2</p>	
		<p>SNR</p> <p>C-E</p>	
		Plots: TT002, TT003, TT004, JK217, CHV011	

TEM Map Code	Site Association		CDFmm & CWHxm Site Series
FC	Fescue—Camas		00
FCcn	coarse-textured soil; fan	FCkv	cool aspect; very shallow soil
FCd	deep soil	FCv	very shallow soil
FCh	hummocky	FCvw	very shallow soil; warm aspect
FChv	hummocky; very shallow soil	FCw	warm aspect
FCK	cool aspect		
Atypical Characteristics/Additional Comments:			
N/A			

Distribution of Map Units Containing FC Units



TEM Map Code	Site Association	CDFmm Site Series	
GO	Garry oak—Oceanspray	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Garry oak-Oceanspray ecosystem occurs mostly on the slopes of Mount Maxwell found on Saltspring Island on steep, south-facing, very rapidly drained thin colluvial material over bedrock. This ecosystem type is defined using a combination of information from Erickson and Meidinger (2007) and Madrone (2003) to determine the site attributes typical for this community. Sites dominated by shrubs and Garry oak with a relatively contiguous herbaceous layer were mapped as GO (structural stage 3b), although species composition varied.</p> <p>This ecosystem unit was only mapped on Saltspring Island.</p> <p>Assumed modifiers: j, m, r</p>		Elevation (m)	50-550
		Slope (%)	40-100
		Aspect (°)	135-270
		Surficial material	C, R
		Drainage	x
		SMR	1-2
		SNR	B
No photo available		Plots: N/A	

TEM Map Code	Site Association	CDFmm Site Series
GO	Garry oak—Oceanspray	00
GOsw shallow soil; warm aspect GOvw very shallow soil; warm aspect GOvz very shallow soil; very steep warm aspect Atypical Characteristics/Additional Comments: N/A		

Distribution of Map Units Containing GO Units




TEM Map Code	Site Association	CDFmm Site Series	
LM	Dunegrass—Beach pea	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>These ecosystems were rare within the CDFmm and were mapped as structural stages 1 to 3. These sites are found along the coast line in several areas of the CDF. Vegetation is generally sparse, where invasive species have not impacted the site. Dunegrass (<i>Elymus mollis</i>) is common on these sites, with beach pea (<i>Lathyris japonica</i>) and other herbaceous species scattered.</p> <p>These sites are rare and are limited to Denman, Savory, Harwood, Hermando, Valdez and Galiano Islands and to a few coastal areas in Nanaimo and on the Saanich Peninsula.</p> <p>Assumed modifiers: j, m, v</p>		Elevation (m)	5-30
		Slope (%)	0-10
		Aspect (°)	variable
		Surficial material	W, WG, C
		Drainage	r
		SMR	1-2
		SNR	B
No photo available		Plots: N/A	

TEM Map Code	Site Association	CDFmm Site Series
LM	Dunegrass—Beach pea	00
LMc coarse-textured soil LMcw coarse textured soil; warm aspect LMw warm aspect Atypical Characteristics/Additional Comments: These sites are vulnerable to invasive species, where Scotch broom and non-native grasses easily become dominant.		

A map of Indonesia and surrounding regions, including parts of Malaysia, the Philippines, and Australia. The Indonesian archipelago is highlighted in light green, while surrounding landmasses are in light yellow and water bodies are in light blue. Ten specific locations are marked with red circles: 1. Sumatra (northern tip), 2. Sumatra (central), 3. Sumatra (southern tip), 4. Java (western coast), 5. Java (central), 6. Java (eastern coast), 7. Sulawesi (northern coast), 8. Sulawesi (central), 9. Sulawesi (southern coast), and 10. Sulawesi (southeastern coast). A north arrow is located in the top left corner.



TEM Map Code	Site Association	CDFmm Site Series	
OM	Garry oak—moss	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Garry oak-Moss ecosystem on Saltspring Island reflects the associations classified by Erickson and Meidinger (2007). The OM ecosystem is dominated by Garry oak with the understory containing a constant dominant or major associate species component of <i>Dicranum</i> moss over thin substrate, or directly over bedrock, and likely to the oak–rock moss association as well. Crustose and foliose lichens are also typically present.</p> <p>OM ecosystems are nutrient-poor and drain rapidly, and litter accumulations likely yield successional transition to more complex and richer ecosystem types. This ecosystem unit was infrequently mapped on Saltspring Island, occurring on warm aspects and was mapped as structural stages, 3b, 4 and 6.</p> <p>From plot data dominant species included Garry oak, heron's-bill moss, Douglas-fir, rock-moss and arbutus. Associate species included Scotch broom, hairy honeysuckle, licorice fern, haircap moss, slender wheatgrass, common foxglove, oceanspray, Pacific sanicle, Idaho fescue and yarrow.</p> <p>These ecosystems are only mapped on Saltspring and Galiano Islands.</p> <p>Assumed modifiers: j, s</p>		<p>Elevation (m)</p>	10-350
		Slope (%)	30-65
		Aspect (°)	140-225
		Surficial material	R
		Drainage	x
		SMR	0-1
		SNR	A-B
		Plots: CHG287, CHG257, JKV121, JCV235, JCG225	

TEM Map Code	Site Association	CDFmm Site Series
OM	Garry oak—moss	00
OMk cool aspect OMv very shallow soil OMvw very shallow soil; warm aspect Atypical Characteristics/Additional Comments: N/A		

Distribution of Map Units Containing OM Units



TEM Map Code	Site Association	CDFmm Site Series	
OR	Oceanspray—Rose	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The shrub-dominated Oceanspray – Rose ecosystem was only mapped infrequently. It is likely more frequent, but occurs within mosaics of other ecosystems and is difficult to distinguish a scale of 1:20000. OR ecosystems could be a seral or disturbance-driven phase of another ecosystem type such as DS.</p> <p>Assumed modifiers: m, s</p>		Elevation (m)	100
		Slope (%)	35
		Aspect (°)	155
		Surficial material	M
		Drainage	w
		SMR	4
		SNR	c
No photo available		Plots: DYVH03, DYVH04, DYVH05	

TEM Map Code	Site Association	CDFmm Site Series
OR	Oceanspray—Rose	00
ORw warm aspect Atypical Characteristics/Additional Comments: N/A		

Distribution of Map Units Containing OR Units




TEM Map Code	Site Association	CDFmm & CWHxm Site Series	
QB	Garry oak—Brome/mixed grasses	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Garry oak-Brome/mixed grasses unit corresponds with the associations proposed by Erickson and Meidinger (2007) where the overstory is sparse and consisting solely of Garry oak, and the understory includes graminoids and forbs. The composition of the herbaceous layer varies with surficial material and thickness, slope, and drainage: relatively mesotrophic sites contain mixtures of graminoids and forbs, while richer sites have fewer graminoids and more species found in drier Garry oak meadows including members of the lily family and yellow montane violet.</p> <p>The Garry oak ecosystem is a distinctive feature of the CDFmm landscape. The Parkland Garry Oak unit was mapped as QB structural stage 5 with no shallow soil modifier. The Scrub Oak unit was mapped as QB with a structural stage 3, 4 or sometimes 5 and always with a shallow or very shallow soil modifier. The Parkland Garry Oak unit most commonly occurred as a minor component in complex with cultivated field and rural landscapes. Deep glaciomarine material produces rich soil and support large, robust Garry oak trees. The Rocky Outcrop Garry Oak unit is more abundant and can be found occurring in areas such as Mt. Tzouhalem, Maple Mountain, Cobble Hill, Mt. Douglas, Mt. Tolmie, Mt Work, Gowlland Tod Provincial Park and around Prospect Lake and Elk Lake. From plot data dominant species included Scotch broom, Garry oak, barren fescue, hedgehog dogtail, Kentucky bluegrass, orchard-grass, blue wildrye and hairy cat's-ear with oceanspary as an associate species. This ecosystem is rare within the CDFmm and has been mapped mostly in Saanich and Victoria, with several sites on Hornby Island, Nanoose Bay, coastal areas of Nanaimo, the Cowichan Valley, and on Saltspring Island.</p> <p>Assumed modifiers: j, m, r</p>		<p>Elevation (m)</p>	20-550
		<p>Slope (%)</p>	20-50
		<p>Aspect (°)</p>	15-260
		<p>Surficial material</p>	R, C (M)
		<p>Drainage</p>	r - x
		<p>SMR</p>	1-2
		<p>SNR</p>	B




Plots: TT002, TT003, TT004, MT039, CHG024, CHV030

TEM Map Code	Site Association		CDFmm & CWHxm Site Series
QB	Garry oak—Brome/mixed grasses		00
QBh	hummocky	QBkv	cool aspect; very shallow soil
QBhs	hummocky; shallow soil	QBw	very shallow soil; warm aspect
QBhv	hummocky; very shallow soil	QBw	warm aspect
QBk	cool aspect	QBz	very steep warm aspect
QBks	cool aspect; shallow soil	QBsw	shallow soil; warm aspect
		QBsz	shallow soil; very steep warm aspect
		QBv	very shallow soil
Atypical Characteristics/Additional Comments:			
N/A			

Distribution of Map Units Containing QB Units



TEM Map Code	Site Association	CDFmm Site Series	
RA	Nootka rose—Pacific crab apple	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Nootka rose – Pacific crab apple ecosystem occurs on the upper portions of estuaries, often between the forest and herbaceous areas. It generally occurs in narrow strips along river estuaries, such as Witty's Lagoon and the Englishman, Cowichan, and Nanaimo River estuaries. These shrubby RA ecosystems are generally dominated by dense (15-80%) cover of Nootka rose with scattered Pacific crab apple and willows. As these sites are often highly disturbed, they can contain numerous non-native and invasive plants as well such as Scotch broom and English ivy. The herb layer cover (10 – 40%) is dominated by grasses and can include reed canary grass, <i>Elymus</i> species, <i>Bromus</i> species, creeping bentgrass, and orchard grass. Mosses were found on one site only. From plot data dominant species included Nootka rose, Himalayan blackberry, lodgepole pine, slender beaked-moss, orchard-grass, Douglas-fir, common hawthorn, Pacific crab apple, creeping buttercup, Oregon beaked-moss and Scotch broom. Associate species included English holly, tufted hairgrass and common snowberry. This ecosystem was identified during the CDFmm mapping project, and is rare within the CDFmm. Known map sites are located in Parksville on the Englishman River Estuary, in Nanaimo on the Nanaimo River Estuary, in the Cowichan Valley on the Chemainus River Estuary and the Cowichan River Estuary, in Saanich on Goldstream Estuary and in areas on Valdez, Galiano, and Thetis Islands.</p> <p>Assumed modifiers: N/A</p>		<p>Elevation (m)</p>	<20
		Slope (%)	0-5
		Aspect (°)	999
		Surficial material	WG, FG
		Drainage	m
		SMR	6
		SNR	B
		Plots: CHG279a, CHG279b	


TEM Map Code	Site Association	CDFmm & CWHxm Site Series	
SC	Cladina—Wallace's selaginella	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Cladina-Wallace's selaginella ecosystem is a non-forested ecosystem occurring on shallow to very shallow soils typical of rock outcrops. Typically it occurs on warm, south facing aspects but can also occur on cool aspects. This ecosystem can be subdivided into several associations, as some sites are dominated by broom moss, others by rock moss and yet others by Cladina and Wallace's selaginella. Non-vascular flora dominates the unit with species such as <i>Cladonia</i> species, hoary and common rock moss, sidewalk moss, and Wallace's selaginella. Field inspection revealed this community varied to include components of diverse vascular species such as stonecrops, and frequently contained low to moderate cover of introduced grass species on richer sites.</p> <p>The SC unit was often mapped in association with DA and QB units and was typically mapped as structural stage 1b, 2a and 2b.</p> <p>From plot data dominant species included Wallace's selaginella and Scotch broom, while associate species included arbutus, Douglas-fir, blue wildrye, long rock-moss, grey rock-moss, awned haircap moss, juniper haircap moss, hedgehog dogtail, hairy Manzanita, early hairgrass, sweet vernalgrass, oceanspray, kinnikinnick, Columbia brome, common camas, curly heron's-bill moss, gorse, rockgossamer lichens, western fescue, freckle pelt and Hooker's onion.</p> <p>This unit is quite common in the CDFmm and its distribution ranges from Nanoose south to Victoria on Vancouver Island and in Powell River, Sechelt, Texada, Lasqueti, Thormanby, Valdes, Thetis and Saltspring Islands. The highest concentration of this ecosystem is in the Saanich/Victoria area.</p> <p>Assumed modifiers: j, m, r, v</p>		<p>Elevation (m)</p> <p>0-550</p>	
		<p>Slope (%)</p> <p>0-100</p>	
		<p>Aspect (°)</p> <p>135-285 (varies)</p>	
		<p>Surficial material</p> <p>R (Mx, W^Gx)</p>	
		<p>Drainage</p> <p>x</p>	
		<p>SMR</p> <p>0-1</p>	
		<p>SNR</p> <p>A-B</p>	




Plots: TTV21, CHG015, CHV026, CHV169, CHV188, CHV211, CHV239, CHG269, HRG227, HRG231, JKG207

TEM Map Code	Site Association		CDFmm & CWHxm Site Series
SC	Cladina—Wallace's selaginella		00
SCh	hummocky	SCs	shallow soil
SChw	Hummocky; warm aspect	SCsw	shallow soil; warm aspect
SCK	cool aspect	SCw	warm aspect
SCq	very steep cool aspect	SCz	very steep warm aspect
Atypical Characteristics/Additional Comments: Herbaceous species more typical of richer sites were noted in depressional microsites and crevices, such as <i>Brodiaea</i> and common camas; monkeyflower species were infrequent in seepage sites.			

Distribution of Map Units Containing SC Units



Wetland Ecosystems in the CDFmm

TEM Map Code	Site Association	CDFmm Site Series	
Ed01	Tufted hairgrass—Meadow barley	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Tufted hairgrass – Meadow barley ecosystem is found at the Englishman River estuary near Parksville and Witty's Lagoon in Metchosin. Ed01 sites occur in the upper intertidal zone on fan estuaries and are briefly flooded each day with brackish water. These sites are characterized by low species diversity dominated by tufted hairgrass and introduced grasses such as meadow barley and bentgrasses. This estuarine meadow ecosystem occurs on gleyed sandy soils where ground seepage is present.</p> <p>From plot data dominant species included Puget Sound gumweed, reed canarygrass, wildrye and fescue. Associate species included yarrow, green sorrel and Scotch broom.</p> <p>This ecosystem is rare in the CDFmm and has been mapped on the Englishman River, Qualicum River, Nanaimo River, Chemanius River, and Cowichan River estuaries, on Gabriola Island, Witty's Lagoon and Island View Beach.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		<p>Elevation (m)</p>	0
		<p>Slope (%)</p>	0
		<p>Aspect (°)</p>	999
		<p>Surficial material</p>	Wp
		<p>Drainage</p>	m - p
		<p>SMR</p>	6
		<p>SNR</p>	D
		Plots: HRV162	

Distribution of Map Units Containing Ed01 Units



TEM Map Code	Site Association	CDFmm Site Series	
Ed03	Arctic rush-Alaska plantain	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Arctic rush – Alaska plantain ecosystems are found where there is a well-developed freshwater lens that reduces the salinity of tidal waters such as estuaries near larger river systems or localized depressions within estuarine meadows. The Arctic rush-Alaska plantain unit typically occurs above the Ed01 or Em05 estuarine sites in the high intertidal zone.</p> <p>These sites are characterized by moderate species diversity dominated by Arctic rush, coast silverweed, Douglas' aster, and Alaska plantain. Silty-textured Gleysols with a humic enriched surface horizon are most common, with minerally enriched Fibrisols also present.</p> <p>This ecosystem was mapped in the Saanich area at Island View Beach.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m)	0
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Wp
		Drainage	i - p
		SMR	5-6
		SNR	C-D
No photo available		Plots: N/A	

Distribution of Map Units Containing Ed03 Units



TEM Map Code	Site Association	CDFmm Site Series	
Em01	Widgeon-grass	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Widgeon-grass ecosystems are rare in the lower tidal zone, occurring in brackish, mud-bottomed pools, lagoons, backwater sloughs, drainage channels, and mudflats that dissect lower portions of estuarine marshes. Tidal inundation is usually prolonged, resulting in silty Rego-Gleysols.</p> <p>This unit is rare in the CDFmm and has been mapped in Nanoose Bay, Nanaimo River estuary and in the northeast of Nanaimo near Piper's Lagoon.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m)	0
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Wp
		Drainage	p
		SMR	7
		SNR	B
No photo available		Plot: N/A	


Distribution of Map Units Containing Em01 Units



TEM Map Code	Site Association	CDFmm Site Series	
Em02	Glasswort—Sea-milkwort	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Glasswort – Sea-milkwort ecosystems are estuarine marshes that are rare in the study area. Em02 sites are confined to flat, imperfectly drained sites in riparian outlets such as McKenzie Bight and are protected from the open ocean in the intertidal zone. The dominant species are the aquatic glasswort and sea-milkwort.</p> <p>This ecosystem is located at Englishman River Estuary, Ladysmith coastal areas and Chemainus River Estuary.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m)	0
		Slope (%)	0
		Aspect (°)	999
		Surficial material	F ^A f
		Drainage	i
		SMR	7
		SNR	B
No photo available		Plot: CHV233	


Distribution of Map Units Containing Em02 Units



TEM Map Code	Site Association	CDFmm Site Series	
Em03	Seashore saltgrass	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Seashore saltgrass ecosystem was found at Witty's Lagoon in the southernmost portion of the study area near Metchosin. Seashore saltgrass communities are often found with Em02 sites. Em03 ecosystems occur on fine-textured, poorly drained tidal sediments in brackish estuaries. The dominant species are seashore saltgrass, glasswort and sea milkwort, with few other species.</p> <p>From plot data dominant and associate species included seashore saltgrass and American glasswort, respectively.</p> <p>This ecosystem is generally difficult to identify through air photo interpretation and these sites are often too small to map. They are distributed along the Vancouver Island and Gulf Island coastlines, but few have been mapped.</p> <p>The coastal areas that have been mapped are: Englishman River Estuary, Witty's Lagoon and the Cowichan River Estuary.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		<p>Elevation (m)</p> <p>0</p> <p>Slope (%)</p> <p>0</p> <p>Aspect (°)</p> <p>999</p> <p>Surficial material</p> <p>Ov, Wp</p> <p>Drainage</p> <p>p</p> <p>SMR</p> <p>7</p> <p>SNR</p> <p>B-C</p>	
		Plots: CHV275, CHV280	

Distribution of Map Units Containing Em03 Units



TEM Map Code	Site Association	CDFmm Site Series	
Em05	Lyngbye's sedge	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Lyngbye's sedge site association is commonly found along the coastal shores of BC. In the CDFmm subzone, this estuarine marsh was found at the Cowichan River estuary. Lyngbye's sedge communities are often the lowest vegetated zone grading into Ed02 communities upslope.</p> <p>The dominant species are Lyngbye's sedge, coast silverweed, tufted hairgrass, and sea-milkwort. Em05 ecosystems occur on fine-sandy layered soils where there are strong fluctuations of brackish water.</p> <p>This unit was mapped in the Nanaimo River and Cowichan River Estuaries.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m) 0	
		Slope (%) 0	
		Aspect (°) 999	
		Surficial material W ^G p	
		Drainage p	
		SMR 6	
		SNR B	
		Plots:HRV250	

Distribution of Map Units Containing Em05 Units



TEM Map Code	Site Association	CDFmm & CWHxm Site Series	
Wb50	Labrador tea—Bog-laurel—Peat-moss bog	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Labrador tea – Bog-laurel – Peat-moss ecosystem is uncommon in the CDFmm and was found only on Lasqueti Island and in the Parksville area. Wb50 sites occur in raised peatlands and closed basins with a stagnant watertable. These sites are characterized by moderate species diversity dominated by low ericaceous species such as Labrador tea, bog-laurel, and sweet gale. Peat-moss species dominate the ground layer with scattered acid-loving herbs. This bog ecosystem occurs on organic veneers of poorly developed peat.</p> <p>From plot data dominant species included peat-moss and Labrador tea.</p> <p>This ecosystem was mapped on Lasqueti Island and in the Cowichan Valley.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m) 0 Slope (%) 0 Aspect (°) 999 Surficial material Ov Drainage p-v SMR 7 SNR D	
No photo available		Plots: DHG019	

Distribution of Map Units Containing Wb50 Units



TEM Map Code	Site Association	CDFmm Site Series	
Wf51	Sitka sedge—Peat-moss fen	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Sitka sedge—Peat-moss sites are uncommon in the CDFmm, but do occur on Lasqueti and Texada Islands. These sites are characterized by low species diversity dominated by Sitka sedge and <i>Sphagnum</i> mosses. Water moss and brown moss species may also be present. This fen ecosystem occurs on organic veneers to deep blankets of poorly developed, sloping peatlands with slow surface water flow.</p> <p>This ecosystem is relatively rare in the CDFmm and the highest distribution is found in the Islands adjacent to Powell River including Texada. Several units are located on Galiano Island and a few are scattered throughout Vancouver Island.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: p</p>		Elevation (m) Slope (%) Aspect (°) Surficial material Drainage SMR SNR	 0 0 999 Ob, Ov p-v 7 C-D
No photo available		Plots: DHG032, JCG018	

Distribution of Map Units Containing Wf51 Units



TEM Map Code	Site Association	CDFmm Site Series	
Wf52	Sweet gale—Sitka sedge fen	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Sweet gale—Sitka sedge sites are uncommon in the CDFmm, but occur in a variety of landscape positions that are shallowly flooded during part of the year. Wf52 sites are characterized by low species diversity, dominated by sweet gale, Sitka sedge, and <i>Sphagnum</i> mosses. This fen ecosystem has a closed and dense thicket of sweet gale and hardhack. Sitka sedge dominates the herb layer. Terric Humisols and Mesisols most common.</p> <p>This unit is mapped only on Texada Island and in the northern portion of the Cowichan Valley.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: p</p>		Elevation (m)	0
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Ob
		Drainage	p-v
		SMR	6-7
		SNR	C
No photo available		Plots: A213, A223, CXV038, JCV021, V986	


Distribution of Map Units Containing Wf52 Units



TEM Map Code	Site Association	CDFmm & CWHxm Site Series	
Wf53	Slender sedge—White-beak-rush fen	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>These fens are richer than Wf51 and Wf52 fens. These sites require permanently saturated soils and are tolerant of prolonged shallow inundation. Lake margins are the most common location but some isolated basins may also have suitable conditions. These sites are characterized by low species diversity dominated by slender sedge, white-beak rush, and scattered sweet gale low shrubs. A variety of water mosses and hook mosses may also be present. Terric Mesisols and Humisols are common.</p> <p>These ecosystems are mapped only on Saltspring Island and Saanich.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: d, s</p>		Elevation (m)	0
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Ob
		Drainage	p-v
		SMR	7
		SNR	D
No photo available		Plots: C180, C181, C186, C184	

Distribution of Map Units Containing Wf53 Units



TEM Map Code	Site Association	CDFmm Site Series	
Wm05	Cattail marsh	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The cattail marsh ecosystem is widespread and locally common throughout the study area in protected embayments at open water areas and permanently saturated wetlands. Cattail dominates the vegetation layers with few other species. Cattail marshes are often a result of nutrient loading adjacent to human development and agricultural lands. This marsh ecosystem occurs on well decomposed organic veneers or saturated mineral soils (Gleysols). From plot data dominant species included common cattail and common rush, while associate species included slough sedge.</p> <p>These marshes are distributed throughout the CDFmm mainly on Vancouver Island south of Nanoose, Sechelt, and Texada, Lasqueti, and Thormanby Islands.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: p</p>		Elevation (m)	0
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Ov, Ob
		Drainage	p-v
		SMR	7-8
		SNR	D
		Plots: DHV030, DHG038, DHV052, DHV054, DHV064, HRV217, CHV202, HRV022	

Distribution of Map Units Containing Wm05 Units




TEM Map Code	Site Association	CDFmm Site Series	
Wm06	Great bulrush marsh	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Great bulrush marsh ecosystem overlaps with cattail marshes (Wm05) along lake margins and wave-exposed lake embayments with significant water movement. Bulrush species dominate the vegetation layers with few other species. Soils are mostly Gleysols and Humic Gleysols.</p> <p>This unit was mapped only in the Cowichan Valley on Quamichan Lake.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m)	0
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Ob
		Drainage	v
		SMR	7-8
		SNR	D
No photo available		Plots: N/A	

Distribution of Map Units Containing Wm06 Units



TEM Map Code	Site Association	CDFmm Site Series	
Wm50	Sitka sedge—Hemlock-parsley marsh	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Sitka sedge – Hemlock-parsley marsh ecosystem is very uncommon in the study area. Sitka sedge is always present with a number of other forb and grass species, depending on the substrate and amount of flowing water. Wm50 sites develop on organic veneers and marine deposits along streams and ponds near coastal waters. From plot data dominant species included common rush, small-flowered forget-me-not and slough sedge, while associate species included common green peat-moss and hardhack.</p> <p>This ecosystem was found on Lasqueti Island, Parksville/Qualicum area, Sechelt, Saanich and on Somenos Lake in the Cowichan Valley.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m)	<60
		Slope (%)	0
		Aspect (°)	999
		Surficial material	W ^G b
		Drainage	p
		SMR	8
		SNR	C
No photo available		Plots: DHG035	


Distribution of Map Units Containing Wm50 Units



TEM Map Code	Site Association	CDFmm Site Series	
Wm51	Three-way sedge marsh	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Three-way sedge marsh ecosystem occurs along protected margins of shallow lakes or sluggish streams on mucky substrates. Wm51 sites are transitional between fen ecosystem interfaces. They occur in permanently flooded conditions on degrading peat or soft muck.</p> <p>Several Wm51 ecosystems are mapped in the Cowichan Valley and Saanich.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m)	<60
		Slope (%)	0
		Aspect (°)	999
		Surficial material	Ob
		Drainage	p
		SMR	8
		SNR	C-D
No photo available		Plots: N/A	


Distribution of Map Units Containing Wm51 Units



TEM Map Code	Site Association	CDFmm & CWHxm Site Series	
Ws50	Pink spirea—Sitka sedge swamp	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Pink spirea-Sitka sedge swamps are a common CDFmm wetland type. The shrub layer is dominated by pink spirea with minor components of willows and red-osier dogwood. Sitka sedge may not always be present, depending on the density of spirea and amount of open water. Ws50 sites typically occur on organic veneers of poorly decomposed plant residue formed over gleyed mineral soil.</p> <p>From plot data dominant species included hardhack and slough sedge, while associate species included bluejoint reedgrass and red alder.</p> <p>This ecosystem is the most common wetland in the CDFmm. It is distributed throughout the entire project area.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: p, s</p>		<p>Elevation (m)</p>	30-500
		<p>Slope (%)</p>	0
		<p>Aspect (°)</p>	999
		<p>Surficial material</p>	Ov, Ob, W ^G u, Lp
		<p>Drainage</p>	p
		<p>SMR</p>	7-8
		<p>SNR</p>	B-E
		<p>Plots: CHV041, CHV046, CHV050, CHV066, CHV073, CHV082, CHV128, CHV203, CHV216, CHV266, HRV008, HRV104, HRV105, HRV211, HRG022, HRG033, HRG080, DHV021, DHV053, DHV063, JCV024</p>	


Distribution of Map Units Containing Ws50 Units



TEM Map Code	Site Association	CDFmm & CWHxm Site Series	
Ws51	Sitka willow—Pacific willow—Skunk cabbage swamp	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Sitka willow-Pacific willow-Skunk cabbage swamp ecosystem occurs throughout the study area. Sitka and Pacific willow are common tall shrubs with a diverse understory of herbs and mosses. Skunk cabbage and lady fern are common herbs that thrive in the rich organic soils. These sites may develop near floodplains or marshes, and are often flooded for many months of the year.</p> <p>From plot data associate species included slough sedge, willow, red alder, salmonberry, skunk cabbage, highbush-cranberry, Pacific crab apple and salal.</p> <p>This wetland type is common throughout the CDFmm and is distributed throughout most of the Vancouver Island portion. It is rare on the Gulf Islands, as well as in Powell River and Sechelt.</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: p</p>		<p>Elevation (m)</p>	<100
		<p>Slope (%)</p>	0
		<p>Aspect (°)</p>	999
		<p>Surficial material</p>	Ov, Op, W ^G p
		<p>Drainage</p>	p-v
		<p>SMR</p>	7
		<p>SNR</p>	B-E
		<p>Plots: CHV112, CHV116, CHV247, HRV002, HRV006, HRV045, HRV099, HRG179, CXG039</p>	

Distribution of Map Units Containing Ws51 Units




TEM Map Code	Site Association	CDFmm Site Series	
Ws52	Red alder—Skunk cabbage swamp	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Red-alder – Skunk cabbage ecosystem was found in the northern portion of the study area near Powell River and on Texada, Denman, and Galiano Islands at low elevations. Ws52 swamps are uncommon, but when present occur adjacent to small creeks and floodplains.</p> <p>Red alder and salmonberry are dominant shrub species with occasional tall willows. A diverse understory of herbs, such as skunk cabbage, lady fern and Sitka sedge may be present with variable amounts of moss. These sites have high watertables and lack elevated microsites that allow for conifer development (MacKenzie and Moran, 2004).</p> <p>Assumed modifiers: N/A</p> <p>Modifiers used: N/A</p>		Elevation (m) Slope (%) Aspect (°) Surficial material Drainage SMR SNR	 <100 0 999 Op, Fv p 6-7 C
		Plots: JCV035, JCV227, JCG097, HRV024	

Distribution of Map Units Containing Ws52 Units




Additional Ecosystems in the CWHxm subzone

TEM Map Code	Site Unit Name	CWHxm Site Series	
HK	Western hemlock–Douglas-fir–Oregon beaked moss	01	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Western hemlock – Douglas-fir – Oregon beaked moss unit occurs typically on gentle slopes, with medium textured soils. The zonal CWH forest is distinguished from the zonal CDF forest by the presence of western hemlock. Western hemlock occurs as the dominant trees species in association with Douglas-fir and western redcedar. Western hemlock occurs in all layers of the tree canopy including regeneration layers (B1 and B2). The understory is well developed with a continuous feathermoss layer carpeting the forest floor. Shrubs such as dull Oregon grape and red huckleberry are present and abundant while the abundance of salal is lower than zonal sites in the CDF.</p> <p>Assumed modifiers: d, j, m</p>		Elevation (m):	150-500
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	M, F ^G
		Drainage:	m-w
		SMR:	3-4
		SNR:	A-C
		Plots: 4827, 4831, 4832, CHG071, CHV142, ERSK_GRV0, GBG027, GBG028, GBG029, GBV030, HRG130, HRV131, HRV174, QAG003, WMG004, WME009, WMV038, WMV044, MTV060, MTV061	


TEM Map Code	Site Unit Name		CWHxm Site Series
HK	Western hemlock–Douglas-fir–Oregon beaked moss		01
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE			
HK4 (Immature Forest)	HK5 (Young Forest)	HK6 (Mature Forest)	Mature Undisturbed HK6
Dominant Species Douglas-fir salal Oregon beaked-moss western hemlock Associate Species red alder oceanspray lodgepole pine western redcedar dull Oregon-grape	Dominant Species Douglas-fir salal dull Oregon-grape western hemlock Associate Species step moss oceanspray lodgepole pine red huckleberry western redcedar electrified cat's-tail moss Oregon beaked-moss red alder Scotch broom stinging nettle arbutus	Dominant Species salal Douglas-fir Oregon beaked-moss step moss Associate Species western redcedar western hemlock dull Oregon-grape sword fern oceanspray red alder	Tree Layer Douglas-fir western hemlock western redcedar Shrub Layer salal dull Oregon-grape red huckleberry baldhip rose oceanspray Herb Layer vanilla-leaf sword fern twinline bracken fern Moss Layer step moss Oregon beaked moss lanky moss

TEM Map Code		Site Unit Name		CWHxm Site Series	
HK		Western hemlock–Douglas-fir–Oregon beaked moss		01	
HKh	hummocky	HKqv	very steep cool aspect; very shallow soil	HKvx	very shallow soil; drier than typical
HKhs	hummocky; shallow soil	HKrs	ridge; shallow soil	HKvz	very shallow soil; very steep warm aspect
HKhv	hummocky; very shallow soil	HKs	shallow soil	HKw	warm aspect
HKk	cool aspect	HKsw	shallow soil; warm aspect	HKxz	drier than typical; very steep warm aspect
HKks	cool aspect; shallow soil	HKv	very shallow soil	HKz	very steep warm aspect
HKkv	cool aspect; very shallow soil	HKvw	very shallow soil; warm aspect		
Atypical Characteristics/Additional Comments: N/A					

TEM Map Code	Site Association	CWHxm Site Series	
DC	Douglas-fir—Lodgepole pine—Cladina	02	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Dougals-fir-Lodgepole pine-Cladina unit was infrequently mapped, generally occurring on water-shedding ridge crests and convex upper slopes with very thin till-derived Brunisols or bedrock outcrops. Douglas-fir and shore pine were dominant in the canopy, with more pine occurring where soil conditions were driest. Canopy cover was sparse (20-40%), with much of the forest floor exposed. Dominant shrubs included salal and dull Oregon-grape. Arbutus and regenerating canopy species were frequent associates that occupied the shrub layer. Herb cover was sparse to nil during winter when sites were surveyed. Rocks and substrate were often covered by <i>Cladina</i> lichen species and dominant mosses including step moss, juniper haircap moss, curly heron's bill-moss, broom moss and red-stemmed feather moss.</p> <p>Assumed modifiers: j, m, r, s</p>		Elevation (m)	250-650
		Slope (%)	5-65
		Aspect (°)	999
		Surficial material	Mv
		Drainage	r
		SMR	0
		SNR	A (-B)
		Plots: WMG001, WM006, WM007, ERSK_GR02, ERSK_GR03, WM036, JK304, JKG309, JKG314, ERSK_GR05	


TEM Map Code	Site Association	CWHxm Site Series
DC	Douglas-fir—Lodgepole pine—Cladina	02
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE		
DC4 (Immature Forest)	DC5 (Young Forest)	Mature Undisturbed DC6
Dominant Species curly heron's-bill moss juniper haircap moss shore pine brome fire-moss salal Associate Species Douglas-fir hoary rock-moss reindeer lichens step moss	Dominant Species salal Douglas-fir Associate Species dull Oregon-grape curly heron's-bill moss juniper haircap moss broom-moss red-stemmed feathermoss arbutus shore pine sweet vernalgrass yellow curl-moss hoary rock-moss prince's pine rock-moss ragged-moss	Tree Layer shore/lodgepole pine Douglas-fir arbutus Shrub Layer salal oceanspray dull Oregon-grape red huckleberry baldhip rose Herb Layer hairy cat's ear Moss Layer Oregon beaked moss step moss juniper haircap moss electrified cat's-tail moss red-stemmed feathermoss lichen

TEM Map Code	Site Association		CWHxm Site Series
DC	Douglas-fir—Lodgepole pine—Cladina		02
DChv	hummocky; very shallow soil	DCv	very shallow soil
DCkv	cool aspect; very shallow soil	DCvw	very shallow soil; warm aspect
DCs	shallow soil	DCvz	very shallow soil; very steep warm aspect
DCsw	shallow soil; warm aspect	DCw	warm aspect
Atypical Characteristics/Additional Comments: N/A			


TEM Map Code	Site Association	CWHxm Site Series	
DS	Douglas-fir—Western hemlock—Salal	03	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Douglas-fir – Western hemlock – Salal sites occurred on well-drained, nutrient very poor to medium upper slopes with mor humus types. Substrate was generally till of various depths, infrequently co-occurring with colluvium. The canopy closure of structural stage 4-5 stands was dense (often greater than 50%), opening up with succession. Dominant tree species included Douglas-fir, western redcedar and western hemlock. Regeneration of hemlock tended to be abundant in the understorey, with little Douglas-fir regeneration, reflecting the low shade tolerance of Douglas-fir and its mineral seedbed requirement. Salal formed dense thickets and was a constant dominant in the shrub layer, with little other plant cover except dull Oregon grape as a frequent associate. Step moss, Oregon beaked-moss, electrified cat's tail moss and curly heron's bill moss were common in the bryophyte layer.</p> <p>Assumed modifiers: d, m, w</p>		Elevation (m)	130-600
		Slope (%)	5-100
		Aspect (°)	variable
		Surficial material	M (C)
		Drainage	w-r
		SMR	1-2
		SNR	A-C
		Plots: 6105, 6107, CAG022, CA057, CA076, CHG068, CHV069, GBG026, JKV013, JK301, JK302, MTV059, WM001, WM002, WM003, WM042, ERSK_GR15, WM043, ERSK_GR13	

TEM Map Code	Site Association		CWHxm Site Series
DS	Douglas-fir—Western hemlock—Salal		03
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE			
DS4 (Immature Forest)	DS5 (Young Forest)	DS6 (Mature Forest)	Mature Undisturbed DS6
Dominant Species salal Douglas-fir Oregon beaked-moss arbutus dull Oregon-grape	Dominant Species Douglas-fir oceanspray dull Oregon-grape salal Oregon beaked-moss western redcedar	Dominant Species Douglas-fir step moss curly heron's-bill moss Oregon beaked-moss salal western redcedar western hemlock dull Oregon-grape knight's plume sibbaldia	Tree Layer Douglas-fir western hemlock western redcedar shore/lodgepole pine Shrub Layer salal red huckleberry dull Oregon-grape baldhip rose
Associate Species western hemlock oceanspray step moss Siberian miner's-lettuce trailing blackberry hairy honeysuckle	Associate Species electrified cat's-tail moss step moss arbutus heron's-bill moss twinflower	Associate Species golden short-capsuled moss juniper haircap moss red-stemmed feathermoss western fescue yellow curl-moss	Herb Layer twinflower bracken fern vanilla-leaf sword fern Moss Layer step moss Oregon beaked moss lanky moss electrified cat's-tail moss

TEM Map Code	Site Association		CWHxm Site Series
DS	Douglas-fir—Western hemlock—Salal		03
DS _{hs}	hummocky; shallow soil	DS _{kv}	cool aspect; very shallow soil
DS _{hv}	hummocky; very shallow soil	DS _{qv}	very steep cool aspect; very shallow soil
DS _j	gentle slope	DS _{rs}	ridge; shallow soil
DS _{ju}	gentle slope, very shallow soil	DS _{rv}	ridge; very shallow soil
DS _k	cool aspect	DS _s	shallow soil
DS _{ks}	cool aspect; shallow soil	DS _{sz}	shallow soil; very steep warm aspect
Atypical Characteristics/Additional Comments: N/A			


TEM Map Code	Site Association	CWHxm Site Series	
DF	Douglas-fir—Sword fern	04	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Douglas-fir – Sword fern ecosystems were mapped on well drained upper to mid-slopes with variable thicknesses of till, typically of finer texture than sites supporting site series 03. Humus forms were mulls to moders. Douglas-fir was the most common canopy dominant, but western hemlock was a frequent associate in all canopy layers, although sparser than expected on Saltspring Island. Grand fir was an infrequent associate. Shrubs occupied low to moderate cover (15-30%), with dull Oregon-grape, red huckleberry, common snowberry, and trailing blackberry constant associates. Sword fern dominated the herb layer, with relatively few other species. The bryophyte layer was dominated by Oregon beaked moss.</p> <p>Assumed modifiers: d, j, m</p>		Elevation (m)	140-575
		Slope (%)	35-55
		Aspect (°)	variable
		Surficial material	M
		Drainage	w
		SMR	1-2
		SNR	C-E
		Plots: GB028, GBV031, BBJK11, CA079, WM030, WM031, ERSK_GR07, JK305, K30	

TEM Map Code	Site Association				CWHxm Site Series
DF	Douglas-fir—Sword fern				04
DFhs	hummocky; shallow soil	DFkv	cool aspect; very shallow soil	DFsz	shallow soil; very steep warm aspect
DFhv	hummocky; very shallow soil	DFqs	very steep cool aspect; shallow soil	DFw	warm aspect
DFk	cool aspect	DFs	shallow soil		
DFks	cool aspect; shallow soil	DFsw	shallow soil; warm aspect		
Atypical Characteristics/Additional Comments: N/A					

TEM Map Code	Site Association	CWHxm Site Series	
RS	Western redcedar—Sword fern	05	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Western redcedar – Sword fern ecosystem was mapped most often on north-facing mid-slopes with well to moderately well-drained soils derived from till and occasionally colluvium. Western redcedar was not always dominant in the canopy, likely reflecting the disturbance history of sites in the study area. Western hemlock, grand fir, and Douglas-fir were frequent, represented in all canopy layers. Coarse woody debris was often moderately abundant to abundant on these sites. The understory was dominated by salal, with variable amounts of salmonberry, and occasionally dull Oregon-grape in the shrub layer. The herb layer featured the dominant species sword fern and bracken fern, occasionally with spiny wood fern and introduced grass species. Both bracken fern and graminoids decreased with increasing canopy closure. The moss layer was dominated by Oregon beaked moss with lesser amounts of constant associates curly heron's bill moss, knight's plume, and with coastal leafy moss on decaying wood. Some sites were influenced by seepage, and developed this ecosystem on warm aspects or shallower soils than typical.</p> <p>Assumed modifiers: d, m</p>		<p>Elevation (m)</p>	125-550
		Slope (%)	10-100
		Aspect (°)	285-135 (135-285)
		Surficial material	M, C
		Drainage	w-m
		SMR	3-4
		SNR	D (E)
		Plots: CHG074, GB029, WM013	


TEM Map Code	Site Association	CWHxm Site Series
RS	Western redcedar—Sword fern	05
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE		
RS3 (Shrub/Herb)		Mature Undisturbed RS6
Dominant Species salmonberry Oregon beaked-moss bracken fern red alder dull Oregon-grape salal Scotch broom trailing blackberry western redcedar Himalayan blackberry Associate Species red huckleberry western hemlock		Tree Layer Douglas-fir western hemlock western redcedar red alder Shrub Layer red huckleberry salal dull Oregon-grape baldhip rose oceanspray Herb Layer sword fern vanilla-leaf bracken fern twinflower wall-lettuce sweet-scented bedstraw three-leafed foam flower Moss Layer step moss Oregon beaked moss lanky moss coastal leafy moss

TEM Map Code		Site Association		CWHxm Site Series	
RS		Western redcedar—Sword fern		05	
RShj	hummocky; gentle slope	RSks	cool aspect; shallow soil	RSsw	shallow soil; warm aspect
RSj	gentle slope	RSkv	cool aspect; very shallow soil	RSvw	very shallow soil; warm aspect
RSjs	gentle slope; shallow soil	RSqv	very steep cool aspect; very shallow soil	RSw	warm aspect
RSk	cool aspect	RSs	shallow soil	RSwx	warm aspect; drier than typical
Atypical Characteristics/Additional Comments: N/A					

TEM Map Code	Site Association	CWHxm Site Series	
HD	Western hemlock—Western redcedar—Deer fern	06	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Western hemlock – Western redcedar – Deer fern site tended to develop on north-facing mid- to lower slopes with moderately to imperfectly drained Brunisols and Podzols derived from till. Douglas-fir and western hemlock were the most abundant tree and constant dominant, with lesser amounts of frequent associates western redcedar and grand fir. Bigleaf maple and red alder were often present in seral stands. Diagnostic species deer fern was infrequent in these stands. The understorey featured salal, salmonberry, and dull Oregon-grape in the shrub layer, with sword fern, bracken dominants in the herb layer. Occasionally spiny wood fern and introduced grass species also occurred. The moss layer was dominated by Oregon beaked moss with lesser amounts of associated species curly heron's bill moss and knight's plume. Coastal leafy moss was common on decaying wood.</p> <p>Assumed modifiers: d, j, m</p>		Elevation (m)	250-575
		Slope (%)	15-45
		Aspect (°)	variable
		Surficial material	M
		Drainage	w-m
		SMR	5-6
		SNR	A-C
		Plots: JK016, JK017, MT058, QAF005, TTG005, WM037	


TEM Map Code	Site Association	CWHxm Site Series
HD	Western hemlock—Western redcedar—Deer fern	06
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE		
<div>HD5 (Young Forest)</div>		<div>Mature Undisturbed HD6</div>
Dominant Species Douglas-fir baneberry sweet-cicely Sitka columbine Oregon beaked-moss Columbia brome western hemlock Associate Species bedstraw Nootka rose prickly rose sword fern wild strawberry		Tree Layer Douglas-fir western hemlock western redcedar grand fir red alder Shrub Layer salal red huckleberry false azalea dull Oregon-grape salmonberry Herb Layer deer fern sword fern Moss Layer step moss lanky moss Oregon beaked moss

TEM Map Code	Site Association		CWHxm Site Series
HD	Western hemlock—Western redcedar—Deer fern		06
HDh	hummocky	HDs	shallow soil
HDk	cool aspect	HDsw	shallow soil; warm aspect
HDks	cool aspect; shallow soil	HDw	warm aspect
HDkv	cool aspect; very shallow soil		
Atypical Characteristics/Additional Comments:			
N/A			


TEM Map Code	Site Association	CWHxm Site Series	
RF	Western redcedar—Foamflower	07	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>The Western redcedar - Foamflower ecosystems occurred on moisture-receiving toe slopes, some seepage sites, and level sites with thick relatively rich, moderately well to imperfectly drained soils. Common soil types were Humo-Ferric Podzols or occasionally Gleysols on level sites grading to gleyed Brunisols derived from till. Humus forms ranged from moders to mulls. Western redcedar was often associated in the semi-open canopy with constant associates western hemlock, red alder, grand fir and bigleaf maple; Douglas-fir was infrequent to absent on most sites. Hemlock also was infrequent in the study area on these site types. Trees on these productive sites tended to be larger their cohorts in most other site series. Shrubs included dense cover of the dominant species dull Oregon-grape, salmonberry and thimbleberry. Herbs also occupied a high cover proportion, with ladyfern, spiny wood fern, foamflower, vanilla-leaf, oak fern, and small-flowered rush typically present in varying amounts on most sites. Step moss, curly heron's-bill moss, lanky moss and Oregon-beaked moss was dominant in the bryophyte layer.</p> <p>Assumed modifiers: d, j, m</p>		Elevation (m) Slope (%) Aspect (°) Surficial material Drainage SMR SNR	 120-325 0-35 variable M m 5-6 D-E
		Plots: JKG014, TTG008, WM040	

TEM Map Code	Site Association	CWHxm Site Series
RF	Western redcedar—Foamflower	07
CHARACTERISTIC VEGETATION WITHIN EACH STRUCTURAL STAGE		
RF4 (Immature Forest)	RF6 (Mature Forest)	Mature Undisturbed RF6
Dominant Species stinging nettle red alder coastal leafy moss small-flowered bitter-cress Associate Species dovefoot geranium red-stemmed feathermoss arbutus bedstraw bigleaf maple Douglas-fir field chickweed Nootka rose Oregon beaked-moss Pacific bleeding heart Scouler's corydalis sweet-cicely	Dominant Species red alder western hemlock western redcedar step moss curly heron's-bill moss knight's plume lanky moss Oregon beaked-moss salal Associate Species dull Oregon-grape flat-moss stinging nettle	Tree Layer Douglas-fir western hemlock western redcedar grand fir red alder bigleaf maple Shrub Layer dull Oregon-grape red huckleberry salmonberry devil's club Herb Layer sword fern vanilla-leaf wall-lettuce three-leaved foamflower sweet-scented bedstraw bracken fern lady fern oak fern Moss Layer step moss coastal leafy moss Oregon beaked moss palm tree moss lanky moss


TEM Map Code	Site Association	CWHxm Site Series
RF	Western redcedar—Foamflower	07
RFh hummocky RFk cool aspect RFs shallow soil RFw warm aspect Atypical Characteristics/Additional Comments: N/A		

TEM Map Code	Site Association	CWHxm Site Series	
SS	Sitka spruce—Salmonberry	08	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Sitka spruce - Salmonberry high bench site series was infrequently mapped in the study area. Fluvial materials and landscape position contribute to highly productive, moderately-well drained sites with relatively open canopies (more open as stands age) Soils were frequently well-developed with moder to mull humus forms. Sitka spruce was not found in the study area; broadleaf species including black cottonwood, red alder and bigleaf maple were dominant, with varying amounts of frequent associate western redcedar. Shrubs were highly variable in cover and diversity, with salmonberry and thimbleberry dominating, and less abundant associates' ninebark, red-osier dogwood, cascara, red elderberry, bitter cherry, and Pacific crabapple. Herbs also varied, with relatively low cover compared to shrubs. The common associate species were vanilla-leaf, ladyfern, sweet-cicely, and false lily-of-the-valley. Season of sampling also influenced cover and species recorded. Coastal leafy moss was a common colonizer on woody substrate.</p> <p>Assumed modifiers: d, j, m</p>		Elevation (m)	150-350
		Slope (%)	5-35
		Aspect (°)	variable
		Surficial material	F
		Drainage	m
		SMR	5-6
		SNR	C-E
		Plots: WM040	


TEM Map Code	Site Association	CWHxm Site Series	
CW	Black cottonwood—Willow	10	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Low fluvial benches and floodplain sites support the Black cottonwood - Willow site series, featuring deep coarse-textured gravelly sandy parent material with relatively high coarse fragment content. Soils vary from Non-soil in recently disturbed locations to gravelly sandy Regosols and Brunisols. This ecosystem type was extremely infrequent on Saltspring Island. Frequent inundation limits the dominant species to red alder, willows (more often in or adjacent to inundation zones) and black cottonwood, with associated flood-tolerant shrubs such as salmonberry. Canopy closure varies with flood regime, terrain, and seral stage. Structural stage 4 supports the densest stands (canopy closure approximately 50%) with fluvial erosion contributing to stand disturbance. This species association is a disturbance-maintained disclimax that would support conifer species in the absence of disturbance. Herbs and mosses are infrequent or absent depending on water table and fluvial characteristics, with blue wildrye most common associate in less frequently flooded sites. Soils are well drained, but the coarse texture limits productivity on sandstone-derived soils, and flooding action often precludes development of an organic soil horizon. Erosion is a typical disturbance agent, leading to a predominance of younger seral stands (structural stages 4-5) in the study area.</p> <p>Assumed modifiers: a, c, d, j</p>		<p>Elevation (m)</p>	150-500
		Slope (%)	<15
		Aspect (°)	variable
		Surficial material	F ^A
		Drainage	w - m
		SMR	5-6
		SNR	C-D
No photo available		Plots: JK136	

TEM Map Code	Site Association	CWHxm Site Series	
RC	Western redcedar—Sitka spruce—Skunk cabbage	12	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar – Sitka spruce – Skunk cabbage are rich, moist to wet sites which occurred in moisture-receiving depressions with fine to medium soils, possibly including a minor component of organics. Although Sitka spruce was not observed on Saltspring Island, western redcedar was a typical dominant with lesser amounts of associated red alder, bigleaf maple, and grand fir on the margins. Alder abundance decreased with stand age. Sites often had abundant moderate to large woody debris. Shrubs observed included salmonberry, thimbleberry, and occasionally salal on hummocks. Species visible in the herb layer were limited by the sampling window, but the most common associate species were vanilla-leaf, lady fern, deer fern, and skunk cabbage. Bryophytes associated with this site series were coastal and large leafy moss and slender beaked moss.</p> <p>Assumed modifiers: d, j, m</p>		Elevation (m)	120-450
		Slope (%)	0-10
		Aspect (°)	variable
		Surficial material	M (0)
		Drainage	p
		SMR	7
		SNR	C-E
		Plots: N/A	


TEM Map Code	Site Association	CWHxm Site Series
RC	Western redcedar—Sitka spruce—Skunk cabbage	12
RCp peaty material RCs shallow soil Atypical Characteristics/Additional Comments: N/A		


TEM Map Code	Site Association	CWHxm Site Series	
RB	Western redcedar—Salmonberry	13	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar – Salmonberry ecosystems are nutrient medium- to-rich, very moist (winter) to fresh (summer) sites and were fairly common in hummocky, imperfectly drained, level terrain or moisture receiving toe slopes as well as in riparian areas. These sites tended to support a mosaic of site series 13 and 07, with the former in hollows and the latter on raised microsites. Soils were often gleyed or weakly mottled Brunisols (typically Gleyed Eluviated Dystric Brunisols, similar to CWHxm/12 but with more pronounced mottling), reflecting the seasonally fluctuating water tables. There were accumulations of fines beneath the organic horizons in depressions. The canopy was open (5-15%), with western redcedar the dominant tree species growing on elevated hummocks. Dense thickets of salmonberry were the dominant shrub component, with occasional Douglas maple, thimbleberry, and red elderberry. Herbs were variable, but few were apparent during sampling (largely in winter). Sword fern grew on hummocks and bleeding heart and sweet-cicely were also common associates. Bryophytes were sparse to absent except on decaying wood.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m)</p>	120-450
		Slope (%)	0-10
		Aspect (°)	variable
		Surficial material	M
		Drainage	i
		SMR	5
		SNR	C-D
		Plots: N/A	

TEM Map Code	Site Association	CWHxm Site Series
RB	Western redcedar—Salmonberry	13
RBh hummocky RBhs hummocky; shallow soil RBhv hummocky; very shallow soil RBks cool aspect; shallow soil RBs shallow soil Atypical Characteristics/Additional Comments: N/A		

TEM Map Code	Site Association	CWHxm Site Series	
RT	Western redcedar—Black twinberry	14	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western recedar – Black twinberry sites are rich, wet (winter) to moist (summer) sites were infrequent, occurring in hummocky, imperfectly drained, level terrain or moisture receiving toe slopes and in riparian areas. These sites tended to support a mosaic of site series 14 and 07 or 12, with the former in hollows and the latter on raised microsites. The canopy was open to sparse, with western redcedar the dominant tree species growing on elevated hummocks. Shrubs had fairly high cover, with salmonberry, black twinberry, red elderberry, and willows the most common dominants, and occasional components of ninebark and Pacific crabapple associated with richer sites. Herbs varied among sites; foamflower, lady fern, vanilla-leaf, and sword fern were the most frequent. Bryophytes were sparse to absent except on decaying wood.</p> <p>Assumed modifiers: d, j, m</p>		Elevation (m)	200-450
		Slope (%)	0-15
		Aspect (°)	variable
		Surficial material	M, O
		Drainage	i
		SMR	6
		SNR	D-E
		Plots: N/A	


TEM Map Code	Site Association	CWHxm Site Series
RT	Western redcedar—Black twinberry	14
RTs shallow soil Atypical Characteristics/Additional Comments: N/A		

TEM Map Code	Site Association	CWHxm Site Series	
CS	Western redcedar—Slough sedge	15	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Western redcedar – Slough sedge are swamp forests which occupy poorly drained flat sites to depressions. Western redcedar and sword fern dominate elevated microsites, while sedges, lady fern, and horsetails occupy hollows with occasional skunk cabbage where there is limited surface flow. Soils are moderately deep to deep (>0.5m) with medium texture, typically gleyed, with seasonally fluctuating water tables, even where bedrock restricts soil depth. Tree species are limited to shade- and moisture-tolerant trees with relatively shallow roots: western redcedar, grand fir on margins, and black cottonwood with minor amounts of red alder. Shrubs in this site series are diverse, with Indian-plum, common snowberry, currants/gooseberry, and thimbleberry the most frequent associates; ninebark, black twinberry, and red-osier dogwood increased in frequency with increased seasonal flooding. Herbaceous species were variable, with slough sedge commonly dominating the herb layer, and Cooley's hedge-nettle in more well-drained and small-flowered rush in more poorly-drained sites also common associates. Moss tended to occupy little of the substrate, and only Oregon beaked moss was a constant associate; large leafy moss and coastal leafy moss were occasionally present.</p> <p>Assumed modifiers: d, j, m</p>		<p>Elevation (m)</p>	150-450
		Slope (%)	0-35
		Aspect (°)	variable
		Surficial material	Ov, M, L
		Drainage	m-p
		SMR	6-7
		SNR	C-E
		Plots: CA008, CA039, MT036	


Tem Map Code	Site Association	CWHxm Site Series	
AM	Arbutus—Hairy manzanita	00	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>These very xeric Arbutus – Hairy Manzanita sites occurred on ridge crests and water-shedding steep upper slopes directly on bedrock. Canopies were very open (<5%) with arbutus and occasionally shore pine dominant, rarely reaching past structural stage 4 with respect to structural development criteria. The shrub layer was dominated by hairy manzanita, arbutus regeneration, and occasionally Scotch broom. Herbs were typically sparse, but annuals may be evident in the spring, particularly graminoids, but comprising low (<20%) cover. The substrate was typically dominated almost completely by curly heron's-bill moss and, to a lesser extent, and on rock outcrops, foliose lichens.</p> <p>From plot data for a shrub/herb (structural stage 3) dominated Arbutus – Hairy Manzanita site the dominant vegetation included sweet vernalgrass, hairy Manzanita, shore pine, curly heron's-bill moss, hoary rock-moss, Douglas-fir, Alaska oniongrass, grey rock-moss and juniper haircap moss. Associate species included green sorrel and lesser green reindeer. From plot data for a young forest (structural stage 5) site the dominant vegetation included arbutus, curly heron's-bill moss, red-stemmed feathermoss, hairy Manzanita, western hemlock, Douglas-fir, Oregon beaked-moss and salal, with oceanspray as an associate species.</p> <p>Assumed modifiers: j, r, s</p>		Elevation (m) Slope (%) Aspect (°) Surficial material Drainage SMR SNR	Variable Variable 120-250 R r 1 A
		Plots: 4839, ERSK_GR01, WMG006	

Tem Map Code	Site Association	CWHxm Site Series
AM	Arbutus—Hairy manzanita	00
AMv very shallow soil AMvw very shallow soil; warm aspect AMvz very shallow soil; very steep warm aspect AMw warm aspect AMz very steep warm aspect Atypical Characteristics/Additional Comments: N/A		


Distribution of Map Units Containing AM Units




Anthropogenic and Non-Vegetated/Sparsely Vegetated Map Units of the CDFmm


TEM Map Code	Site Unit Name	Site Series	
BE	Beach	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Beach units are characterized by sorted sediments reworked by wave action. All beach units were mapped along the ocean edge, beach units were not present along fresh water bodies. This unit is typically void of vegetation and consists of either sand or coarse fragments such as gravels, cobbles and stones, with scattered driftwood, seaweed and various washed up items along the shoreline.</p>		Elevation (m):	0-3 m
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	W
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: N/A	


TEM Map Code	Site Unit Name	Site Series	
CF	Cultivated Field	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Cultivated fields are non forested, open areas that are subject to agricultural practices including plowing, fertilization, and non native crop production which often results in long-term soil and vegetation changes. Cultivated fields are common throughout the CDFmm, and extensive throughout the Cowichan Valley and Central and North Saanich. The typical structural stage for cultivated fields is 2b, graminoid, infrequently 3. Small islands of forest occur in fields, as do narrow shrub dominated riparian channels and scattered large trees. Noteworthy are the cultivated fields of Cowichan and Saanich with scattered Garry oak trees and Garry oak groves. Rural residential is also a very common association with cultivated fields. Although cultivated fields are typically agriculture based CF was mapped for other green spaces as well. These included city parks, baseball fields, and residential lawns. This unit was often complexed with Rural (RW) and Urban (UR) units.</p>		Elevation (m):	50-150
		Slope (%):	<15
		Aspect (°):	Variable
		Surficial material:	W ^G , W (Ox)
		Drainage:	i - w
		SMR:	2-6
		SNR:	B-D
		Plots: BB_HR_G1, BB_HR_G2, BB_HR_G3, BB_HR_G6, DHV022, GBG044, GBV055, CHV093, JCV101, JCV060, DHV050, HRV077, HRV209	

TEM Map Code	Site Unit Name	Site Series	
CL	Cliff	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Cliffs are steep, vertical or overhanging rock faces. Cliff units are most common along the coast complexed with coastal bluffs or steep rock controlled gully sidewalls and infrequently along mountain slopes. Cliffs are non-vegetated ecosystems.</p>		Elevation (m):	Variable
		Slope (%):	>130
		Aspect (°):	Variable
		Surficial material:	R
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: JCV036	

TEM Map Code	Site Unit Name	Site Series	
CO	Cultivated Orchard	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Agricultural areas composed of single or multiple tree species planted in rows. Typically this would include, for example, apple orchards or mixed species orchards. Old homesteads are common in the CDF and although many of them have become overgrown, these have been mapped as CO wherever possible. It also includes tree farms such as Christmas tree farms, or any coniferous tree farm where conifer trees are planted in rows for commercial production.</p>		Elevation (m):	Variable
		Slope (%):	<15
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: N/A	

TEM Map Code	Site Unit Name	Site Series	
CV	Cultivated Vineyard	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Cultivated vineyards are agricultural areas composed of single or multiple species of grapes planted in rows, usually supported on wood or wire trellises. Although in the grand scheme of things within the mapping area, cultivated vineyards are not common, they do occur in the Cowichan Valley and to a less extent in Central Saanich. Wine vineyards quite often do not occur as a major polygon component, instead they are a minor component complexed with cultivated fields and rural residential. Occasionally cultivated vineyards have been mapped as pure polygons.</p>		Elevation (m):	Variable
		Slope (%):	<15
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
No photo available		SNR:	n/a
		Plots: N/A	


TEM Map Code	Site Unit Name	Site Series	
ES	Exposed Soil	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Exposed soil is any area of exposed soil that is devoid of vegetation and is not bedrock but mineral soil. Exposed soil was most often mapped near rural and urban areas, predominantly disturbance sites associated with new housing developments. Exposed soil was also mapped for pull-outs along forestry roads and to a lesser extent along steep side slopes. For example steep side slopes composed of glaciomarine sediments, were mapped as exposed soil. Due to surface erosion, vegetation was not present.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	F ^G , W ^G , A
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: CHV108, CHV166, HRV106, JCV160	


TEM Map Code	Site Unit Name	Site Series	
GB	Gravel Bar	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Gravel bars typically occur along active river channels. Gravel bars are non-vegetated and characterized by a dynamic consortium of well rounded cobbles, pebbles, stones and sand. Notable rivers with gravel bars include Englishman River, Chemainus River, Cowichan River and Koksilah River.</p>		Elevation (m):	Variable
		Slope (%):	<15
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: N/A	

TEM Map Code	Site Unit Name	Site Series	
GC	Golf Course	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Golf courses are a common feature particularly along southeastern Vancouver Island - a favorite pastime of island dwellers. Golf courses are characterized by flat to gently rolling grass-covered throughways and open areas for playing golf. The fairways are typically separated by isolated rows or patches of trees and ponds. More often than not, forested areas were too small to identify to site series and were mapped as GC with a structural stage 5 or 6. Although golf courses have altered the native landscape and vegetation communities to a large extent, scattered trees that remain are noteworthy. For example, scattered trees found in golf courses of the Victoria area were large diameter and of significant height and age whereas other golf course were dotted with healthy specimens of Parkland Garry Oak and arbutus.</p>		Elevation (m):	Variable
		Slope (%):	<15
		Aspect (°):	Variable
		Surficial material:	W ^G
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
No photo available		Plots: JCV069, HRV113, HRV215	


TEM Map Code	Site Unit Name	Site Series	
GP	Gravel Pit	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
Gravel pits are areas of exposed soil through the commercial removal of sand and gravel.		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	W ^G , F ^G , A
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
No photo available		Plots: CAG029, CHV072, JCV062, HRV034, HRV059	

TEM Map Code	Site Unit Name	Site Series	
IN	Industrial	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
Industrial sites were added as an anthropogenic unit due to the requirement to address sites that were dominated by industrial development namely, pulp and paper, lumber mills, oil/gas refineries and so on. These units are characterized by a high degree of ground disturbance; concrete parking lots, large commercial buildings, work yards and other specialized industry infrastructure.		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
No photo available		Plots: N/A	

TEM Map Code	Site Unit Name	Site Series	
LA	Lake	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>A lake is a naturally occurring body of water, greater than 2 m deep. Several lakes occur within the mapping area such as Elk, Beaver, Prospect, Cowichan, Quamichan, Somenos, Long and Brannen Lake to name a few.</p>		Elevation (m):	Variable
		Slope (%):	0
		Aspect (°):	999
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: N/A	


TEM Map Code	Site Unit Name	Site Series	
MI	Mine	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Mine sites are un-vegetated areas used for the purpose of extracting mineral ore or other materials. Mine sites were mapped in the study area, some of which were limestone quarries.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	A
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: JCV023	


TEM Map Code	Site Unit Name	Site Series	
MU	Mudflat Sediment	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
Mudflat sediment is a flat plain-like area dominated by fine textured sediments. These areas were mapped at the interface of salt and fresh water, namely active channel sediment located at the mouth of a creek as they drain into saltwater bays.		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
No photo available		SNR:	n/a
		Plots: N/A	

TEM Map Code	Site Unit Name	Site Series	
OW	Shallow Open Water	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Shallow open water is a wetland class composed of permanent, shallow (less than 2 m at midsummer levels), standing water that has less than 10% surface cover of emergent vegetation (plants rooted in the bottom). Open water with more than 10% surface cover of emergent vegetation are classified as marsh wetlands. Open water is typically a minor polygon component in association with other non-forested wetlands such as marsh ecosystems (e.g. Wm05 & Wm06) and swamp forests (Cw-Skunk cabbage (11) and Cw-Slough sedge (14)).</p>		Elevation (m):	Variable
		Slope (%):	0
		Aspect (°):	999
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: DHV013, DHV039	


TEM Map Code	Site Unit Name	Site Series	
PD	Pond	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Ponds are small bodies of water greater than 2 m deep yet smaller than lakes (less than 50 ha). Ponds are scattered across the landscape and were often surprisingly difficult to distinguish between natural ponds and man-made impoundments. Man-made impoundments were typically mapped as reservoirs however berms & dykes were often not readily visible and left some doubt as to their origin. Ponds are a common feature throughout the rural residential areas and in association with cultivated fields.</p>		Elevation (m):	Variable
		Slope (%):	0
		Aspect (°):	999
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
No photo available		Plots: N/A	

TEM Map Code	Site Unit Name	Site Series	
RE	Reservoir	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>A reservoir is an artificial basin created by the impoundment of water behind a human-made structure such as a dam, berm, dyke, or wall. As noted above, reservoirs were often difficult to distinguish from ponds. Reservoirs most commonly occur throughout cultivated fields and rural residential areas.</p>		Elevation (m):	Variable
		Slope (%):	0
		Aspect (°):	999
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
No photo available		Plots: N/A	


TEM Map Code	Site Unit Name	Site Series	
RI	River	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>A river is a natural stream of water, usually freshwater, flowing toward the ocean, a lake, or another stream. In some cases a river flows into the ground or dries up completely before reaching another body of water. The water in a river is usually confined to a channel, made up of a stream bed between banks. Notable rivers include Englishman River, Chemainus River, Cowichan River and Koksilah River.</p>		<p>Elevation (m):</p>	Variable
		<p>Slope (%):</p>	Variable
		<p>Aspect (°):</p>	Variable
		<p>Surficial material:</p>	n/a
		<p>Drainage:</p>	n/a
		<p>SMR:</p>	n/a
		<p>SNR:</p>	n/a
		Plots: N/A	


TEM Map Code	Site Unit Name	Site Series	
RN	Railway Surface	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Railway surface is a long, linear feature that is characterized by a level surface with fixed rails to carry trains. The Esquimalt and Nanaimo (E&N) railway dissects a portion of the mapping area on southeastern Vancouver Island. The E&N is a short line railway that runs from Victoria to Courtenay, with branch lines to Parksville and south of Nanaimo to the E&N's main railyard and barge slip (dock) on the Nanaimo waterfront.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
			

Plots: N/A


TEM Map Code	Site Unit Name	Site Series	
RO	Rock Outcrop	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Rock outcrops are bedrock outcroppings with little soil development and sparse vegetation cover. Rock outcrops can be steep such as the one in the photo below or gentle such as those found along ridges. Often rock outcrops are associated with dry O2 sites or QB and SC units. Most importantly SEI units classified as coastal bluffs were mapped as rock outcrops with aspect modifiers. These coastal rock outcrops begin at the waters edge and are exposed to salt spray and high winds.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	R (Mvx)
		Drainage:	x-r
		SMR:	n/a
		SNR:	n/a
		Plots: CORV003, JCV086, JCV251, CHV261, JCG004, JCV008, JCV012, CXV034	

TEM Map Code	Site Unit Name	Site Series	
RW	Rural	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Rural areas are characterize by areas that have residences and other human development scattered and intermingled with forests, range, farm land, cultivated fields or native vegetation. Rural areas are very common in the CDF and often form the dominant polygon component. Minor components include cultivated fields, zonal forests and shrubby riparian creek draws.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: JCV032B, DHV043, DHV058, HRV010, HRV012, WMG021	

TEM Map Code	Site Unit Name	Site Series	
RZ	Road Surface	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Road surfaces are defined as areas cleared and compacted for the use of vehicles. Road surfaces are linear features that cross the landscape and occur in varying degrees. From small farms roads, to gravel forestry roads, to two-lane residential paved roads & highways and to main thoroughfares such as the Island highway (Hwy 1 & 19) with a significant portion being four-lanes. The Island highway was for the most part, pulled out and mapped as its own long, narrow and continuous corridor. Other, lesser roads were not.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: CHV098	

TEM Map Code	Site Unit Name	Site Series	
TA	Talus	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
Talus is angular, colluvial rock fragments that have accumulated at the foot of steep rock slopes. Talus ecosystems are not a common feature of the CDFmm.		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	C
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: N/A	

TEM Map Code	Site Unit Name	Site Series	
TZ	Mine Tailings	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Mine tailings are solid waste materials directly produced and deposited during the mining and milling of ore bodies. Mine tailings are not common however did occur on occasion.</p> <p>For example, near Ladysmith there is an old site where coal was hauled in, then exported out. There is a thick layer of coal debris underneath with disturbed shrub vegetation.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
No photo available		Plots: N/A	

TEM Map Code	Site Unit Name	Site Series	
UR	Urban	N/A	
SITE DESCRIPTION		SITE CHARACTERISTICS	
<p>Urban units are characterized by an almost continuous covering over the landscape by residences and human development. The greater Victoria area is the most significant urban area mapped. Given the extensive cover of concrete and altered landscape, bioterrain labels were intended to reflect this by labeling urban polygons Anthroprogenic (A). Whereby the landform is man-modified and drainage is no longer applicable. Other urban areas occurred in Duncan, Crofton, Ladysmith, Nanaimo, Parksville, Qualicum and Powell River for example. Urban units most often formed pure polygons however sometimes patches of forest are present as a minor component (10%). For example the QB unit was most often found to occur in urban areas of Victoria and Saanich while the DS unit more often occurred as a minor component with urban centers up island.</p>		Elevation (m):	Variable
		Slope (%):	Variable
		Aspect (°):	Variable
		Surficial material:	n/a
		Drainage:	n/a
		SMR:	n/a
		SNR:	n/a
		Plots: JCV031, HRV137	



APPENDIX VII

Vascular and Non-Vascular Plant Species Observed During Field Assessments in the CDFmm Study Area

Appendix VII. Vascular and Non-Vascular Plant Species Observed During Field Assessments in the Study Area

<i>Abies amabilis</i>	amabilis fir
<i>Abies grandis</i>	grand fir
<i>Acer macrophyllum</i>	bigleaf maple
<i>Achillea millefolium</i>	yarrow
<i>Achlys triphylla</i>	vanilla-leaf
<i>Adenocaulon bicolor</i>	pathfinder
<i>Agoseris grandiflora</i>	large-flowered agoseris
<i>Agrostis scabra</i>	hair bentgrass
<i>Agrostis</i> sp.	bentgrass
<i>Agrostis stolonifera</i>	creeping bentgrass
<i>Aira caryophylla</i>	silver hairgrass
<i>Aira praecox</i>	early hairgrass
<i>Allium amplexans</i>	slimleaf onion
<i>Allium cernuum</i>	nodding onion
<i>Alnus rubra</i>	red alder
<i>Amelanchier alnifolia</i>	saskatoon berry
<i>Anthoxanthum odoratum</i>	sweet vernalgrass
<i>Aphanes arvensis</i>	field aphanes
<i>Arabis hirsuta</i>	hoary rock-cress
<i>Arbutus menziesii</i>	arbutus
<i>Arctostaphylos columbiana</i>	hairy manzanita
<i>Arctostaphylos uva-ursi</i>	kinnikinnick
<i>Asarum caudatum</i>	wild ginger
<i>Asplenium trichomanes</i>	common spleenwort
<i>Aster</i> sp.	Aster sp.
<i>Athyrium filix-femina</i>	lady fern
<i>Atriplex</i> sp.	Orache
<i>Bellis perennis</i>	English daisy
<i>Blechnum spicant</i>	deer fern
<i>Brachythecium frigidum</i>	golden short-capsuled moss
<i>Bromus carinatus</i>	California brome
<i>Bromus hordeaceus</i>	soft brome
<i>Bromus vulgaris</i>	Columbia brome
<i>Bryum</i> sp.	Bryum sp.
<i>Calamagrostis canadensis</i>	bluejoint reedgrass
<i>Callitriche</i> sp.	water starwort
<i>Calypso bulbosa</i>	fairy-slipper
<i>Camassia leichtlinii</i>	great camas
<i>Camassia quamash</i>	common camas
<i>Campanula rotundifolia</i>	common harebell
<i>Cardamine occidentalis</i>	western bitter-cress
<i>Cardamine oligosperma</i>	little western bitter-cress
<i>Carex aquatilis</i>	water sedge
<i>Carex deweyana</i>	Dewey's sedge
<i>Carex lasiocarpa</i>	slender sedge
<i>Carex obnupta</i>	slough sedge
<i>Carex sitchensis</i>	Sitka sedge
<i>Castilleja hispida</i>	harsh paintbrush



<i>Cerastium arvense</i>	field chickweed
<i>Cerastium vulgatum</i>	mouse-ear chickweed
<i>Chimaphila umbellatum</i>	prince's pine
<i>Cirsium arvense</i>	Canada thistle
<i>Cirsium vulgare</i>	bull thistle
<i>Cladina mitis</i>	lesser green reindeer lichen
<i>Cladina portentosa</i>	coastal reindeer lichen
<i>Cladina rangiferina</i>	grey reindeer lichen
<i>Cladonia chlorophaea</i>	peppered pixie-cup
<i>Claytonia perfoliata</i>	miner's lettuce
<i>Claytonia sibirica</i>	Siberian spring beauty
<i>Claytonia spathulata</i>	pale montia
<i>Collinsia parviflora</i>	small-flowered blue-eyed Mary
<i>Cornus nuttallii</i>	western flowering dogwood
<i>Crataegus douglasii</i>	black hawthorn
<i>Crepis tectorum</i>	annual hawksbeard
<i>Cytisus scopulorum</i>	Scotch broom
<i>Dactylis glomerata</i>	orchardgrass
<i>Daphne laureola</i>	spurge-laurel
<i>Dicranum fuscescens</i>	curly heron's-bill moss
<i>Dicranum scoparium</i>	broom moss
<i>Dicranum sp.</i>	heron's bill moss
<i>Digitalis purpurea</i>	common foxglove
<i>Distichlis sp.</i>	saltgrass
<i>Dryopteris expansa</i>	spiny wood fern
<i>Elymus glaucus</i>	blue wildrye
<i>Epilobium angustifolium</i>	fireweed
<i>Epilobium ciliolatum</i>	purple-leaved willowherb
<i>Epilobium sp.</i>	willowherb
<i>Equisetum arvense</i>	common horsetail
<i>Equisetum hyemale</i>	scouring-rush
<i>Equisetum pretense</i>	meadow horsetail
<i>Equisetum sylvaticum</i>	wood horsetail
<i>Erythronium oreganum</i>	white fawn lily
<i>Eurhynchium oreganum</i>	Oregon beaked moss
<i>Eurhynchium praelonga</i>	Oregon slender moss
<i>Festuca occidentalis</i>	western fescue
<i>Festuca rubra</i>	red fescue
<i>Festuca sp.</i>	fescue
<i>Festuca subulata</i>	bearded fescue
<i>Fragaria chiloensis</i>	beach strawberry
<i>Fragaria sp.</i>	strawberry
<i>Fragaria vesca</i>	wood starwberry
<i>Fragaria virginiana</i>	wild strawberry
<i>Fritillaria lanceolata</i>	chocolate lily
<i>Galium aparine</i>	cleavers
<i>Galium palustre</i>	marsh bedstraw
<i>Galium trifidum</i>	small bedstraw
<i>Galium triflorum</i>	sweet-scented bedstraw
<i>Gaultheria shallon</i>	salal
<i>Gentiana sp.</i>	gentian



<i>Geranium bicknellii</i>	Bicknell's geranium
<i>Geum macrophyllum</i>	big-leaved avens
<i>Gnaphalium microcephalum</i>	slender cudweed
<i>Gnaphalium</i> sp.	cudweed
<i>Goodyera oblongifolia</i>	rattlesnake-plantain
<i>Geranium molle</i>	dove-foot geranium
<i>Gratiola ebracteata</i>	bractless hedge-hyssop
<i>Grindelia integrifolia</i>	entire-leaved gumweed
<i>Heracleum maximum</i>	cow-parsnip
<i>Heuchera micrantha</i>	small-flowered alumroot
<i>Hieracium albiflorum</i>	white-flowered hawkweed
<i>Holcus lanatus</i>	common velvet-grass
<i>Holodiscus discolor</i>	oceanspray
<i>Hylocomium splendens</i>	step moss
<i>Hypericum</i> sp.	St. John's wort
<i>Hypochaeris glabra</i>	smooth cat's ear
<i>Hypochaeris radicata</i>	hairy cat's-ear
<i>Ilex aquifolium</i>	English holly
<i>Juncus articulatus</i>	jointed rush
<i>Juncus balticus</i>	Baltic rush
<i>Juncus covillei</i>	Coville's rush
<i>Juncus effuses</i>	common rush
<i>Juncus</i> sp.	rush
<i>Juniperus communis</i>	common juniper
<i>Juniperus scopulorum</i>	Rocky Mountain juniper
<i>Kindbergia oregano</i>	Oregon beaked moss
<i>Koeleria macrantha</i>	junegrass
<i>Lactuca muralis</i>	wall lettuce
<i>Leucolepis acanthoneuron</i>	palm tree moss
<i>Lilaeopsis occidentalis</i>	western lilaeopsis
<i>Lilium columbianum</i>	Columbia lily
<i>Linnaea borealis</i>	twinflower
<i>Listera caurina</i>	northwestern twayblade
<i>Lithophragma parviflora</i>	woodland star
<i>Lonicera ciliosa</i>	western trumpet honeysuckle
<i>Lonicera hispidula</i>	hairy honeysuckle
<i>Lonicera</i> sp.	honeysuckle
<i>Lotus micranthus</i>	small-flowered birds-foot trefoil
<i>Luzula multiflora</i>	many-flowered woodrush
<i>Luzula parviflora</i>	small-flowered woodrush
<i>Lycopus uniflorus</i>	northern water-horehound
<i>Lysichiton americanus</i>	skunk cabbage
<i>Madia</i> sp.	tarweed
<i>Mahonia aquifolium</i>	tall Oregon-grape
<i>Mahonia nervosa</i>	dull Oregon-grape
<i>Maianthemum stellatum</i>	star-flowered false Solomon's-seal
<i>Malus fusca</i>	Pacific crab apple
<i>Melica subulata</i>	Alaska onion-grass
<i>Mentha arvensis</i>	field mint
<i>Mimulus alsinoides</i>	chickweed monkeyflower
<i>Mimulus guttatus</i>	common monkeyflower



<i>Moehringia macrophylla</i>	big-leaved sandwort
<i>Monotropa uniflora</i>	Indian pipe
<i>Montia parvifolia</i>	small-leaved montia
<i>Myosotis laxa</i>	small-flowered forget-me-not
<i>Nemophila parviflora</i>	small-flowered nemophila
<i>Nuphar lutea</i>	yellow pond-lily
<i>Oenanthe sarmentosa</i>	Pacific water-parsley
<i>Oplopanax horridus</i>	devil's club
<i>Opuntia fragilis</i>	prickly-pear cactus
<i>Orthocarpus pusillus</i>	tiny owl-clover
<i>Osmorhiza chilensis</i>	sweet cicely
<i>Pachystima myrsinites</i>	false-box
<i>Panicum occidentale</i>	western witchgrass
<i>Peltigera</i> sp.	pelt lichen
<i>Pentagramma triangularis</i>	golden-back fern
<i>Petasites frigidus</i>	palmate coltsfoot
<i>Philonotis Fontana</i>	spring moss
<i>Physocarpus capitatus</i>	Pacific ninebark
<i>Picea sitchensis</i>	Sitka spruce
<i>Pinus monticola</i>	western white pine
<i>Pinus contorta</i> var. <i>contorta</i>	shore pine
<i>Plagiomnium insigne</i>	coastal leafy moss
<i>Plagiomnium</i> sp.	leafy moss
<i>Plagiothecium undulatum</i>	flat-moss
<i>Plantago elongata</i>	dwarf plantain
<i>Plantago lanceolata</i>	ribwort plantain
<i>Plantago major</i>	common plantain
<i>Plantago maritima</i>	seashore plantain
<i>Platanthera orbiculata</i>	large round-leaved rein orchid
<i>Platanthera unalascensis</i>	Alaska rein orchid
<i>Plectritis congesta</i>	sea blush
<i>Pleurozium schreberi</i>	red-stemmed feathermoss
<i>Poa annua</i>	annual bluegrass
<i>Poa compressa</i>	Canada bluegrass
<i>Poa pratensis</i>	Kentucky bluegrass
<i>Polypodium glycyrrhiza</i>	licorice fern
<i>Polystichum munitum</i>	sword fern
<i>Polytrichum juniperinum</i>	juniper haircap moss
<i>Polytrichum piliferum</i>	awned haircap moss
<i>Polytrichum</i> sp.	haircap moss
<i>Polytrichum strictum</i>	bog haircap moss
<i>Populus tremuloides</i>	trembling aspen
<i>Potentilla anserina</i> ssp. <i>pacifica</i>	Pacific silverweed
<i>Prunella vulgaris</i>	self-heal
<i>Prunus emarginata</i>	bitter cherry
<i>Pseudotsuga menziesii</i>	Douglas-fir
<i>Pteridium aquilinum</i>	bracken fern
<i>Racomitrium canescens</i>	grey rock moss
<i>Racomitrium lanuginosum</i>	hoary rock moss
<i>Ranunculus acris</i>	meadow spearwort
<i>Ranunculus repens</i>	creeping buttercup



<i>Ranunculus</i> sp.	buttercup
<i>Ranunculus uncinatus</i>	small-flowered buttercup
<i>Rhamnus purshiana</i>	cascara
<i>Rhizomnium</i> sp.	leafy moss
<i>Rhytidiadelphus loreus</i>	lanky moss
<i>Rhytidiadelphus triquetrus</i>	electrified cat's-tail moss
<i>Ribes lacustre</i>	black gooseberry
<i>Ribes sanguineum</i>	red flowering currant
<i>Rosa gymnocarpa</i>	baldhip rose
<i>Rosa nutkana</i>	Nootka rose
<i>Rubus laciniatus</i>	evergreen blackberry
<i>Rubus parviflorus</i>	thimbleberry
<i>Rubus discolor</i>	Himalayan blackberry
<i>Rubus pedatus</i>	five-leaved bramble
<i>Rubus spectabilis</i>	salmonberry
<i>Rubus ursinus</i>	trailing blackberry
<i>Rumex acetosella</i>	sheep sorrel
<i>Rumex obtusifolius</i>	dock
<i>Sagina procumbens</i>	procumbent pearlwort
<i>Salix lucida</i>	Pacific willow
<i>Salix sitchensis</i>	Sitka willow
<i>Salix</i> sp.	willow
<i>Sambucus racemosa</i>	red elderberry
<i>Sanguisorba canadensis</i>	Sitka burnet
<i>Sanicula crassicaulis</i>	Pacific sanicle
<i>Satureja douglasii</i>	yerba buena
<i>Saxifraga ferruginea</i>	Alaska saxifrage
<i>Scapania bolanderi</i>	yellow-ladle liverwort
<i>Schoenoplectus acutus</i>	hard-stemmed bulrush
<i>Sedum oreganum</i>	Oregon stonecrop
<i>Sedum spathulifolium</i>	broad-leaved stonecrop
<i>Selaginella wallacei</i>	Wallace's selaginella
<i>Senecio sylvaticus</i>	wood groundsel
<i>Sherardia arvensis</i>	spurwort
<i>Sonchus arvensis</i>	field sowthistle
<i>Sonchus asper</i>	prickly sow-thistle
<i>Sphagnum girgensohnii</i>	common green peat-moss
<i>Sphagnum</i> sp.	peat-moss
<i>Spiraea douglasii</i>	hardhack
<i>Stellaria crispa</i>	crisp starwort
<i>Stellaria media</i>	chickweed
<i>Stellaria</i> sp.	starwort
<i>Symphoricarpos albus</i>	common snowberry
<i>Symphoricarpos hesperius</i>	trailing snowberry
<i>Taraxacum officinale</i>	common dandelion
<i>Taxus brevifolia</i>	western yew
<i>Tellima grandiflora</i>	fringe-cup
<i>Thuja plicata</i>	western redcedar
<i>Tiarella trifoliata</i>	three-leaved foamflower
<i>Timmia australica</i>	false-polytrichum
<i>Tolmiea menziesii</i>	piggy-back plant



<i>Trientalis latifolia</i>	broad-leaved starflower
<i>Trifolium microcephalum</i>	woolly clover
<i>Trifolium oliganthum</i>	few-flowered clover
<i>Trifolium repens</i>	white clover
<i>Trifolium variegatum</i>	white-tip clover
<i>Tsuga heterophylla</i>	western hemlock
<i>Typha latifolia</i>	cat-tail
<i>Ulex europaeus</i>	gorse
<i>Urtica dioica</i>	stinging nettle
<i>Vaccinium ovalifolium</i>	oval-leaved blueberry
<i>Vaccinium ovatum</i>	evergreen huckleberry
<i>Vaccinium parvifolium</i>	red huckleberry
<i>Veronica arvensis</i>	wall speedwell
<i>Veronica officinalis</i>	common speedwell
<i>Viburnum edule</i>	highbush-cranberry
<i>Vicia americana</i>	American vetch
<i>Vicia gigantea</i>	giant vetch
<i>Vicia</i> sp.	vetch
<i>Xanthoparmelia cumberlandia</i>	questionable rockfrog
<i>Zygadenus venenosus</i>	death camas

